



22101825290





Digitized by the Internet Archive
in 2015

<https://archive.org/details/b20411479>

SURGICAL EMERGENCIES

By the same Author.

INJURIES AND DISEASES OF THE KNEE-JOINT,
and their Treatment by Amputation and Excision Contrasted.
Jacksonian Prize Essay of the Royal College of Surgeons.
With Wood Engravings. 8vo. 9s.

PRELIMINARY MEDICAL EDUCATION AT PROVINCIAL HOSPITALS. Sixpence.

SURGICAL EMERGENCIES

TOGETHER WITH THE

EMERGENCIES ATTENDANT ON PARTURITION
AND THE TREATMENT OF POISONING

A MANUAL

FOR THE USE OF GENERAL PRACTITIONERS

BY

WILLIAM PAUL SWAIN, F.R.C.S.

SURGEON TO THE ROYAL ALBERT HOSPITAL, DEVONPORT; HONORARY FELLOW
OF KING'S COLLEGE

THIRD EDITION



LONDON

J. & A. CHURCHILL, NEW BURLINGTON STREET

1880

19530

30049182

M19520

WELLCOME INSTITUTE LIBRARY	
Coll.	WOMOmec
Call	
No.	W0700
	1880
	S97s

TO
CHRISTOPHER HEATH, Esq., F.R.C.S.,
SURGEON TO UNIVERSITY COLLEGE HOSPITAL,
ETC. ETC.

This Book is Dedicated

AS A TRIBUTE OF SINCERE PERSONAL REGARD,
AND IN REMEMBRANCE OF MANY KINDNESSES RECEIVED,

BY
THE AUTHOR.



PREFACE TO THE THIRD EDITION.

IN preparing a Third Edition of this Manual, it has been my endeavour to embrace within its pages all that is most recent in the Surgery of Emergencies. At the same time, I have removed all that my own experience or that of others has found to be lacking in practical use. Many new woodcuts have been added, notably those illustrating the use of Esmarch's triangular cloth. For the description of its use I am indebted to Dr. Thomas Guy, who has translated it from the German. The chapter on Antiseptics has again been kindly revised by Dr. Bishop, and may therefore be taken to represent the latest improvements in Antiseptic Treatment.

WILLIAM PAUL SWAIN, F.R.C.S.

DEVONPORT, *February* 1880.

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY

JOHN BURNET

OF

THE UNIVERSITY OF OXFORD

IN TWO VOLUMES

LONDON

Printed by J. Streater, at the Sign of the Gun, in St. Dunstons Church-yard

1704

Vol. I.

THE HISTORY OF THE

REIGN OF

CHARLES THE FIRST

BY

JOHN BURNET

OF

PREFACE TO THE FIRST EDITION.

THIS Manual pretends to be little more than a compilation from the best and most recent works on Surgery. It is intended to supply what the Author believes is a wide-spread want—viz. a small book containing directions for the immediate treatment of all those various emergencies with which the General Practitioner may be called upon to deal at any moment. The aim has been throughout to condense as much as possible, without sacrificing precision and clearness in details. A chapter on the various emergencies attending Parturition, and also one on the treatment of Poisoning, have been introduced as affording information upon conditions particularly liable to arise at any moment, and requiring precise and immediate treatment. The chapter on Antiseptic Treatment has been kindly written by Dr. Bishop, at the request of Professor Lister, and embodies the most recent and exact directions for the effectual carrying out of this method.

I have sincerely to thank the various gentlemen who have permitted me to use their woodcuts for the purposes of this Manual, particularly Sir William Fergusson, Mr. Bryant, and Mr. Christopher Heath. I must also acknowledge, with thanks, the kind assistance rendered to me by Mr. G. Lawson in the compilation of the chapter on "Injuries to the Eye;" and by Dr. Alfred Meadows, for similar help in the chapter on "Emergencies connected with Parturition."

WILLIAM PAUL SWAIN.

DEVONPORT, *May* 1874.

CONTENTS.



CHAPTER I.

INJURIES TO THE HEAD.

Wounds of the Scalp—Concussion and Compression—Fractures of the Skull—Trephining—Fractures of the Base—Injuries to Cerebral Nerves—Injuries to the Face—Fracture of the Nasal Bones—Foreign Bodies in the Ear and Nostril—Epistaxis—Fractures and Dislocations of the Jaw—Injuries to the Spine
Pages 1-18

CHAPTER II.

INJURIES TO THE EYE.

Injuries and Acute Diseases of the Conjunctiva, Cornea, and Sclerotic—The Iris—The Lens—The Choroid—Injuries from Chemicals—Acute Glaucoma—Foreign Bodies within the Eyeball—Excision of the Eyeball—Artificial Eyes—Injuries to the Orbit—Injuries to Optic Nerve external to Eyeball . . . 19-31

CHAPTER III.

INJURIES TO THE MOUTH, PHARYNX, ŒSOPHAGUS, AND LARYNX.

Wounds of Tongue—Foreign Bodies in Pharynx and Œsophagus—Foreign Bodies in Windpipe—Scald of Larynx—Laryngitis—Laryngotomy—Laryngo-tracheotomy—Tracheotomy—Cut-throat—Stabs in the Throat—Ligature of Common Carotid—Drowning—Hanging—Respiration of Noxious Gases—Artificial Respiration 32-45

CHAPTER IV.

THE CHEST.

Fracture of the Ribs—Wounds of the Chest—Contusion of the Lung—Hernia of the Lung—Wounds of the Lung—Emphysema—Pneumothorax—Hæmothorax—Paracentesis Thoracis—Wounds of the Heart—Gunshot Wounds of the Chest

Pages 46–50

CHAPTER V.

THE UPPER EXTREMITY.

Wounds of Wrist-joint—Wounds of Palmar Arch—Needle in the Hand—Fracture and Dislocation of the Clavicle; of the Scapula; of the Humerus—Compound Injuries to the Shoulder-joint—Resection of Shoulder-joint—Amputation at Shoulder-joint—Fractures and Dislocations of the Radius; the Ulna; the Bones of the Hand—Compound Injuries to the Elbow-joint—Excision of Elbow—Compound Injuries to Wrist-joint—Amputation of the Thumb and Fingers 51–82

CHAPTER VI.

THE ABDOMEN.

Rupture of Muscles—Rupture of the Peritoneum—Injury to the Viscera—Non-penetrating Wounds—Penetrating Wounds—Protruded Viscera, Constriction, Distension with Flatus, Gangrene, Wounds—Protruded Omentum—Sutures in Wounded Parietes—Foreign Bodies—Obstructed, Inflamed, and Strangulated Hernia—Umbilical, Inguinal, Femoral, and Obturator Hernia 83–94

CHAPTER VII.

THE PELVIS.

Fractures and Dislocations—Injuries to the Bladder—Foreign Bodies in the Urethra and Bladder—Dilating the Female Urethra—Injuries to the Urethra—Retention of Urine—Use of Catheter—Puncture of the Bladder per Rectum; by Aspiration; above Pubes—Cock's Operation for Perineal Section—Retention from Enlarged Prostate—Extravasation of Urine—Injuries to the Penis—Paraphimosis—Phimosis—Injuries to the Scrotum; Testicle—Rupture of Vas Deferens—Injuries to the Pregnant Uterus; to the Vagina—Foreign Bodies in the Rectum—Hæmorrhage from the Rectum 95–110

CHAPTER VIII.

THE LOWER EXTREMITY.

Contusions—Rupture of Muscles—Sprains—Wounds of the Knee-joint—Ligature of Superficial Femoral ; of Posterior Tibial ; of Anterior Tibial—Fractures and Dislocations of the Femur—Compound Injuries to the Shaft of the Femur and Knee-joint—Excision of Knee-joint—Amputation of Thigh—Fractures and Dislocations of Patella ; of Tibia ; of Fibula ; of both Bones—Compound Comminuted Fractures—Compound Dislocation of Ankle-joint—Fractures and Dislocations of Tarsal Bones ; of Metatarsal ; of Phalanges—Amputation of Leg—Chopart's Amputation—Hey's Amputation—Amputation of Toes
Pages 111–146

CHAPTER IX.

EMERGENCIES CONNECTED WITH PARTURITION.

Signs of Pregnancy—Accidental Hæmorrhage—Placenta Prævia—Retained Placenta—Flooding—Induction of Premature Labour—Cæsarean Section—Craniotomy—Cephalotripsy—Embryulcia—Decapitation—Displacements of the Gravid Uterus—Malpresentations—Turning—Prolapse of the Cord—Puerperal Convulsions—Rupture of the Uterus ; of the Bladder ; of the Perinæum—Inversion of the Uterus . . . 147–159

CHAPTER X.

POISONING.

Poisoning by Sulphuric Acid—Nitric Acid—Oxalic Acid—Salt of Sorrel—Hartshorn—Phosphorus—Arsenic—Scheele's Green—Orpiment—Sugar of Lead—Blue Vitriol—Tartar Emetic—Chloride of Zinc—Cantharides—Mussels—Opium—Godfrey's Cordial—Dalby's Carminative—Paregoric Elixir—Prussic Acid—Cyanide of Potassium—Essential Oil of Bitter Almonds—Chloroform—Tobacco—Strychnia—Hemlock—Water-Hemlock—Monkshood—Deadly Nightshade—Fungi—The Use of the Stomach-Pump . . . 160–170

CHAPTER XI.

ANTISEPTIC TREATMENT.

General Remarks—Carbolic Acid—Antiseptic Gauze—Gauze Bandage—Antiseptic Atmosphere—Steam Spray Apparatus—

Boric Acid—Salicylic Acid—Chloride of Zinc—Protective Oil-Silk—Drainage Tube—Catgut Ligature—Carbolized Silk Sutures—Abscesses—Precautions before and during an Operation—Dressing after Operations—Compound Fractures—Dressing in Compound Fractures—Operations on Parts affected with Sinuses—Abscess by the side of Rectum . Pages 171-184

CHAPTER XII.

APPARATUS AND DRESSINGS.

Extemporary Apparatus—Conveyance of Injured Persons—Esmarch's Cloth—Swing for Patients in Bed—Pasteboard Splints—Plaster-of-Paris—Gum, Chalk, and Starch—Glue—Gutta-Percha Splints—Extension by Weight—The Use of the Aspirator—Transfusion of Blood—Esmarch's Bloodless Operation—The Elastic Tourniquet—Hunter's Tourniquet—Method of Opening Deep-seated Abscesses—Counter-openings—Burns and Scalds
185-208

LIST OF ILLUSTRATIONS.


FIG.	PAGE
1. The operation of trephining (Bryant)	4
2. Elevator	5
3. Hey's saw	5
4. Needle-holder	6
5. Syphon-tube (Heath)	8
6. Ear-forceps	9
7. Plugging posterior nares (Fergusson)	9
8. Instrument for plugging posterior nares	10
9. Adam's forceps for fracture of the nose	11
10. „ steel screw compressor	12
11. „ ivory plugs	12
12. Interdental wire splint (Lyons)	15
13. Bandage for fracture of the lower jaw (Bryant)	15
14. Instrument for removing foreign bodies from the eye	21
15. Eye speculum (Lawson)	21
16. Broad needle	21
17. Fine needle	21
18. Liebreich's eye bandage (Lawson)	23
19. Scissors for extirpation of eyeball (Lawson)	29
20. Weiss' tongue depressor	33
21. Horse-hair probang (Bryant)	34
22. Money probang	34
23. Tracheotomy canula	38
24. Pilot trocar	38
25. Tracheotomy tube	39
26. Sylvester's method (Heath)	43
27. „ „ „ „	44
28. Strapping for fractured ribs	47
29. Treatment of fractured clavicle by strapping	53
30. „ „ „ „	53
31. „ „ „ „	54
32. Splint for fracture of the neck of humerus (Heath)	59
33. Method of treating fracture of the shaft (Bryant)	60

FIG.	PAGE
34. Aitkin's Crimean splints	60
35. Gutta-percha splint for fracture at elbow-joint	61
36. Sub-glenoid dislocation of humerus	62
37. Sub-coracoid dislocation of humerus	63
38. Reduction of dislocated humerus by manipulation (Bryant)	64
39. Reduction of dislocated humerus by manipulation (Bryant)	64
40. Reduction pads	65
41. Stromeyer's cushion for compound fracture of humerus (Porter)	67
42. Stromeyer's cushion for compound fracture of humerus (Porter)	67
43. Small saw for resection of elbow-joint	68
44. Method of treating fracture of the forearm (Heath)	71
45. Colles' fracture	72
46. Gordon's splint for Colles' fracture	73
47. " applied to the arm	73
48. Pistol-shaped splint	74
49. The clove-hitch (Fergusson)	78
50. Forceps for reducing dislocation of phalanges	78
51. Method of amputating phalanges of the thumb (Ferg.)	80
52. " the metacarpal bone "	81
53. " the forefinger "	81
54. " the ringfinger "	82
55. " the third phalanx of finger (Heath)	82
56. Urethral forceps	98
57. Thompson's probe-pointed catheter	100
58. Perineal catheter (Bryant)	103
59. Hutchinson's silver prostatic catheter	105
60. India-rubber catheter	106
61. Rod for introducing india-rubber catheter	106
62. Nozzle and stiletted plug for india-rubber catheter	106
63. Method of treating rupture of the tendo Achillis (Ferg.)	113
64. Fracture of the neck of the femur (Fergusson)	118
65. " " " "	118
66. Liston's long splint applied " "	122
67. Fractured leg put up in MacIntyre's splint (Bryant)	123
68. Dislocation of the femur on the dorsum ilii	125
69. " into the great sciatic notch	125
70. Method of reduction by manipulation (Bryant)	126
71. Dislocation of the femur into the foramen ovale	127
72. Method of reduction by manipulation (Bryant)	128
73. Dislocation of the femur on the pubis	129
74. Reduction of dislocation by extension (Fergusson)	130
75. " " " "	130
76. Treatment of fractured patella	135
77. Fracture of the fibula	136
78. Dupuytren's splint	137

LIST OF ILLUSTRATIONS.

xix

FIG.	PAGE
79. Fracture of tibia and fibula	137
80. Arnold's splint and cradle	139
81. McNalty's splint	139
82. " opened for use	140
83. " with leg fixed in it	140
84. Dislocation of tibia inwards	142
85. " forwards (Bryant)	142
86. " backwards (Fergusson)	142
87. Diagram showing various amputations of the foot (Bryant)	145
88. Perforator for craniotomy (Meadows)	153
89. Craniotomy forceps "	153
90. Cephalotribe "	154
91. Leaves of hemlock (Taylor)	166
92. Leaves of parsley "	166
93. <i>a</i> , Root of aconite ; <i>b</i> , root of horseradish	167
94. Stomach-pump	169
95. Tosswell's stomach-pump	170
96. Extemporized litter (Porter)	186
97. Esmarch's triangular cloth	187
98. " " " applied to the head	187
99. " " " sling for arm	188
100. " " " applied to shoulder	189
101. " " " " chest	190
102. " " " " hip	191
103. " " " " hand	191
104. " " " " foot	191
105. Swing for patients in bed	192
106. Coxeter's pliers for cutting starch bandage	195
107. Foot-piece for extension by weight	198
108. Plaster strip for ditto	198
109. " " with foot-piece fixed	198
110. " " applied to leg	199
111. Pneumatic aspirator	200
112. Apparatus for transfusion	202
113. Modification of Esmarch's bandage	204
114. Elastic tourniquet, method of fixing	205
115. " " applied to limb	205
116. Dr. Hunter's tourniquet, unfitted	206
117. " " buckled	206



SURGICAL EMERGENCIES.

CHAPTER I.

INJURIES TO THE HEAD.

Wounds of the Scalp—Concussion and Compression—Fractures of the Skull—Trephining—Fractures of the Base—Injuries to Cerebral Nerves—Injuries to the Face—Fracture of the Nasal Bones—Foreign Bodies in the Ear and Nostril—Epistaxis—Fractures and Dislocations of the Jaw—Injuries to the Spine

WOUNDS of the scalp are generally flap-wounds, and, however extensive, the flap should be thoroughly cleansed, replaced, and kept in position by silver-wire sutures and strapping. Care must be taken not to include any part of the occipito-frontalis muscle in the sutures. The same treatment will be pursued if the bones of the skull are exposed. As a rule, these wounds are not followed by erysipelas. Punctured wounds are sometimes followed by diffuse inflammation, and if suppuration takes place, very free incisions down to the bone must be made.

Wounds of
the Scalp.

Concussion of the brain, following a blow on the head, may be very slight and transitory, or its effects may be severe and lasting. In a severe case the principal signs are—loss of motion and consciousness, or if capable of being roused, the patient relapses into unconsciousness; the pulse is also feeble, the pupils contracted, and the sphincters relaxed. This condi-

Concussion

tion is terminated favourably by an attack of vomiting, with restoration of heat and recovery of the pulse. The treatment consists in restoring heat by wrapping the patient in hot blankets, applying hot-water bottles to the feet, and using friction to the surface. If the collapse is severe, an enema of beef-tea and brandy should be administered.

Compression.

Compression, as the immediate result of a blow, arises from the pressure of a depressed portion of bone, or from extravasated blood within the cranium. The symptoms are complete coma, stertorous breathing, dilated pupils, full, slow pulse, passage of fæces involuntarily, retention of urine, and a hot, moist skin. Should these symptoms supervene shortly after the accident, they are probably due to extravasated blood. The general treatment of compression should be to relieve the bowels by the administration of a drop or two of croton oil, mixed with mucilage, and the use of turpentine enemata. The head should be shaved and ice applied, and the urine drawn off night and morning. The rules for any operative procedure are given below.

Fractures of the vault.

Fractures of the skull may occur in the vault or at the base. In the vault, fractures are generally the result of direct violence. The external table alone may be driven in, especially over the frontal sinuses, or a very slight injury to the external table may be accompanied by extensive fracture of the inner table. Very extensive depressions of bone may exist, especially under the temporal muscles, without the possibility of discovery, whilst extravasations of blood under the scalp may be mistaken for depressed fractures. If there be a wound down to the bone, care must be taken not to confound the cranial sutures with fissured fracture. The distinguishing point is, that a fissure presents a red line along the course of the fracture.

Simple fractures with depression, and comminuted fractures unaccompanied with a wound of the scalp, without brain symptoms, are not to be interfered with. Neither should an operation be proposed if the symptoms are slight, as they may arise from mere concussion. If, on the other hand, the nature of the accident and the severity of the symptoms indicate severe compression and contusion of the brain, trephining for the local injury is worse than useless. In children, compound depressed fractures, if no brain symptoms are present, should be let alone.

When not to trephine.

In compound comminuted fractures, with depression, the bone should be at once elevated, without waiting for symptoms, and if needful the trephine should be used. Any spicula of bone penetrating the substance of the brain should be removed, if this can be accomplished without damage to the brain tissue. In injuries to the skull, arising from blows with sharp instruments, it is not uncommon to find the inner table extensively splintered. In the event of any bad symptoms threatening, the trephine should be used at the earliest moment. In fact, it is advised by some authors in all cases of punctured fracture to apply the trephine at once. An instrument with a large crown should be used, and a circle of bone containing the puncture should be removed. If the injury be such as to make it probable that the middle meningeal artery is wounded, the trephine may be used near the anterior inferior angle of the parietal bone, for the purpose of letting out the effused blood.

When to trephine.

The instruments required for the operation of trephining are—1, Scalpel; 2, Trephine; 3, Hey's Saw; 4, Elevator; 5, Bone forceps; 6, Probe, with flattened or pointed extremity.

How to trephine.

The patient's head having been shaved, make a free crucial incision down to the bone over the site of the fracture, and dissect back the flaps. Place the

central pin of the trephine, fixed as at A, on the edge of the sound bone, and cut into the outer table by a




FIG. 1.

semi-rotatory movement. The teeth of the trephine are so set as to cut from left to right. Pressure on the instrument should therefore only be exerted when it is rotated in that direction. When a sufficient groove is made in the bone, withdraw the pin of the trephine, as at B, and complete the section of the outer table: cut through the inner table with great caution. The approach to the inner table through the diploë is indicated by the detritus being soft and bloody, instead of being dry as when the outer table was being divided. Whilst the inner table is being divided the groove in the bone should constantly be examined with a probe, to ascertain that in no part has it been quite cut through. The bits of bone being removed, raise the depressed portion with the elevator (Fig. 2). It may be possible by the use of Hey's saw (Fig. 3), or a pair of bone forceps, to remove a projecting bit of sound bone, and thus get room for

the use of the elevator, without recourse to the trephine. If, after removal of the bone, considerable



FIG. 2.



FIG. 3.

hæmorrhage should ensue from a wounded meningeal artery, a bit of sponge should be tucked in under the bone, so as to exert pressure on the wounded artery. Bring the wound together without sutures, and apply water-dressing. If trephining for effused blood, and none be found between the bone and dura mater, but that membrane bulges into the wound, it is advisable to open it, as blood may be effused beneath it.

Fractures of the base are generally the result of indirect violence, except in the nostrils and orbit, through which foreign bodies are sometimes thrust into the brain. In fractures through the orbital plate of the frontal, blood appears *first* beneath the ocular conjunctiva, then it spreads to the eyelids, the lower one being generally the first affected. If the hæmorrhage be excessive, the eyeball may be protruded. Bleeding from the nose and mouth, *if continuous*, indicates fracture through the ethmoid, body of the sphenoid, or basilar process. Large and continuous bleeding from the ears, especially if the blood filling the meatus pulsates, points to fracture through the petrous portion of the temporal. So, too, the profuse discharge of cerebro-spinal fluid indicates rupture of

Fractures of,
the base.

Discharge of cerebro-spinal fluid from.

Injuries to cerebral nerves.

Injuries to the face.
Wounds.

Wounds of the eyelids.

the cerebral membranes. The fluid will run out of the ears in large quantities so as to saturate the pillow, and may even be collected in a cup. It is a clear limpid fluid, containing a large quantity of chloride of sodium and a little sugar. It is not coagulated by heat or nitric acid. In fractures of the base the following symptoms indicate injury to nerves:—Loss of smell and sight indicate injury to the 1st or 2d pair; ptosis and dilated pupil to the 3d; paralysis of the face and loss of sensation on the injured side, loss of taste and sensation on the same side of the tongue, and loss of sensation in the nostril, to the 5th; internal strabismus, to the 6th; loss of hearing and facial paralysis, to the 7th. The 8th and 9th pair are seldom injured; but when they are, aphonia, dysphagia, and dyspnoea are the prominent symptoms.

Wounds of the face should be brought together very accurately, if possible with plaster only, the surface being painted over with flexile collodion. Leslie's narrow "tape plaster" will be found very convenient for this purpose. The strapping should not be removed for some days, unless swelling and redness appear. When the wound cannot be kept in apposition with plaster alone, serrefines will be found useful, as avoiding the cicatrix left by a suture. For sewing wounds about the eyelids a very fine curved needle is the best, and the operation is much



FIG. 4.

facilitated by the use of a needle-holder such as that shown in Fig. 4. In wounds of the eyelids fine ophthalmic silk is the most convenient for use when the skin only is involved. If the tarsal cartilage is

cut through by a clean incision, a very fine needle should be inserted close to the tarsal edge, and the edges drawn together with a twisted suture of fine silk. If the cut through the cartilage is jagged, it is better to pare the edges with a small scalpel, and then bring them together as above. In the eyebrow care should be taken to preserve the line. In wounds of the ears and nose the greatest care should be taken to replace the parts and fix them with sutures. This should be done under the most unpromising circumstances. Wounds through the thickness of the lips should be brought together with harelip pins, care being taken to keep the exact red-line of the lips; in order to effect this the first pin introduced should be at the margin of the lips. When the lip is cut internally against the teeth, considerable hæmorrhage may occur, the blood being swallowed and finally vomited in large quantities. The bleeding will most likely be found to proceed from the coronary artery.

Wounds of
the lips.

Foreign bodies may be removed from the nose with ordinary polypus forceps, or by those figured below (Fig. 6). Or a current of water may be injected up the opposite nostril, when if the patient is directed to keep the mouth open, it will return behind the foreign body and eject it. For this and other like purposes the syphon-tube (Fig. 5) will be found very useful, the force of the current being regulated by the height at which the jug containing the water is placed. The perforated weight, *a*, is placed at *m* in the bottom of the jug. The india-rubber tube is then curled up under the water, and left for a couple of minutes. Then raise the jug to the required height, and pinching the tube at *c* between the forefinger and thumb, draw it out over the brim of the jug to *c'*, when, being converted into a syphon, water will run continuously through the nozzle *v*. For injecting

Foreign
bodies in
nose and
ears.

the nostrils salt and water should always be used, water alone being very painful.




FIG. 5.

For removing foreign bodies from the ear a current of water will very frequently suffice, care being taken to direct the stream along the roof of the canal ; but if this fails, then the forceps (Fig. 6) made by Mayer and Meltzer are the best I know. They combine very great fineness with wonderful power and strength. In young and timid children it is better to administer chloroform, if any difficulty occurs in removing the foreign body.

Epistaxis.

Epistaxis may, when slight, be arrested by raising the head, applying cold to the forehead and nape of

the neck, making the patient snuff up tannic acid, or by injecting a current of iced salt and water through




FIG. 6.

the nostril with the syphon-tube, as described above.




FIG. 7.

Plugging
posterior
nares.

These plans failing, pressure on the facial artery as it passes over the edge of the lower jaw will sometimes succeed. If it be needful to plug the posterior nares, the usual instrument recommended is Bellocq's canula (Fig. 7). By this a plug of lint or compressed sponge is drawn up into the posterior nares, through the mouth. The plug should be guided behind the palate with the finger, and should lie with its long diameter upwards and downwards. The ends of the string hanging from the nostrils are separated, and tied over a plug of lint placed in the anterior nares,

whilst one string should be brought out through the mouth, to assist in withdrawing the plug if needful. A flexible gum elastic catheter is a good substitute for the canula, the silk or whip-cord being passed down through the eye of the catheter. An ingenious instrument, made by Mayer and Meltzer, is here figured (Fig. 8). In its collapsed condition it is introduced along the floor of the nostril. The thin india-rubber bag may then be blown up, thus exercising pressure on the whole interior of the cavity; or it may be distended with iced water, a constant current

Mayer's
instrument.




FIG. 8.

being sent through it, passing in at the upper tube and out at the lower one. Before plugging, care should be taken to syringe out all clots, which are often the cause of the hæmorrhage being kept up.

In very extreme cases the subcutaneous injection of ergotine may be had recourse to. The solution of ergotine I have used for this and other purposes is that prepared by Messrs. Gale, of Bouverie Street, London. The usual dose by subcutaneous injection is five minims containing two grains of ergotine.

In fractures of the nasal bones with displacement, the bones should be replaced as accurately as possible, by pressure from within the nostril. Should emphysema appear about the root of the nose, and below the eyes, the skin over the parts affected should be painted with collodion. A female catheter is the instrument usually recommended to effect the replacement of the bones, but its point is too large, especially when the parts are swollen. The end of a small sound will be found better adapted. No plugging of the nostrils is of any use. In displacement of the septum, a plug in both nostrils with a well-fitting gutta-percha shield on the nose, outside, will be found of advantage.

In cases where there is much displacement of the septum upon the subsidence of the swelling, Mr. Adams has devised the following plan of treatment:—

Fractures of
the nasal
bones.




FIG. 9.

The patient being placed under the influence of chloroform, a pair of strong forceps (Fig. 9) are used

to straighten the bent cartilaginous septum. The same forceps, with the blades closed, may be forced up each nostril under the lower portion of the nasal bones, should they be displaced, and the bones raised from within, whilst lateral pressure is made at the same time externally with the thumb.




FIG. 10.

In order to support the septum in its place, a steel screw compressor (Fig. 10) is used, one blade being introduced into each nostril, and the screw sufficiently tightened to hold it in place. This may be worn for a few days, after which its place may be supplied by two ivory plugs (Fig. 11), one placed in each nostril, which can be removed and replaced by the patient himself.




FIG. 11.

Fractures
of the
upper jaw.

The superior maxillæ are subject to various injuries, but there is seldom room for more treatment than that afforded by evaporating lotions and ordinary

retentive bandages. The two bones are sometimes separated from one another, but reunion takes place without much interference on the part of the surgeon. It is possible that the central incisors with the alveolar may be pushed by direct violence up into the nostril. In such a case the fragment will be felt in the nostril and can be easily pushed down into its place, and retained there by wiring it to the adjacent teeth, at the same time placing a firm plug in the nostrils. When the fragments are much displaced, an attempt should be made to place them in position by running the fingers along the alveoli, and restoring any irregularity. This may be facilitated by introducing a pair of dressing forceps into the nostril. The teeth may then be wired together, or a gutta-percha shield may be moulded round the alveolar process, care being taken not to let the molars of the lower jaw become imbedded in the gutta-percha whilst it is soft. If there is much comminution, the greatest care should be taken not to remove any fragments. Hæmorrhage from the branches of the internal maxillary is sometimes very excessive. It must be controlled by the application of ligatures, if possible, the actual cautery, ice, or the perchloride of iron. Fracture of the nasal process is treated in the same way as fracture of the nasal bones. A very serious deformity is caused by depression of the malar bone and fracture of the antrum. The deformity may be remedied by making an opening into the antrum through the alveolar process, and then pressing up the malar bone with a sound, or, as recommended by Hamilton, making a small incision through the cheek at the anterior margin of the masseter muscle, and then introducing a hook under the zygoma, and thus lifting it up. Before having recourse to these severe measures, an attempt should be made to replace the bone by pressing with the thumb, through the mouth, against

Fracture of
the antrum
and depres-
sion of
malar bone.

Fractures of,
the zygoma.

the zygoma. Fractures of the zygoma are only of consequence when the fragments, projecting inwards, enter the tendon of the temporal muscle and impede the movements of the lower jaw. Reduction should then be attempted from within the mouth. In fractures of the alveolar process the teeth in the broken fragment should be secured to a firm tooth, if there is one, by silver wire; or a mould of gutta-percha may be fitted to the fragments. On no account should the fractured portion be removed.

Fractures of
the lower
jaw.

Fractures of the lower jaw are generally the result of much violence. Fractures are more frequent in the body of the bone, and they generally happen at or near the mental foramen. If it be a single fracture it is easily detected by pressing the posterior fragment inwards, when mobility and crepitus will be discovered. If the fracture—as is not unfrequently the case—be double, then it is discovered by taking the central fragment between the finger and thumb and moving it. Fractures at the angle and ascending ramus present the same features as those of the body, only there is less displacement and mobility. Fractures of the coronoid are rarely met with, but a severe blow on the chin will sometimes fracture the neck of one or both condyles. This may be discovered by the pain at the seat of fracture, on moving the jaw, and by crepitus, which the patient can himself generally feel and hear. The chin is also turned towards the affected side, which will distinguish it from dislocation. In the treatment of fracture of the jaw, all teeth that are very much loosened from their sockets, in the immediate neighbourhood of the fracture, should be removed. It should also be ascertained that no tooth, or portion of a tooth, has fallen between the fragments. If there is much displacement, and the teeth contiguous to the fracture are sound, they should be tied together either with silver

Treatment
of fractures
of the
lower jaw.

wire or thick silk; but this should be avoided if possible; and if a ligature is used it should be tied as high up as possible, to prevent the gum from being




FIG. 12.

cut and irritated. In severe cases, especially when the fracture is compound, an interdental splint of iron wire, invented by Mr. Hammond, may be used. It is made of a piece of wire bent so as to embrace the whole of the teeth and connected together between the teeth by bits of thinner wire (Fig. 12). The simplest and most efficacious form of bandage is the one figured (Fig. 13). It is made with a piece of bandage or muslin, a yard long and four inches wide. A slit four inches long is made longi-

Interdental
splint.



FIG. 13.

tudinally in the centre, and the ends of the bandage are torn down to within three or four inches of this slit. The chin is then placed in the slit, the anterior tails are tied behind the nape of the neck, the posterior ones over the top of the head, and the two connected together.

Dislocations
of the
lower jaw.

The lower jaw may be dislocated on one or both sides. In both the displacement is forwards. The double dislocation is the most frequent. The condyles are drawn upwards and forward by the action of the pterygoid and temporal muscles, and the coronoid processes rest on the back of the superior maxillæ. The symptoms are—widely open mouth, and immobility of the jaw, with considerable pain, loss of power of deglutition and articulation. Immediately in front of the ears a distinct hollow is perceptible, and the condyle is felt in front of this. When one side only is dislocated, the symptoms are not so prominent, and it may be overlooked. The chief points are that the chin is usually directed to the sound side, and the hollow in front of the ears is found on the injured side. The dislocation may be reduced in two ways. Place the patient on the floor with the head between your knees. Then put two pieces of cork as far back as possible between the molars on either side. Press the point of the chin steadily and directly upwards. Or, standing in front of the patient, place your two thumbs well guarded with a towel between the last molars. Grasp the jaw firmly with the hands. Press downwards and backwards with the thumbs, and at the same time elevate the chin.

Reduction.

Injuries to
the spine.
Sprains.

Sprains of the spine are found principally in the cervical and lumbar regions. In the neck they are accompanied with intense pain, and partial loss of power in the arms and legs, with numbness and pricking throughout the body. In the lumbar region there is general pain across the loins, with consider-

able swelling, and frequently hæmaturia. But the line of spinous processes is found level, and the pain on pressure cannot be limited to any particular spot. Perfect rest and hot fomentations are alone required.

Concussion of the spinal cord may arise from direct or indirect violence. Injury from direct violence in the cervical region may cause instant death ; or there may be a sense of suffocation, irregular action of the heart, and vomiting, from injury to the vagus. If the spinal accessory is involved, there may be spasmodic action of the trapezius and sterno-mastoid. Hiccup and a sense of constriction round the body indicate injury to the phrenic. Paralysis of one or both arms may be induced. Concussion.

In the dorsal and lumbar regions concussion may produce complete paraplegia. There is, however, no priapism, as in laceration of the cord. Concussion from indirect violence frequently takes place in the severe shock sustained in railway accidents, and it is to be remarked that the remote injury produced is quite out of proportion to the apparent slowness of the accident. It is also to be remarked that the symptoms of spinal concussion seldom appear when serious injury has been inflicted on other parts of the body, unless the spine itself has been directly injured. The principal treatment consists in perfect and prolonged rest, the patient lying as much as possible on the face. The administration of chloral and bromide of potassium will be found of service. Dry cupping and the application of ice-bags will also give relief.

Fractures of the spine above the level of the fourth cervical vertebra cause, as a rule, instant death. In the cervico-dorsal region fracture frequently takes place. All below the point of fracture is paralysed, but there is sensation as low as the nipples. Breathing is entirely carried on by the diaphragm and the serratus magnus. In the dorsal region, between the Fractures and dislocations.

fourth and the tenth vertebræ, there is generally complete paralysis of the lower extremities, of the abdominal muscles, and of the bladder and rectum. Respiration is imperfectly performed by the diaphragm and upper part of the thorax.

In the lumbar region there may be an absence of paralysis, or if it is present in the erect position, when the patient is placed on his back the fractured parts are replaced, and he recovers the use of his lower extremities.

Dislocation of the axis from the atlas sometimes occurs, and is followed by instant death. Dislocation of any one of the five lower cervical vertebræ may take place from forcible flexion and rotation combined. Dislocation of the transverse processes of the cervical vertebræ sometimes occurs, the head being fixed immovably and turned to the opposite side to that on which the injury has taken place. Reduction may be effected by placing the two knees on the patient's shoulders, extending the head, and forcibly turning it into position.

Treatment.

In such injuries all motion of the spine should be guarded against during the removal of the patient, by placing him on a door, or board (see Chap. XII.) If the fracture be cervical, the head should be surrounded with a sack filled with sand, to prevent all motion, and the clothes should be cut off, to avoid disturbance. Bleeding should not be resorted to. The patient should at once be placed on a water-bed, a catheter introduced at the earliest moment, and the urine drawn off.

CHAPTER II.

INJURIES TO THE EYE.

Injuries and Acute Diseases of the Conjunctiva, Cornea, and Sclerotic—The Iris—The Lens—The Choroid—Injuries from Chemicals—Acute Glaucoma—Foreign Bodies within the Eyeball—Excision of the Eyeball—Injuries to the Orbit—Injuries to Optic Nerve external to the Eyeball.

SUDDEN effusions of blood sometimes take place beneath the conjunctiva, following a sneeze, or some other violent exertion. They produce considerable alarm, but call for no treatment. Lacerations of the conjunctiva produce much swelling, but it is seldom needful to do more than keep the eye closed, and apply some evaporating lotion. If a large flap is torn, it may be sometimes necessary to put in one or two sutures of fine ophthalmic silk. Very small curved needles are the best for the purpose, used with a needle-holder (Fig. 4, p. 6).

Injuries of the conjunctiva.
Effusion of blood.
Lacerations.

Catarrhal ophthalmia is an acute inflammation of the conjunctiva arising without apparent cause, or from exposure to cold. It is epidemic and very contagious. The characteristic symptoms are bright redness, large secretion of muco-pus, the affection of both eyes simultaneously, or one following the other rapidly. Treatment:—Bathe the eye every three hours with a lotion of alum (gr. vj ad ʒj). Drop in two or three drops of a solution of nitrate of silver (gr. j ad ʒj) night and morning. Keep the lids from sticking together by smearing a little vaseline

Acute diseases of the conjunctiva.

over the edges. Give a smart purge and then administer tonics.

Ophthalmia
neonatorum.

Ophthalmia neonatorum, or the purulent ophthalmia of new-born infants, commences from the second to the seventh day after birth. Both eyes are simultaneously affected. It commences with slight muco-purulent discharge, glueing the lids together. The discharge rapidly increases and changes to pure pus. The lids become red and swollen, and the cornea in some cases is rapidly involved. Treatment:—Wash away the discharge as soon as it collects, and syringe out the eyes with an alum lotion every half-hour. Twice a day drop in nitrate of silver drops (gr. ij ad ʒj). Above all things keep the child in a darkened room.

Gonorrhœal
ophthalmia.

Gonorrhœal ophthalmia is a specific disease induced by the contact of gonorrhœal matter. The symptoms are—profuse purulent discharge, great chemosis, swelling of the lids, which become red, tense, and closed; rapid involvement of the cornea. Treatment:—Full administration of stimulants, and liberal diet. A free purge, and then quinine, in 2 gr. doses every four hours, with opium, if required, to allay pain. Nitrate of silver (gr. x ad ʒj) applied on the everted lids with a camel's hair brush, and then washed off with a stream of water, once or twice daily. If the lids cannot be everted, the solution may be dropped in. Wash the eye every hour with an alum or zinc lotion; cold applications are generally the most grateful.

Foreign
bodies in the
cornea.

Method of
removal.

Very minute portions, generally of iron, become imbedded in the epithelium of the cornea. They cause great pain, lachrymation, and intolerance of light. To remove them, seat the patient in a chair, and, standing behind him, draw up the upper lid, and steady the globe of the eye with the fore and middle finger of one hand, whilst with the other pick

out the foreign body with a small spud or chisel (Fig. 14). This little instrument for the pocket



FIG. 14.

contains on one side a curette, which is very useful for everting the lids, and other purposes, whilst on the other side is the spud. Care must be taken, if the foreign body is deeply fixed, not to press it into the anterior chamber; and if there be danger of this, a broad needle must be passed into the anterior chamber from the corneal edge, and pressed against the cornea immediately behind the foreign body, which can then be picked out with another needle.




FIG. 15.

For this operation there will be required a spring speculum (Fig. 15), a broad needle (Fig. 16), a fine



FIG. 16.



FIG. 17.

needle (Fig. 17), and conjunctival forceps to steady the eye with. After this operation a little castor-oil dropped into the eye is very soothing. Atropine is

soluble in castor-oil, and is a useful addition when there is much pain and irritation.

Abrasions of
the cornea.

Abrasions of the cornea cause considerable pain, intolerance of light, and lachrymation. They appear as little facets where the cornea has been abraded. The application of castor-oil, and the closure of both eyes, affords very great relief.

Foreign
bodies
beneath the
eyelid.

Foreign bodies very frequently become lodged beneath the upper lid, and abrade the cornea. They cause much distress, and are frequently overlooked, the inflammation of the eye caused by their presence being put down to cold, etc. It is wise always to evert the upper lid when there is the least suspicion of a foreign body being present. To do this, seat the patient on a chair, and standing behind him, take hold firmly of the eyelashes, and by them lift the upper lid off the globe: then placing the curette (Fig. 14) parallel to the upper margin of the tarsal cartilage, tilt the lid upwards over the curette. Keep it in place with the fore and middle fingers, withdraw the curette, and with it remove any little foreign body you may find lodged on the conjunctiva of the lid.

Wounds of
the cornea
and
sclerotic.

Wounds of the cornea, if simple, heal rapidly, the eye being kept closed with a pad of cotton-wool and a bandage. Wounds accompanied with much contusion, especially those involving the sclerotic, are very dangerous, and perfect rest should be at once enjoined, both eyes being closed, and three or four leeches applied to the temples. In extensive wounds of the sclerotic, even when much vitreous has been lost, the wound should be closed by suture. A very fine suture of platinum wire is the best.

For closing the eye, in this and other cases requiring such treatment, "Liebreich's Eye Bandage" (Fig. 18) will be found the most useful. It consists of a knitted band, A, ten inches long and two and a half

inches broad. The tape, B, twelve inches long, passes over the top of the head, from ear to ear; the second




FIG. 18.

tape, C D, passing through a loop at the end of B, and tying with E at the side of the temples. The bandage is thus applied:—Make the patient close both eyes gently, and over each place a square of linen, and again on this a round pad of cotton-wool. The bandage being fitted to the head, is drawn across the eyes, and fastened on the temple, opposite to the injured eye.

Diffuse suppurative corneitis commences with pain and heat in the eye, congestion of the conjunctiva and sclerotic, photophobia, and lachrymation. The cornea becomes steamy, and pus is soon effused in its layers. If the disease advances, the deeper tissues become involved, and pus is effused into the anterior chamber, settling down into its lower part and constituting hypopion. The treatment consists of the application of warm belladonna fomentation (ext. belladon. $\overline{3}$ j, aq. ferv. Oj). The fomentation should be applied as hot as possible and changed frequently. If there is increased tension, iridectomy should be performed or paracentesis. The latter operation is performed as follows:—Introduce the wire speculum,

Acute diseases of the cornea.

Paracentesis of the cornea.

and seizing a fold of conjunctiva with the conjunctival forceps, perforate the cornea at its edge with a broad needle (Fig. 16), keeping the point well towards the posterior surface of the cornea. Then turn the needle on its edge, so as to stretch the wound and allow the aqueous to escape. When the iris approaches the cornea, as it will do on the escape of the aqueous, turn the needle again on the flat and withdraw it. Remove the speculum and close the eye with a fold of wet lint and a bandage.

Sloughing
ulcer of
cornea.

A sloughing ulcer of the cornea presents an excavated surface, with a whitish-yellow base, the cornea all round being hazy. This ulceration frequently runs on to perforation and prolapse of the iris. Paracentesis will frequently prevent this, and iridectomy is also highly recommended. The use of nitrate of silver is to be avoided.

Crescentic
ulceration
of cornea.

Crescentic ulceration appears exactly at the margin of the cornea, and follows the curve of the corneal edge. It is at first transparent, but afterwards becomes nebulous. Perforation of the cornea frequently takes place. This form of ulceration is very difficult to cure. The treatment consists of the application of belladonna fomentation, dropping in liq. atropiæ, and keeping both eyes closed. Sometimes nitrate of silver is of use. It should be applied with a camel's hair brush, moistened with water, and drawn several times over a stick of nitrate of silver. In both these forms of ulceration, liberal diet, stimulants, tonics, and sedatives should be administered.

Injury to
the iris.

Hæmor-
rhage into
anterior
chamber.

The iris is sometimes injured by sharp blows, as from a cork discharged from a soda-water bottle, etc. The immediate result may be hæmorrhage into the anterior chamber, from rupture of some of the vessels of the iris. As a rule the blood is rapidly absorbed, rest and an evaporating lotion being the treatment. Another effect of a blow may be the separation of the

iris from its ciliary border. This is accompanied by considerable hæmorrhage into the anterior chamber, but if uncomplicated by other damage, generally does well.

Detachment
of iris.

Wounds at the edge of the cornea are frequently accompanied with prolapse of the iris. Immediately after the accident, atropine drops should be instilled, and a firm compress applied. This should be removed two or three times a day, the eye bathed with warm water, and fresh drops put in.

Prolapse of
iris.

When the prolapse is small, and the wound in the cornea clean and not contused, the nodule of iris may be cut off with a pair of scissors. If the prolapse is large, and the corneal wound inclined to gape, the nodule should be frequently punctured with a needle. The general treatment should be—the application of two or three leeches to the lid of the injured eye, belladonna lotion, and the use of atropine drops (gr. j ad ʒj) twice a day.

Syphilitic iritis commences as the secondary rash is fading. Its peculiar characteristic is the deposit of nodules of lymph on the edge and surface of the iris. Photophobia and pain are neither well marked. Treatment:—Give mercury until the gums are tender, with iodide of potassium; drop in atropine drops (gr. ij ad ʒj) twice a day; or use belladonna lotion (ext. belladon. ʒij, aq. ad ʒviiij).

Acute dis-
eases of the
iris.
Syphilitic
iritis.

In rheumatic iritis the aqueous becomes yellow. The pain is very severe and neuralgic, with great photophobia. Treatment:—Iodide of potassium, in small doses, with bicarbonate of potash, or 2 gr. doses of quinine, every four hours. Slight mercurial inunction to the temples. Keep the pupil dilated with atropine.

Rheumatic
iritis.

Acute traumatic iritis may come on four or five days after an injury. The symptoms are oedema of the lids and chemosis of the conjunctiva, with rapid exudation

Traumatic
iritis.

of lymph on the surface of the iris and in the pupil. Treatment :—Leeches to the temple, atropine drops, belladonna lotion or fomentation, a purge to relieve the bowels, and opium to relieve pain. If suppuration should occur, with much distension, paracentesis of the cornea should be performed (see p. 23). In all these cases the use of a shade covering *both* eyes is of consequence. By far the best shade is one cut out of brown paper. It should reach from ear to ear, and should be from four to six inches deep. A strip of brown paper passing round the back of the head and pinned to the shade above either ear, will secure it in place.

Shading the eye.

Injury to the lens.

The lens is frequently wounded, in company with the cornea and iris. When this is the case the lens tissue rapidly swells and becomes opaque. The iris should be kept fully dilated with atropine to prevent the swollen lens pressing upon it. Should symptoms of "traumatic glaucoma" come on, indicated by increased pain, hardness of the globe, diminution of the anterior chamber, and pink tinge of the sclerotic, the lens should at once be removed. As it is generally soft, the suction operation will be applicable, or the entire lens may be removed through a small corneal wound, by the simple introduction of the curette, accompanied with slight pressure on the cornea.

Traumatic glaucoma.

If urgent symptoms, however, do not manifest themselves, atropine drops, belladonna lotion, and perfect rest by closing both eyes, should be the plan of treatment. It must be remembered that opacities of the lens sometimes come on after a blow on the eye, unaccompanied by any external wound.

Dislocation of the lens into the anterior chamber.

Dislocation of the lens into the anterior chamber may be the result of a blow. The lens appears "like a large drop of oil lying at the back of the cornea, the margin exhibiting a brilliant yellow reflex." The iris is pushed backwards, and widely dilated. Great

pain and inflammation attend this accident, as a rule. It is advisable at once to remove the lens, linear extraction being the operation most suitable. When the lens is dislocated into the vitreous, the symptoms are, a falling backwards and dilatation of the iris, which also becomes tremulous. If great pain and irritation come on, it is better to endeavour at once to remove the lens—the traction operation being in this case the best.

Dislocation
of the lens
into the
vitreous.

Partial dislocations of the lens are the result of severe blows, and generally terminate in cataract. There is also sometimes a glaucomatous state produced, which intermits, but is highly dangerous to sight, and the lens should be removed by the traction operation, with a large iridectomy. In all these cases the operation is sure to be accompanied with loss of vitreous.

Partial
dislocations.

A severe blow on the eye will frequently cause hæmorrhage to take place between the choroid and the retina. If the hæmorrhage is severe all useful vision is lost, but if slight, recovery may be looked for with a blind spot corresponding to the detached portion of retina. The treatment should consist of a couple of leeches to the temple, ice over the eye, atropine drops, and perfect rest to both eyes.

Injuries to
the choroid.

Injury to the eye by lime or mortar, if *seen immediately*, should be treated by washing out with a tepid solution of vinegar (3j ad aq. 3ij). The upper lid should be everted, and all portions remaining in the eye carefully removed; a drop of castor-oil may then be put in, and the eye closed with a pledget of wet lint.

Injuries to
the eye
from chemi-
cals, etc.:
lime.

When the eye is injured by a strong acid it should be syringed out with an alkaline solution (sodæ bicarb. gr. v, aq. 3j), the after-treatment being the same as that above described. If the lids are injured with the acid they should be dressed with the following liniment:—

Strong
acids.

Olei lini,

Liq. calcis, āā fl. oz. iv.

Cretæ preparatæ, oz. ij.

Lint dipped in this liniment should be applied over the eye, and a pad of cotton-wool placed over it with a bandage.

Gunpowder.

When the eye is injured by an explosion of gunpowder, all the loose powder should be carefully removed with a syringe and tepid water. The cornea should also be carefully searched, and all particles embedded in it removed with the spud. Castor-oil should then be dropped in, and belladonna lotion applied.

Acute
glaucoma.

Acute glaucoma is sudden in its attack. The eye presents all the appearance of acute inflammation. The anterior chamber is greatly diminished, so that the iris is almost in contact with the cornea. The pupil is dilated and very sluggish, if not entirely inactive. The patient sees rainbows round the candle or gas light. The field of vision is much diminished and sight greatly impaired. The tension of the globe is increased even to stony hardness. The pain is most intense, especially at the back of the head, and as it is frequently accompanied by sickness, the attack is too often mistaken for a simple bilious one. If the fundus can be seen with the ophthalmoscope, the optic entrance will be found cupped, the retinal arteries pulsating, and the veins tortuous and distended. Small bloodshots may also be seen scattered over the retina. A few hours will frequently accomplish the destruction of the eye. Treatment :—The immediate performance of an iridectomy gives the only chance of saving the eye.

Foreign
bodies
within the
eye.

The fragments caused by the explosion of percussion-caps, and small shot, very frequently penetrate the eyeball. If the foreign body can be seen, an attempt should be made to remove it. If its presence is only

suspected, great watchfulness must be exercised. The following symptoms indicate the presence of a foreign body:—

1. Continuation of primary inflammation.
2. Subacute choro-iritis, after the subsidence of the primary inflammation.
3. Non-union of the corneal wound.
4. Severe and continued pain in the eye.

The continuance of these symptoms should lead to excision of the eyeball. But if at the time of the accident sight is destroyed, and no doubt exists as to the presence of a foreign body within the eye, the eyeball should be at once extirpated.

The instruments required for this operation are—1, a spring speculum; 2, conjunctival forceps; 3, a blunt-pointed pair of scissors, curved on the flat (Fig. 19); and 4, a strabismus-hook. The patient being

Extirpation
of the
eyeball.



FIG. 19.

placed under the influence of chloroform, the spring speculum is introduced between the eyelids. A fold of conjunctiva is then nipped up with the forceps close to the corneal edge. One blade of the scissors being introduced into this opening, they are run round the edge of the cornea, snipping as they go, and thus dividing the conjunctiva completely. The forceps are then laid aside, and the recti muscles are taken up on the squint-hook in succession, and divided with the scissors close to the globe. By pressing the speculum backwards into the orbit the globe is now made to protrude from the socket. The remaining oblique muscles and the fascia are then divided. The closed scissors are next passed backwards along the inner side of the eyeball, and on being opened, are made to include the optic nerve, which is divided, and the

entire globe lifted from the orbit. Brisk hæmorrhage ensues from the central artery, which is easily controlled by a stream of cold water. If it continues to bleed, the best plan is to leave the speculum between the lids, and press a small dry sponge into the orbit, retaining it there firmly with a bandage for an hour or two.

Artificial
eye.

An artificial eye should not be introduced until six weeks or two months after extirpation. The following instructions are useful with regard to wearing the eye:—It should be taken out every night and replaced in the morning.

To Take the Eye Out.—The lower lid must be drawn downwards with the middle finger of the left hand; and then, with the right hand, the end of a small bodkin must be put beneath the lower edge of the artificial eye, which must be raised gently forwards over the lower eyelid, when it will readily drop out. At this time care must be taken that the eye does not fall on the ground, or other hard place, as it is very brittle and may readily be broken by a fall.

To Put the Eye In.—Place the left hand flat upon the forehead, with the fingers downwards, and with the two middle fingers raise the upper lid towards the eyebrow; and then with the right hand push the upper edge of the artificial eye beneath the upper eyelid, which may now be allowed to drop upon the eye. The eye must then be supported with the middle fingers of the left hand, whilst the lower eyelid is raised over its lower edge with the right hand.

Injuries to
the orbit.

Foreign bodies are sometimes impacted in the orbit. To remove them, the outer canthus should be slit with a pair of scissors or a scalpel. If the body be in the upper part of the orbit, the upper lid must be turned up, and the fold of mucous membrane reflected from the globe, divided. The foreign body being felt, may then be removed with a pair of forceps. If it is

in the lower part of the orbit, the lower lid must be drawn down, and the conjunctiva divided. Should the foreign body be small and metallic, anything like prolonged search is to be deprecated, as frequently little or no harm comes from its presence in the orbit, and considerable mischief may follow the attempt to remove it.

It sometimes happens in gunshot wounds that a shot will enter the orbit, and, passing outside the eyeball, will either divide or so injure the optic nerve as to entirely destroy vision. The accident is accompanied with much hæmorrhage into the orbit, which protrudes the eyeball. It is a point of considerable importance as to whether the shot has entered the eyeball or not ; as in the former case extirpation of the eyeball would be an undoubted necessity ; whilst in the latter no interference is needful. Ophthalmoscopic examination will clearly demonstrate the presence of a shot within the globe. If it is found not to be there, then the other alternative must be adopted, viz. that the shot has entered the orbit external to the globe, and, passing backwards, either divided or injured the optic nerve external to the globe.

Injury to
optic nerve
external to
eyeball.

CHAPTER III.

INJURIES TO THE MOUTH, PHARYNX, ŒSOPHAGUS, AND LARYNX.

Wounds of Tongue—Foreign Bodies in Pharynx and Œsophagus
—Foreign Bodies in Windpipe—Scald of Larynx—Laryngitis
—Laryngotomy — Laryngo-tracheotomy — Tracheotomy — Cut
Throat—Stabs in the Throat—Ligature of Common Carotid—
Drowning—Hanging—Respiration of Noxious Gases—Artificial
Respiration.

Wounds of
tongue.

WOUNDS of the tongue, if the parts gape, must be drawn together with fine silk. The suture should be placed deeply, with a curved needle set in a handle. In young children it is better to give chloroform, the mouth being opened with a screw gag, and the tongue drawn well forward by seizing it at the tip with a pair of hooked forceps. Ice, or the application of perchloride of iron, will generally stop all bleeding; but if it persists, a ligature may be passed with a curved needle round the bleeding point, enclosing the muscular tissue.

Foreign
bodies in
the pharynx.

Small sharp bodies, such as bristles, fish bones, or pins, are generally found sticking between the pillars of the fauces and the tonsils. The laryngoscopic mirror is a great aid in finding them; and a pair of fine urethral forceps are the best with which to remove them. Larger substances, such as pieces of meat, coins, or false teeth, are generally arrested at the junction of the pharynx with the Œsophagus, or between the base of the tongue and the epiglottis.

If the foreign body can be reached with the fingers, it may be hooked up, or it may be seized with a pair of curved forceps. It should never be pushed down. If suffocation is impending, laryngotomy should be immediately performed (see below), and then further attempts made to remove the obstruction. If the body is rough or angular, a blunt hook will often catch it. A common skewer, heated in the fire, and the point turned, has served for this purpose. A new tongue-depressor, made by the Messrs. Weiss (Fig. 20), seems a most useful instrument. The spatula is passed into the mouth, the handle being held beneath the chin. Thus no light is lost, and the patient can himself hold the instrument.




FIG. 20.

Foreign bodies stick in the oesophagus generally, either opposite the cricoid cartilage, or just above the diaphragm. If they are soft, and not likely to damage the stomach, they may be pushed on with an ordinary sponge probang, or stomach tube. Sharp bodies, like pins, should not be pushed on, but an endeavour should be made to catch them with the "horse-hair probang" (Fig. 21). This being pushed gently, unexpanded, beyond the point where the pin or bone is stuck, is expanded by pulling up the handle, and then withdrawn with a slight rotatory motion. If a coin is impacted, a "money probang" (Fig 22) may succeed in catching it. This may be easily im-

In the
oesophagus.

provided by turning a loop of stout wire sharply on itself.

In the
larynx.

In the
trachea.




FIG. 21.

A sudden inspiration, when the mouth is full, may draw a large piece of food into the larynx and cause instant death. Smaller bodies may be arrested in the cavity of the larynx, causing suffocation by spasm; or they may pass onwards into the trachea, the symptoms being intermittent cough, with pain at the point of impaction; or they may become




FIG. 22.

impacted, generally in the right bronchus, absence of respiration in the lung indicating this fact (see p. 40). To be sure that the impediment to respiration is not in the œsophagus, if possible make the patient drink some water, which will be swallowed with ease if the obstruction be in the windpipe. The sudden onset of the symptoms, the intermission of the paroxysms, and the dyspnœa during *expiration*, distinguish the symptoms due to the presence of a foreign body from croup or laryngitis.

Scald of
larynx.

In scalds of the larynx the respiration is affected two or three hours after the injury, the voice becoming hoarse, and the inspiration croupy. Sometimes

fatal spasm occurs. A hot sponge should be applied at once to the throat, warm vapour inhaled, and if swallowing be possible, two or three grains of calomel should be given every hour. The cedematous tissue about the fauces should be well scarified with an ordinary tenaculum, and leeches freely applied to the neck. Should the respiration become rapidly worse, tracheotomy must be performed (see below), and the earlier the better, before the lungs become involved. The same treatment is applicable to injury by strong acids. When taken suicidally, the windpipe is rarely injured, but when taken by mistake, the acid frequently finds its way into the larynx, in the efforts made to expel it from the pharynx.

Strong
acids.

Laryngitis generally attacks adults, and is the result of exposure to cold, to the poison of erysipelas or of syphilitic infection. The acute form generally commences with difficulty of swallowing, and pain and tenderness on pressure over the pomum Adami. The voice, at first harsh, becomes stridulous; dyspnoea, often spasmodic, sets in; the face becomes pale, and the lips livid. Cedematous laryngitis is generally associated with some specific cause, as erysipelas or syphilis. There is less tenderness on pressure, but the fauces are swollen and dusky-red. The voice, too, is more rapidly lost. In either form the immediate treatment should consist of plentiful leeching round the neck, the administration of large doses of calomel with antimonials, and the inhalation of steam. In the cedematous variety, the root of the epiglottis and the sides of the glottis should be freely scarified with a probe-pointed bistoury. When, however, dyspnoea and spasm have fairly set in, the sooner tracheotomy is performed the better.

Acute
laryngitis.

Laryngotomy is applicable in adults, when the foreign body is above or in the larynx. It is not to be recommended in children, in disease of the larynx,

Laryn-
gotomy.

Position of
patient.

or when the foreign body is in the trachea or bronchus. The following rules apply to the three operations of Laryngotomy, Laryngo-tracheotomy, and Tracheotomy. If possible give chloroform. Lay the patient on his back, with a pillow under his shoulders, and the head thrown back over the pillow. Place a pad under the neck, to increase the anterior convexity of the cervical spine. Elevate the chin, and depress the shoulders. The head being held perfectly straight, a line drawn through the middle point of the lower jaw, and the notch in the upper border of the thyroid, down to the upper border of the sternum, will give the direction for the incision. These two points should be marked with iodine paint, or by making an indentation with the thumb-nail. The instruments required are—1, a scalpel; 2, dissecting forceps; 3, retractors, button-hooks being as good as any; 4, blunt hooks (for which bent probes answer the purpose); 5, sharp hook; 6, tracheotomy tube.

Laryn-
gotomy.

Laryngotomy may be performed, in great emergency, by the simple thrust of a penknife horizontally through the skin and crico-thyroid membrane, at a point just an inch (in the adult) below the pomum Adami, the wound being then enlarged laterally. When the operation is performed more deliberately, an incision is made through the skin perpendicularly in the middle line, an inch and a half long, extending from the middle of the thyroid cartilage to the base of the cricoid. The sterno-hyoid muscles are then separated, and held aside by retractors, and the crico-thyroid membrane exposed. The knife is thrust well through the mucous membrane into the larynx—air, blood, and mucus being expelled. The opening is enlarged transversely, and an oval canula, curved on the flat, introduced. If time allows, bleeding vessels should be ligatured before the larynx is

opened, some branches of the superior thyroid (the crico-thyroid arteries) being the only vessels likely to be wounded. The canula must be secured by tapes passing round the neck, and may be repeatedly cleared with a feather or with a camel's hair brush. A flannel or sponge, wrung out of hot water, should be placed over the throat, and the patient, if possible, kept in an atmosphere impregnated with steam.

The first steps of the operation for Laryngo-tracheotomy are the same as laryngotomy, except that the external incision must be a little lower. The incision in the windpipe is prolonged through the cricoid cartilage and the first ring of the trachea. This latter incision must be made from below upwards, to avoid wounding the thyroid body. The operation is useful in children, when the obstruction to respiration is either in or above the larynx, and is more rapidly performed in great emergencies than tracheotomy. It should be avoided, if possible, in the adult.

Laryngo-
tracheotomy

In Tracheotomy the steps of the operation are as follow :—Standing on the right side of the patient, make—1st, A free external incision in the median line, from just above the cricoid cartilage to the upper edge of sternum. 2dly, Clear away fat and cellular tissue, and making out the division between the sterno-hyoid muscles, separate them with the handle of the knife, and let them be gently held aside by retractors. If needful secure bleeding arteries, and let large veins be hooked aside with bent probes. The isthmus of the thyroid lies across the third and fourth rings of the trachea, and if possible a spot should be selected for the opening above the isthmus, as near the cricoid ring as possible. The division of the isthmus in children is of little or no consequence. 3dly, On getting the rings of the trachea in view, pass a sharp hook into the upper part of it to steady

Trache-
otomy.

it, and pull it well forward. 4thly, Turning the edge of the knife upwards, puncture the trachea at the lowest point, being careful not to plunge the knife too far in, and incise it freely upwards. The trachea should be opened as soon as possible, and in spite of free venous hæmorrhage, which will most probably cease immediately respiration is restored. 5thly, In-




FIG. 23.

roduce the canula (Fig. 23) fitted on the "pilot .




FIG. 24.

trocar" (Fig. 24), and withdrawing the trocar, introduce the inner tube (Fig 25) ; tie the canula in with tapes passing to the back of the neck. This canula is constructed so as to lengthen or shorten the straight portion, by changing the position of the collar, which is fixed by the screw shown in the figure. The intro-

duction of the tube is necessary when the obstruction is *above* the opening in the trachea. When it is




FIG. 25.

below, the edges of the incision in the trachea should be kept widely open, a spring speculum used for separating the eyelids being a good form of instrument for the purpose, or a large strong hair-pin, somewhat stretched, the free ends bent into hooks. An attempt may be made to catch the foreign body with a pair of tracheal forceps, but it should not be persevered in. Better to keep the tracheal wound open and wait for the subsidence of the spasm, and then renew the attempt.

If during the operation respiration should cease, the tube must be introduced as rapidly as possible, and artificial respiration immediately commenced, and persevered with for a considerable time, until natural respiration is restored. If any blood gets into the canula and obstructs the entrance of air, it must be sucked out with the mouth, if it cannot otherwise be disposed of.

In the absence of good assistants, and when every moment is of consequence, the following method of performing tracheotomy will be found useful:—Place the patient as before described, and standing on the right side, put the forefinger of the left hand on the left side of the trachea, and the thumb on the right, and make uniform and deep pressure until the carotids on both sides are felt pulsating. Firmly grasp the

Alternative
plan.

trachea between the forefinger and thumb. Then, with the right hand, make the incision in the median line, carefully dissecting down to the trachea; and, if needful, from time to time laying aside the knife and feeling your way with the right forefinger. Until the trachea is opened the left forefinger and thumb should not be removed, but should fix the trachea and press it forwards into the wound. If no tracheotomy tube is at hand, one may be improvised by bending a bit of glass tubing in the flame of a spirit lamp and making a shield out of a piece of tin.

Simple
tracheotomy
tube.

Foreign
body in
bronchus.

If there is reason to believe that a foreign body has become impacted in either bronchus, the patient should be inverted and slapped smartly on the back, by which means the foreign body is sometimes dislodged and ejected through the wound. This failing, and it having been ascertained by the stethoscope in which bronchus it is lodged, a bit of pliant wire doubled so as to form a loop, and bent with its convexity slightly towards the obstructed bronchus, may be passed down and beyond the foreign body, when on the withdrawal of the wire it may be caught and brought out of the wound.

Cut throat.

In cut throat the first thing to be done is to search for all bleeding vessels, veins as well as arteries, and either twist or ligature them. If hæmorrhage is severe, and the bleeding point cannot at once be seen, compress the common carotid against the transverse process of the fifth or sixth cervical vertebra, and then continue the search. If it is impossible to arrest the hæmorrhage, put a ligature on the common carotid. (See below.) In the next place, all coagula should be cleared from the wound. If the air-passages are not opened, the wound in the integuments must be brought together with sutures and strapping. The fact of the air-passages being opened will be ascertained by the air being seen and heard

to bubble in and out of the wound during respiration. If the wound is inflicted above the hyoid bone, it is possible that the tongue may fall backward, and cause suffocation. A ligature should be passed through the tip, and secured to the cheek by sticking-plaster. If the wound is just above the thyroid cartilage, a portion of the epiglottis may be divided, and obstruct respiration. It should be either cut off altogether, or else secured in its place with a suture. If the trachea is entirely divided, the divided edges may be kept together by some sutures. *On no account apply sutures to the external wound*, except at the angles, where the wound is superficial. If there is considerable venous hæmorrhage, a silver tube should be inserted through the wound in the trachea, and the surrounding parts plugged with lint or cotton-wool. Place the patient in bed, with the shoulders well raised, and the head thrown forward on the chest, and secured there by tapes passing from a nightcap to a roller round the chest. The temperature of the room should be raised to 80°, if possible, or at any rate the bed should be placed near the fire and surrounded with a screen. Steam should then be conveyed for the patient to breathe by a piece of india-rubber tubing, placed on the spout of a kettle boiling on the fire. If the pharynx or œsophagus is wounded, the patient must be fed with the stomach-pump tube passed through the mouth, and if needful, guided down the gullet with the finger placed in the wound. Should the patient be much depressed, beef tea and brandy must at once be administered in this way, or by the rectum.

In stabs and punctured wounds of the throat, the general treatment is the same as for cut throat. If the trachea is wounded, emphysema may ensue. A pad of dry lint, firmly applied, will suffice. When any large vessel is wounded, the wound must be

Stabs in the throat.

Ligature of
common
carotid
below the
omo-hyoid.

enlarged, if needful, and the bleeding vessel secured. If this cannot be done, the common carotid must be ligatured, generally below the omo-hyoid. For this operation the following instruments are required:—1, Scalpel; 2, dissecting forceps; 3, blunt hooks and retractors; 4, grooved director; 5, aneurism needle; 6, ligatures.

Place the patient on his back, with raised shoulders, the head being thrown back and turned to the opposite side.

1st, Make an incision through the skin, platysma, and superficial fascia, extending along the anterior border of the sterno-mastoid, from a point opposite the thyroid cartilage, to the sternum.

2dly, Relax the sterno-mastoid, by turning the head slightly, and pull this muscle outwards, the sterno-hyoid and thyroid being drawn inwards, with blunt hooks. Look for the anterior belly of the omo-hyoid in the upper part of the wound, its fibres being seen passing downwards and outwards.

3dly, Divide the dense fascia attaching this muscle to the sheath on the director. The descendens noni will then be seen on the sheath, and must be drawn to the inner side.

4thly, Pinch up the sheath with forceps, and divide it with the knife, cutting on the flat. Enlarge the opening on the director.

5thly, Insert the aneurism-needle, armed with the ligature, from without inwards, between the vein and artery; care being taken to keep close to the under surface of the artery, so as to avoid the vagus.

6thly, Disengage the ligature from the needle, and tie, taking care not to raise the artery from its sheath. The wound is then to be brought together with sutures, and dressed. During the operation, all veins should, if possible, be held aside with bent probes, to avoid wounding them.

In cases of suspended animation from drowning, Drowning.
the following rules should be observed :—

1st, Remove from the mouth all dirt, saliva, etc., and, pulling the tongue well forward, cause it to be retained in that position either with a pair of hooked forceps, or by passing an elastic band over it and under the chin.

2dly, Remove from the body all clothing that may constrict the chest, and lay the body on its back, with the shoulders raised on a firm cushion placed under the shoulder-blades.

3dly, Adopt Sylvester's method to restore respiration :— Artificial
respiration.

(a) Standing, or kneeling behind the patient, grasp the arms above the elbows, and draw the arms



FIG. 26.

steadily upwards above the head, and keep them so stretched for a couple of seconds (Fig. 26).

(β) Turn the arms down, and press them firmly for a couple of seconds against the sides of the chest (Fig. 27).

Repeat these movements from fifteen to twenty times in a minute, and persevere in them, if needful,




FIG. 27.

for one or even two hours. The expiratory act is much aided by an assistant pressing on the lower border of the ribs and the diaphragm.

4thly, During these efforts, have the wet clothes removed from the body, and let the limbs be rubbed in an upward direction with hot flannels. Dash hot and cold water alternately on the chest. If a warm bath is procurable, the body may be immersed in it for five minutes, Sylvester's method being still pursued.

If there is much congestion about the head and face, the external jugular vein, or the temporal artery, may be opened.

The operation of bleeding from the external jugular vein is thus performed. Select a spot where the vein passes over the sterno-mastoid muscle. Compress the vein just above the clavicle with the left thumb, and open it with a lancet, making the incision in the

Venesection
from jugular
vein.

direction of the fibres of the sterno-mastoid. Care must be taken not to admit air into the vein, by keeping the thumb well pressed on it during the operation until the wound is closed by a firm pad. The external wound in the skin should be rather larger than that in the vein, to avoid the formation of thrombus.

5thly, As soon as natural respiration is established, the patient should be removed to a warm bed, the room being kept thoroughly well ventilated, and free from all crowding. An enema of brandy and beef-tea may be administered. If any fluid is given by the mouth, it must be in very small quantities, and with considerable care, lest suffocation should ensue.

In cases of suffocation from hanging, the treatment is substantially the same as is adopted in case of drowning. The body should be instantly cut down, the ligature taken from the throat, and Sylvester's method commenced. Bleeding from the external jugular may also be useful, and a galvanic current may be passed from the nape of the neck to the pit of the stomach to excite the diaphragm.

In asphyxia from the inhalation of noxious gases, as carbonic acid, etc., the body should be immediately exposed to cold air, water dashed on the face, and artificial respiration commenced. Both in this case and in hanging, if it is difficult to establish respiration through the mouth, tracheotomy or laryngotomy should be performed, and the lungs inflated through the opening.

Hanging.

Noxious
gases.

CHAPTER IV.

THE CHEST.

Fracture of the Ribs—Wounds of the Chest—Contusion of the Lung—Hernia of the Lung—Wounds of the Lung—Emphysema—Pneumothorax—Hæmothorax—Paracentesis Thoracis—Wounds of the Heart—Gunshot Wounds of the Chest.

Fracture of
the ribs.

FRACTURE of the ribs may be from direct violence, the ends being driven inwards ; or it may be from indirect violence, as from a squeeze against a wall, when the rib gives way at its most convex part, and the ends are driven outwards. The symptoms are—1, Pain at the seat of injury ; 2, Increased pain on taking a deep inspiration ; 3, Arrest of the movements of the ribs in breathing ; 4, Crepitus, which may be detected by pressing one hand over the seat of the injury, and the other on the opposite side of the chest. Crepitus is also detected on deep inspiration, or the stethoscope may reveal it when it cannot in any other way be discovered. In simple fracture, the best plan is to apply strips of plaster two inches broad, stretching from the spine behind to the sternum in front, and extending some inches above and below the site of the fracture (Fig. 28). By this means, the ribs are kept perfectly at rest on the injured side.

Treatment
of fractured
ribs.

The occurrence of crackling under the skin, and of hæmoptysis, are indications that the lung has been wounded by the point of the broken rib. The im-

mediate treatment will not vary from that for ordinary fracture of the ribs.




FIG. 28.

A severe blow or fall may cause contusion of the lung, without fracture of the rib. The symptoms are, paroxysmal difficulty of breathing, mucous expectoration, coarse crepitation, and dulness on percussion. The treatment consists of perfect rest in bed, salines and antimonials, turpentine stupes, and linseed-meal poultices.

Contusion of
the lung

In wounds which do not penetrate the chest cavity, but involve the muscular tissue in front, the edges should be approximated with sutures, and the arm brought forward in front of the chest, and secured there with a bandage. In extensive wounds posteriorly, the elbows should be as much as possible approximated behind.

Non-
penetrating
wounds of
the chest.

In wounds of the intercostal arteries, the application of ice and a compress should first be tried. To apply this compress, lay a large piece of linen over the wound, and press the central portion into the wound with the finger. Into this depression insert pieces of sponge, until firm pressure is made. Thus all fear of the compress slipping into the cavity of the chest is avoided. If this plan fails, the wound must be enlarged, and if possible the divided ends of the vessel

Wounds of
intercostal
arteries.

secured. This is difficult to effect, and if the hæmorrhage cannot in any other way be arrested, digital pressure must be kept up.

Penetrating
wounds of
the chest.

Penetrating wounds of the chest are easily diagnosed, the symptoms being the passage of blood and air-bubbles through the wound, the expulsion of air from it, or the protrusion of the lung itself. Emphysema and hæmoptysis are also pretty sure signs of a wounded lung. In deep punctured wounds, when these prominent signs are not present, it is impossible to say for certain whether the wound is a penetrating one or not. No attempt must be made to ascertain this, either by introducing the finger or a probe. A careful search for a foreign body must, however, be made, and its removal effected, if present. Hæmorrhage must be stopped, and the external wound closed. The parts should then be kept perfectly at rest with strapping or a rib bandage.

Hernia of
the lung.

The lung may protrude through a wound, forming a round tumour covered with pleura. It should be returned by gentle manipulation, and if there is much constriction at the neck, the wound may be cautiously enlarged with a blunt-pointed bistoury. If symptoms of gangrene are present, the protruded lung should not be returned, but ligatured and cut off.

Wound of
the lung.

The symptoms of wounded lung are—1. Passage of air and blood through the wound. 2. Expectoration of bright red, frothy mucus. 3. Deep pain in the chest, with cough and dyspnoea and abdominal breathing. If the chest be examined with the stethoscope at the point of injury, immediately after the accident, loud, rough crepitation may be detected. There is generally also great collapse. The treatment must be perfect rest of the parts, by strips of plaster from the spine to the sternum, arrest of all hæmorrhage externally, immediate closure of the external wound, application of ice to the chest externally, and

exhibition of it internally. Stimulants should not be administered in the collapse stage, unless under very urgent circumstances.

Emphysema presents a pale, puffy swelling, which crackles when pressed upon. When resulting from fractured rib, the swelling commences over the site of fracture and spreads up over the trunk and neck. It is sometimes the result of a ruptured bronchus or air-vessel, without any external wound being present. The air escapes into the posterior mediastinum, and thence up into the neck, along the course of the vessels and nerves.

In emphysema resulting from wounded lung, either by a punctured wound or fractured rib, if the pressure of the hand relieves the patient, a bandage should be applied. If, however, pressure increases the dyspnœa, no application should be made. In penetrating wounds it is better to close the external wound at once, and if possible make pressure. If the emphysema should become so extensive as to encumber the patient's breathing, incisions through the skin may be made with a scalpel, or scarificators, and cupping-glasses applied to various parts over the incisions. Antimony and ipecacuanha should be given in full and repeated doses.

Pneumothorax is easily recognised by auscultation. There is diminution, or entire absence, of respiratory murmur, and vocal fremitus, with great tympanitic resonance on percussion. It only calls for immediate treatment when it occasions great compression of the lung, and consequent dyspnœa. The original wound may then be enlarged, or the air may be drawn off with a small trocar and canula, or by the aspirator. (For use of aspirator, see Chap. XII.)

Hæmothorax may be diagnosed by the presence of the symptoms of excessive hæmorrhage, together with dulness on percussion, absence of respiratory

murmur, and immobility of the ribs on the affected side. If the symptoms become urgent, paracentesis must be performed. Owing to the coagulation of the blood, it is difficult to remove the accumulation through a canula ; therefore it is better to enlarge the wound, or make counter-openings in a dependent position..

Paracentesis
thoracis.

The operation of paracentesis is best performed with the pneumatic aspirator. But if this is not to be had, an ordinary trocar and canula will serve. The spot to be selected should be the sixth or seventh intercostal space, midway between the sternum and the spine. An incision should be made with a scalpel through the skin close to the lower rib, and the trocar or needle thrust in. If a canula is inserted, care must be taken to exclude the entrance of air as the fluid ceases to flow ; and an assistant should firmly compress the ribs whilst the canula is being withdrawn. The opening is then to be closed with a pledget of lint and a strap of plaster.

Wounds of
the heart.

Where wounds of the heart are not immediately fatal, the symptoms are those of internal hæmorrhage, with small intermitting pulse. Absolute repose, ice externally, and early venesection must be employed, with the administration of belladonna and digitalis internally.

Gunshot
wounds.

In penetrating gunshot wounds of the chest, arrest external hæmorrhage, remove foreign bodies and spiculæ of bone. If there is only one wound, place the patient on the wounded side. If there is a wound of exit as well, let the upper wound be closed, and the other left for drainage.

CHAPTER V.

THE UPPER EXTREMITY.

Wound of Wrist-Joint—Wound of Palmar Arch—Needle in the Hand—Fracture and Dislocation of the Clavicle; of the Scapula; of the Humerus—Compound Injuries to the Shoulder-Joint—Resection of Shoulder-Joint—Amputation at Shoulder-Joint—Fractures and Dislocations of the Radius; the Ulna; the Bones of the Hand—Compound Injuries to the Elbow-Joint—Excision of Elbow—Compound Injuries to Wrist-Joint—Amputation of the Thumb and Fingers.

IN wounds near the wrist-joint, the ulnar or radial arteries are frequently divided. If possible, both ends of the bleeding vessels should be secured, and the hand and arm placed on a long splint, to secure perfect rest. In incised wounds of the palmar arch, the vessels can generally be secured. In punctured wounds it is not advisable to enlarge the wound, in order to secure the vessel, but a compress must be applied to the wound, and a hard ball being placed in the palm, the fingers must be firmly bandaged over it. Before applying the compress, it is advisable to place a tourniquet on the brachial, and then thoroughly cleanse the wound from all clots and foreign bodies. Each individual finger should then be carefully bandaged. A ready method of compressing the brachial is as follows:—Take two sticks, each about eight inches long: arrange them, one on the outer and the other on the inner side of the arm, with a pad of lint intervening, and tie or strap the ends of the two sticks firmly together. This failing, extreme

Wounds
near
wrist-joint.

Wounds of
the palmar
arch.

flexion of the forearm on the arm, with a pad at the bend of the elbow, will effectually compress the brachial, and prevent further hæmorrhage.

Needle in
the hand.

Needles, or portions of them, are frequently run into the hand. The ordinary practice of cutting down on the needle is a troublesome, and frequently a useless proceeding, as the needle recedes before the knife. A plan, which will rarely fail, is to pass the point of a long curved bistoury under the needle, and cut out a flap. The needle will generally be found lying in the flap, and may be removed with a pair of forceps. The flap is then replaced and secured with a strip of plaster and a bandage.

Fracture of
the clavicle.

The clavicle is generally fractured by indirect violence, such as a fall on the point of the shoulder. The fracture is usually found near the outer side of the middle third. Symptoms:—1, The patient leans towards the injured side; 2, the shoulder falls downwards, forwards, and inwards; 3, the sternal fragment is tilted upwards, and can be distinctly felt; 4, pain when the arm is raised towards a right angle with the body; 5, crepitus, when the shoulders are forcibly drawn backwards. Sometimes the fracture takes place at the acromial extremity, when the external deformity is wanting. In fractures from direct violence the bone is often comminuted, and the intervening fragments sometimes fall out of the line of the bone, and are difficult to restore.

Treatment
of fractured
clavicle.

The usual treatment with the pad in the axilla and the figure-of-eight bandage is quite inadequate to keep the fragments in place; and, if no other means are at hand, bandaging the arm to the side, and supporting the forearm in a sling, is sufficient. The following plan is simple and efficacious:—Two strips of adhesive plaster (Leslie's Holland plaster is the best) are cut each about three and a half inches wide, and sufficiently long—the one part to encircle

the arm, and then the thorax—the other to reach from the opposite shoulder, obliquely across the chest under the elbow of the injured side, and across the back to the point of starting. The first strip is looped round the arm just below the axillary borders, a bit of lint being placed between the plaster and the skin. The arm is then drawn downwards and backwards, until the sternal fragment falls into place.




FIG. 29.



FIG. 30.

The strip of plaster is then carried round the back, and passing completely round the body is stitched to itself as in Fig. 29. The second strip is then applied, commencing at the opposite shoulder, and passing down behind under the elbow, a slit being cut to receive the point of the elbow, thence across the front of the chest to the point of starting, where it is stitched as in Fig. 30. Before passing the plaster below the elbow, the elbow must be pressed well forward and inward and so held until the plaster is fixed in position. The arrangement in front is shown in Fig. 31.

A good pad of cotton-wool should be placed in the palm of the hand.



FIG. 31.

Dislocation
of the
clavicle.

Dislocation of the sternal end of the clavicle forwards is the most common displacement. Symptoms:—1, The head of the bone can be felt on the sternum; 2, the shoulder falls a little back; 3, the clavicular portion of the sterno-mastoid is very prominent; 4, placing the knee against the spine, and drawing the shoulders back, the deformity may be reduced.

It has been mistaken for a tumour of bone, and fractures of the sternal end sometimes very much resemble it.

Dislocation upwards is rare. Symptoms:—1, Depression of the shoulder; 2, elevation of the sternal end.

Dislocation backwards is still more rare. Symptoms:—1, Disappearance of the head of the bone behind the sternum; 2, difficulty in respiration.

The reduction of these dislocations is very simple—viz. by placing the knee against the spine, and

drawing back the shoulders. But in the first two varieties it is almost impossible to retain the bone in position. In dislocation backward, the bone may be retained in position, by placing the patient on his back, with a thick pillow placed between the shoulders, or by bandaging the shoulders to a splint placed across behind, with a thick pad intervening between it and the spine.

Treatment
of disloca-
tions of the
clavicle.

Fractures of the scapula are the result, generally, of direct violence, and are difficult to diagnose on account of the accompanying contusion. Fractures through the spine of the scapula are easily felt on passing the finger over the ridge. Fracture of the posterior angle is best discovered by placing the hand of the affected side on the opposite shoulder. In examining for fractures below the spine, the forearm must be laid across the back.

Fractures of
the scapula.

Crepitus may be detected, by placing the hand on the scapula, and moving the arm in various directions. A fracture of the neck through the supra-scapular notch is rare. The symptoms are described in column 2, p. 56.

Fractures of the body of the scapula are treated by bandaging firmly to the trunk. Fractures of the neck require the shoulder to be raised by a sling round the elbow (see p. 187), and to be kept out from the side by a pad in the axilla, the arm being confined to the side by a bandage.

Treatment
of fractures
of the
scapula.

The scapula is sometimes dislocated from its acromial attachment to the clavicle. The displacement is usually downwards. Symptoms:—1, Falling of the shoulder; 2, prominence of acromial end of clavicle. The treatment is to raise the shoulder, and depress the clavicle, by a belt passing round the elbow and over the clavicle.

Dislocations
of the
scapula.

Injuries about the shoulder-joint are frequently difficult to diagnose. The Tables, from "Hamilton

Injuries
at the
shoulder-
joint.

HAMILTON'S DIFFERENTIAL DIAGNOSIS.

COLUMN I.

Signs of a dislocation. (Cause, a fall upon the elbow, or sometimes a direct blow.)

1. Preternatural immobility.
2. Absence of crepitus.
3. When the bone is brought to its place it will usually remain without the employment of force.
4. Inability to place the hand upon the opposite shoulder, or to have it placed there by an assistant, while at the same time the elbow touches the breast.
5. Depression under the acromion process; always greatest underneath the outer extremity, but more or less in front or behind, according as the dislocation may be into the axilla, forwards or backwards.
6. Round smooth head of the bone sometimes felt in its new situation, and very plainly removed from its socket; moving with the shaft. Absence of the head of the bone from the socket.
7. Elbow carried outwards, and in certain cases forwards or backwards, and not easily pressed to the side of the body.
8. Arm shortened in the dislocation forwards, and slightly lengthened, or its length not changed when in the axilla.

COLUMN II.

Signs of a fracture of the neck of the scapula. (Cause, generally a direct blow.)

1. Preternatural mobility.
2. Crepitus, generally detected by placing the finger on the coracoid process, and the opposite hand upon the back of the scapula, while the head of the humerus is pushed outwards and rotated.
3. When reduced it will not remain in place.
4. The hand may generally, but with difficulty, be placed upon the opposite shoulder with the elbow resting upon the front of the chest.
5. Depression under the acromion process, but not so marked as in dislocation.
6. Head of the bone may be felt in the axilla, but less distinctly than in dislocation; never much forwards or backwards; head of the bone moves with the shaft. Head of the bone not to be felt under the acromion process, although it has not left its socket.
7. Elbow carried a little outwards, but not so much as in dislocation. Easily brought against the side of the body.
8. Arm lengthened.
9. The coracoid process carried a little toward the sternum and downwards.
10. Pressing upon the coracoid process, it is found to be movable, and it is also observed that it obeys the motions of the arm.

HAMILTON'S DIFFERENTIAL DIAGNOSIS.

COLUMN III.

*Signs of separation of the epiphysis.
(Cause, direct blows.)*

1. Preternatural immobility.
2. Feeble crepitus ; less rough than the crepitus produced when broken bones are rubbed against each other.
3. Fragments replaced are not easily maintained in place.
4. The hand can be easily placed upon the opposite shoulder, while the elbow rests against the front of the chest.
5. The depression is not immediately under the acromion, yet higher than in most fractures of the surgical neck, perhaps one inch below the acromion process.
6. Head of the bone in its socket, and not moving with the shaft. Upper end of lower fragment projecting in front when displacement exists, and feeling less sharp and angular than in case of a broken bone ; indeed, being slightly convex and rather smooth, it may easily be mistaken for the head of the bone.
7. Elbow hanging against the side when the fragments are not displaced, but away from the side when displacement exists.
8. Length of arm not changed, unless the fragments are overlapped, or both fragments are tilted upon each other. When the fragments are overlapped, the arm is shortened.

COLUMN IV.

Signs of fracture through the surgical neck. (Cause, generally direct blows, but in aged people, frequently caused by falling on the elbow.)

1. Preternatural mobility often, but not constantly present.
2. Crepitus produced easily when there is no impaction, or when the displacement is not complete, but with difficulty when impaction exists, or the displacement is complete.
3. When once the fragments have been displaced, it is exceedingly difficult ever afterwards to maintain them in place.
4. The hand can be easily placed upon the opposite shoulder, while the elbow rests against the front of the chest.
5. A slight depression below the acromion, not immediately underneath its extremity, but an inch or more below.
6. Head of bone in the socket, and moving with the shaft when impacted. The upper end of the lower fragment being often felt distinctly pressing upwards toward the coracoid process, its broken extremity being easily distinguished by its irregularity from the head of the bone.
7. Elbow hanging against the side when the fragments are not displaced, but away from the side when displacement exists.
8. Length of arm unchanged, unless the fragments are impacted or overlapped, or both fragments are much tilted inwards. If the fragments are completely displaced, the arm is shortened.

on Fractures and Dislocations" (pages 56, 57), present the points of similarity and difference between the various injuries.

Fracture of
the ana-
tomical neck
of the
humerus.

Fracture of the humerus at the anatomical neck is rare, and the result of direct violence. It may happen that the head of the bone is thrust through the capsular ligament into the axilla. If it remains in the glenoid cavity there is little displacement of the fragments, and little deformity. Pain at the seat of fracture, and crepitus, with the absence of the signs of other injury, are the chief indications. Sometimes impaction occurs, and then the humerus is shortened, the acromion projects more than is natural, and the globular head of the bone is not distinctly felt. This accident is usually accompanied with fracture of the tuberosity, so that crepitus is felt on firmly grasping the head of the bone and rotating the arm.

For symptoms of fracture at the epiphysis and surgical neck, see columns 3 and 4, p. 57.

Treatment
of fractures
of the upper
extremity of
humerus.

Fractures of the anatomical neck, without displacement, and impacted fractures, may be treated in the simplest manner, by bandaging the arm to the side, and securing perfect rest. Great care should be taken, when the fracture is impacted, not to displace the fragments by forcible manipulation. In fracture at the epiphysis, and at the surgical neck, when there is displacement, the splint (Fig. 32) is useful. The crutch is placed in the axilla, the strap and pad passing over the shoulder. The arm and forearm are then bandaged to the splint, and extension made with the key. Care must be taken, in making extension, not to press too firmly on the axillary vessels. A more simple plan is to mould a firm gutta-percha splint to the outer side of the arm, taking care to make it fit as a cap, well over the shoulder. The arm is then bandaged to this splint, and the upper turns of the bandage passing over the

shoulders and pressing against the acromion, draw the lower fragment outwards. The arm is then placed in a sling.




FIG. 32.

Fractures of the shaft of the humerus are usually the result of direct violence, sometimes of muscular action. If, as is usual, the fracture is transverse, there is little displacement; when oblique, the fragments over-ride. If the fracture is above the insertion of the deltoid, the upper fragment inclines inwards, and the lower one outwards; if below the insertion, the reverse effect is observed. An angular splint should be applied on the inner side, extending from the axilla to the hand, with short splints from the shoulder to the elbow externally, and posteriorly, if needful (Fig. 33). The ribbed "Crimean arm splints" made by Aitken, of York, are admirably suited for this purpose (Fig. 34).

Fractures of
the shaft.

Treatment.

The lower extremity of the humerus is fractured above the condyles, by falls upon the bent elbow.

Fractures of
the lower
extremity of
the humerus.

In young subjects, the lower epiphysis is separated in the same way. In consequence of the great and




FIG. 33.




FIG. 34.

rapid swelling, considerable difficulty sometimes arises in diagnosing between fracture of the humerus and dislocation of the radius and ulna backwards. The former injury is always caused by falls on the elbow, the latter never so, but by falls on the palm of the hand. Symptoms:—1, Preternatural mobility, which, however, ceases after a few hours; 2, crepitus, discovered on extending the forearm; 3, reduction on extension, but immediate return of the deformity on flexion; 4, shortening on measurement from the internal condyle to the tip of the acromion; 5, detection of the ends of the fragments, particularly the lower one, beneath the triceps; 6, unnatural relations between the condyles and the olecranon.

Treatment of
fractures
at the elbow-
joint.

These fractures are best treated with gutta-percha splints. The bones being placed in position, the elbow is bent at right angles, and a firm gutta-percha splint (Fig. 35) moulded to the back part of the arm, as far down as the wrist. In order to

secure perfect symmetry it is better to mould the splint to the sound limb. The arm and forearm are firmly bandaged to the splint, care being taken to put plenty of cotton-wool at the bend of the elbow, and to protect all the prominent points of bone with



FIG. 35.

thick padding. Passive motion should be commenced as early as the seventh or eighth day. If considerable swelling exists, and fracture into the joint is suspected, the arm should be placed in a semiflexed position on a pillow, and evaporating lotions applied, to reduce the swelling.

Dislocations of the humerus downwards into the axilla (Fig. 36) are caused by direct blows upon the upper part of the humerus, or by violent leverage

Dislocations
of the
humerus.

Downwards,
or sub-
glenoid.

carrying the elbow away from the side. Symptoms :—
1, A depression immediately under the acromion ; 2,



FIG. 36.

the elbow carried outwards from the side and a little backwards ; 3, the head of the humerus felt in the axilla ; 4, numbness of the arm, and pain ; 5, the elbow being placed against the side, it is impossible to place the hand on the opposite shoulder (common to all) ; 6, lengthening of the arm, from half an inch to an inch.

Forwards,
or sub-
coracoid.

Dislocations forward (Fig. 37) arise from the same causes, except that the force is generally from behind, as, for example, from a fall on the elbow,

when the axis of the limb is directed backwards. Symptoms :—1 and 2, same as above ; 3, axis of the



FIG. 37.

limb directed towards the middle of the clavicle ; 4, fulness underneath the clavicle on the sternal side of the coracoid process.

Dislocations backwards are rare. They arise from falling forwards on the extended arm, or from pushing violently with the arm elevated. Symptoms :—1, Projection under the spine of the scapula ; 2, wide space between the head of the bone and the coracoid process ; 3, forearm carried across the chest ; 4, axis of humerus upwards and outwards, to a point posterior to the glenoid cavity. One

Backwards,
or sub-
spinous.

point of diagnosis common to all is, "That in taking the vertical circumference of any shoulder in which dislocation exists, by means of a tape carried over the acromion and under the axilla, an increase of about two inches over the sound side is an invariable concomitant."

Treatment.

Whenever it is possible, give chloroform in the



FIG. 38.



FIG. 39.

Reduction
by manipu-
lation.

reduction of these dislocations. The method by manipulation may then be used. When the patient is fully under the influence of chloroform, grasp the arm below the elbow with one hand, and carry it well out from the side. Place the other hand on the shoulder, pressing with the thumb upon the head of the bone (Fig. 38). Next, firmly extending the limb, make it describe a semicircle towards the sternum, at the same time pressing the head of the bone towards the glenoid cavity. Then suddenly bring the arm down to the side (Fig. 39).

Reduction
by exten-
sion.

When chloroform is not given, the reduction is accomplished as follows:—The patient being placed on his back, the surgeon places the heel of his foot in the axilla, at the same time grasping the wrist, and making firm extension. The arm should be slightly rotated, and drawn towards the body. After reduction, the arm is to be bandaged to the side, and the forearm placed in a sling. An

evaporating lotion should be applied to the shoulder, if much swelling is present. To facilitate this operation, and give the surgeon more power, I have had



FIG. 40.

two pads made by Messrs. Mayer and Meltzer, of Great Portland Street. The pad consists of a metallic shield, into which is fixed a handle. The shield is well padded, the pad extending well beyond its edges. It is fixed on directly above either condyle, and connected to its fellow by straps and buckles (Fig. 40). The handles give the surgeon considerable power over the limb in making extension, and the pulleys may be attached to them as

Compound
injuries to
the
shoulder-
joint.

well. By the use of these pads, it is likely that the terrible excoriation of the skin, sometimes produced by the use of a towel or rope, may be avoided.

When there is a dislocation of the shoulder, together with a wound into the joint, if the soft tissues are not much injured, the dislocation must be reduced, and the wound treated antiseptically (see chapter on Antiseptic Treatment). A bag of ice should be *continuously* applied, and the patient kept under the influence of opium. If the soft parts are much involved, it is better to resect the joint at once. (See below, p. 68.)

In severe injuries, especially in gunshot wounds, where the bone is very much comminuted, the soft parts lacerated, or the main vessels involved, amputation at the shoulder-joint may be required (see below, p. 68). But every possible attempt should be made to save the limb, and the following plan of treatment, extracted from Surgeon-Major Porter's "Surgeon's Pocket-Book," is worthy of notice. The shattered limb is to be placed on a cushion devised by Dr. Stromeyer, and thus described by MacCormac:—¹

"The cushion may be described as a right-angled isosceles triangle, four inches thick at the apex, which rests against the chest and supports the elbow, the forearm being bent at a right angle with the arm. The cushion gradually thins down till the base is a mere edge, and of the two other angles, one is passed up into the axilla, while the other rests on the chest under the wrist, as represented in Fig. 42. The cushion is readily fastened in its place by a tape round the neck and one round the body, as seen in Fig. 41. When this simple apparatus is applied the arm rests upon it, beautifully supported and in excellent position. Whilst lying in bed, nothing beyond the ordinary dressings are required for the

¹ "Notes and Recollections of an Ambulance Surgeon," p. 36.

wound, and if the patient needs to be transported




FIG. 41.

from one place to another, or is fit to walk about,




FIG. 42.

this can be arranged for with the utmost facility, as cushion, arm and all, can be bound by a broad

bandage to the body, and thus form an immovable whole."

Resection of
shoulder-
joint.

In resection of the shoulder-joint for compound injuries, the instruments required are:—1, Large bistoury; 2, retractors; 3, lion-forceps; 4, saw; 5, bone-forceps. The operation is thus performed:—1st, Enlarge the original wound in a direction which must accord with the condition of the soft parts. A line from the tip of the acromion to the insertion of the deltoid is the usual direction. 2dly, The arm being carried across the chest, rotate it first inwards, and then outwards to allow of the division of the scapular muscles attached to the tuberosities. 3dly, Open the capsule of the joint, the long head of the biceps being, if possible, slipped aside. 4thly, Carry the elbow backwards, and thrust the head of the bone out through the wound. 5thly, The head of the bone being held firmly with the lion-forceps, remove it with the saw, cutting through the tuberosities. The saw best adapted for this, and many other purposes, is one made by Matthews, of Portugal




FIG. 43.

Street (Fig. 43), with a bent shaft and a movable back. The wound being dressed, the arm is abducted and laid on a pillow.

Amputation
at the
shoulder-
joint.

In amputation at the shoulder-joint, the instruments required are:—1, A long stout bistoury; 2, a catling, nine inches long; 3, lion-forceps. The operation is thus performed:—1st, The artery being compressed

by an assistant on the first rib, form an anterior, or antero-posterior flap, according to circumstances, by cutting from without inwards with the bistoury, and dissecting the deltoid upwards. 2dly, Open the capsule with the heel of the knife. 3dly, As in these cases the bone is generally much shattered, and leverage is impossible, grasp the head of the bone with the lion-forceps, and divide, in succession, the scapular muscles attached to the tuberosities. 4thly, Change the bistoury for the catling, and, still holding the head of the bone with the lion-forceps, slip the knife behind it and cut close to the bone. 5thly, Let an assistant follow the knife, with his thumb feel for the artery, and grasp it between the thumb, in the wound, and the fingers outside the flap. 6thly, Cut outwards, dividing the artery, and completing the flap. 7thly, At once secure the axillary artery.

Amputations at lower points of the arm are best performed by the ordinary flap operation—by transfixion. In making the anterior flap, the brachial artery should be avoided, so as not to cut it until the last sweep of the knife in the formation of the posterior flap.

Amputation
of the arm.

Fractures of the lower end of the humerus, involving the elbow-joint, have been noticed at p. 60.

Injuries to
the elbow-
joint.

Fracture of the olecranon is caused by falls on the elbow, or direct blows upon the part. Symptoms:—

Fracture of
the ole-
cranon.

1, Crepitus when the fragments are not separated, or when they are brought together by extension of the forearm; 2, depression between the fragments; 3, semi-flexion of the forearm, and inability to straighten it. Before the age of fifteen, this fracture seldom or never happens. Great swelling soon follows the injury, and obscures the diagnosis. If suspected, the arm should be kept in the straight position, and evaporating lotions applied. Treatment:—The arm must be placed on a straight splint, extending from

near the upper part of the humerus to the wrist-joint, and placed on the front of the arm. It should be fixed in position with strips of adhesive plaster, and then a bandage should be carried upwards from the hand, which, on reaching the elbow, must be made to embrace the upper fragment in a figure of eight, and thus draw it down into position.

Fracture of
the coronoid
process.

Fracture of the coronoid process is very rare. Symptoms:—1, Luxation backwards of the ulna, or the ulna and radius; 2, probable crepitus; 3, the fragment felt in front of the trochlea. Treatment:—The arm must be placed at right angles, and so secured on a rectangular splint.

Fracture of
the neck of
the radius.

Fracture of the neck of the radius is rare. Symptoms:—1, The upper end of the lower fragment is carried forward by the action of the biceps; 2, the displacement is increased on extension; 3, pronation, and inability to supinate. Treatment:—The forearm must be flexed on the arm, and supported by a single dorsal splint, whilst a compress must be placed over the upper end of the lower fragment, and secured with a bandage.

Fracture of
the shaft of
the ulna.

Fracture of the shaft of the ulna alone is generally the result of direct violence. It is frequently found in conjunction with other serious injuries—*e.g.* dislocation of the head of the radius, or dislocation of both bones backwards. Symptoms:—1, Crepitus and mobility, discovered by seizing the bone above and below the point of contusion, and moving it in opposite directions. The direction taken by the fragments is uncertain. When outwards towards the radius, there is slight projection of the styloid process. When accompanied with dislocation of the head of the radius, there is shortening of the forearm. Treatment:—If the fragments fall towards the radius, the finger must be forcibly pressed between the bones in order to restore the fragments; whilst at the

same time extension is made, and the hand is supinated. If they press forwards or backwards, direct force must be used to their extremities in order to reduce the deformity. The arm is then to be put up in splints, as in Fig. 44.




Fig. 44.

If dislocation of the head of the radius is also present, this must, if possible, be reduced. Let an assistant seize the humerus, not in front, but by the condyles, and at the same time flex the forearm, and extend it. Then push the head of the radius towards its socket. Fix the arm to an angular splint, placed upon the palmar surface. Should the fracture be compound, as it not unfrequently is with dislocation, the same plan must be followed, the wound being dressed antiseptically (see chapter on Antiseptic Treatment).

With dislocation of the head of the radius.

In fractures of the shaft of the radius at the middle third, the lower end of the upper fragment is tilted up by the biceps and pronator teres, whilst the lower fragment is tilted in against the ulna by the pronator quadratus and supinator longus.

Fracture of the shaft of the radius.

In fractures of the lower third, the ends of the fragments are generally tilted in towards the ulna. The treatment of these fractures is the same as that indicated above for fractures of the shaft of the ulna.

Colles' fracture.

Colles' fracture takes place an inch above the carpal end of the bone, and is generally caused by falls on the palm of the hand. Symptoms:—1, The patient cannot pronate or supinate, except with great pain; 2, the hand is turned backwards, and to the radial side; 3, the styloid process of the ulna is very prominent; 4, prominence on the dorsum of the wrist, and a corresponding depression in front; 5, crepitus on powerful extension. Fig. 45, from a cast



FIG. 45.

Gordon's splint.

in King's College, shows one aspect of this deformity. Treatment:—The reduction of this fracture, and its treatment by Gordon's splint, is thus accomplished. Supposing the left radius to be broken, reduce the deformity by laying hold of the forearm, placing the thumb of the left hand upon the lower end of the upper fragment, and the thumb of the right hand upon the upper end of the lower fragment, as much to the inner side as possible. Both fragments are thus pushed backwards and outwards, the hand being at the same time strongly adducted.

Apply the bevelled portion of the under splint (Fig. 46), previously padded, against the lower end of the upper fragment. Then place a thick pad over the metacarpus, carpus, and the back of lower fragment, and apply the upper or back splint, its lower end resting upon the dorsal pad.

Both splints are now held in their position by three

straps with buckles attached, the upper strap being placed as high as possible, and the second strap low




FIG. 46.

down, almost at the end of the body of the splint (Fig. 47). This strap serves the double purpose of




FIG. 47.

pressing the bevelled portion of the splint against the lower end of the upper fragment, pushing it backwards, whilst at the same time, acting upon the back splint, the carpus and the carpal fragment are pushed forwards, restoring and maintaining the natural form of the radius.

As the main displacement of the lower fragment is backwards and outwards, it is necessary that the hand should be maintained in a state of abduction :

therefore place a moderately thick pad over the outer margin of the carpus, the third strap being applied encircling the dorsal splint, and the ulnar portion of the front splint (Fig. 47). The indications fulfilled by the last strap are, to force the carpus inwards towards the ulna, and remove the pressure of the upper surface of the carpus from the outer part of the carpal surface of the radius.¹ A more simple method of treating this fracture is with the pistol-



FIG. 48.

splint. To apply this splint the arm should be firmly grasped, and the hand adducted. The splint is then applied to the palmar surface (Fig. 48).

Fracture of
radius and
ulna.

Both bones of the forearm are sometimes fractured together. The cause is, generally, direct violence, or, more rarely, a fall on the palm of the hand. Symptoms:—1, Crepitus and mobility; 2, pain and loss of power; 3, visible bend in the limb.

Treatment:—If there is but little displacement, the limb must be put up in the ordinary manner (Fig. 44, p. 71). If the fracture is oblique and there is shortening, extension must be made. If there is much tendency for the bones to fall together, the limb should be put up in a position of complete

¹ The above description of the application of the splint is from the pen of Dr. Gordon, Professor of Surgery at Queen's College, Belfast.

supination, and the pads should be so adjusted as to well fill the interstices between the bones.

Fractures of the carpal bones are generally compound. When it is a simple fracture, little or no treatment is required beyond rest. Fractures of the metacarpal bones are the result of direct violence, or of blows upon the clenched fist. Crepitus may be detected, and in some cases an angular projection on the dorsum. Treatment:—A ball or a roller is placed in the palm, and the fingers firmly fixed over it, either with a bandage or adhesive plaster. If the fragments are much displaced, traction on the fingers of the broken metacarpal bones will bring them into place, and they may be retained there by a gutta-percha splint, moulded either to the back or palm of the hand, and well padded. Care must be taken to make the splint project on either side of the hand, in order to keep off the pressure of the bandage from the lateral metacarpal bones.

Fracture of the carpal and metacarpal bones.

Dislocation of the head of the radius forwards is caused by falls on the elbow, or on the extended forearm when pronated. It may be caused also in young children, by lifting them up by the arm. Symptoms:—1, The head of the bone can be felt in its new position, and its absence from its natural situation discovered; 2, the arm is inclined outwards and pronated; 3, the tendon of the biceps is relaxed. Treatment:—Extension should be made with the forearm slightly flexed, and whilst this is kept up the elbow must be grasped, and the head of the radius forcibly pushed back into its position. No attempt to straighten the forearm should be made, but it should at once be put up flexed at right angles, and placed in a sling.

Dislocation of the head of the radius.

Dislocation backward is generally the result of a direct blow on the front and upper part of the bone. It is rare without accompanying fracture. Symptoms:

—1, The head of the bone is felt rotating behind the condyle ; 2, the forearm is slightly flexed and pronated ; 3, supination cannot be performed. Treatment :—Extension being made, the forearm is forcibly supinated, the head of the bone being at the same time pressed forward into its place.

Dislocation
of the radius
and ulna
backwards.

Dislocation of both bones of the forearm is an accident to which youth is particularly liable. Dislocation backwards is caused by a fall on the hand, or inner side of the ulna ; or by a violent twist, or a direct blow on the front and upper part of the forearm. Symptoms :—1, The forearm is slightly flexed ; 2, the lower end of the humerus can be felt in front of the elbow ; the olecranon projects behind the head of the radius, becoming more prominent on flexion and less so on extension (just the reverse of what occurs in fractures of the humerus above the condyles) ; 3, the olecranon, instead of being below the internal condyle, is carried about an inch above it ; 4, the forearm is shortened, on measurement from the internal condyle to the styloid process of the ulna ; 5, there is no shortening from the acromion to the internal condyle. Treatment :—Seat the patient sideways on a chair. Place your foot on the chair, and your knee in the bend of the elbow. Take firm hold of the wrist, and pressing the knee firmly against the radius and ulna, forcibly, but slowly, bend the forearm. Thus the coronoid process is brought forward in front of the elbow, and is pulled by the muscles into its place. When reduction is supposed to have taken place, the joint should be flexed and extended, and if this can be done with ease, it is evident that the bones are in place.

Dislocation
outwards.

Dislocation outwards is usually incomplete. Symptoms :—1, Great prominence on the inner side of the internal condyle, with a deep depression below ; 2, prominence on the outer side of the cup of the head

of the radius. Treatment:—Extension and counter-extension being kept up, lateral pressure must be made on the radial side.

Dislocation inwards is a rare accident. Symptoms:—1, The olecranon is felt projecting on the inner side, and the head of the radius indistinctly in the bend of the arm; 2, the external condyle is very prominent; 3, the forearm is flexed and the hand pronated. Treatment:—As above, by extension, counter-extension, and pressure in the direction of the natural position of the bones.

Dislocation inwards.

Dislocation forwards is very rare. Symptoms:—1, The forearm flexed at right angles, immovable and supinated; 2, absence of the olecranon from its natural position; 3, head of radius in the coronoid fossa. Treatment:—Flexion on the knee, as described above. The ulna may be reduced without the radius, which must then be pressed into its position, as before described.

Dislocation forwards.

Dislocations at the wrist are rare, fracture of the lower end of the radius being generally the accident mistaken for it. Dislocation backwards may be caused by falls on the palm, or on the back of the hand. Symptoms:—1, A very abrupt angle caused by the projection of the carpus; 2, absence of crepitus; 3, easy reduction. Treatment:—Simple extension will generally be sufficient.

Dislocation of the carpal bones backwards.

Dislocation of the carpus forwards may be the result of a fall on the palm. The radius and ulna are felt projecting posteriorly, whilst the carpus forms a tumour anteriorly. It can be reduced by extension.

Dislocation forwards.

The most usual dislocation of the thumb is that of the first phalanx backward on the metacarpal bone. Symptoms:—1, The end of the metacarpal bone projects in the palm; 2, the proximal end of the first phalanx is found on the dorsum of the metacarpal. Treatment:—The thumb being protected

Dislocation of the thumb.

with lint, a clove-hitch (Fig. 49) is fixed on it, and extension made; or reduction may be secured by




FIG. 49.

bending the phalanx forcibly backwards, at the same time pressing the base towards the articulation. Dislocation forwards may be reduced by bending the phalanx forwards towards the palm. In cases of




FIG. 50.

considerable difficulty, a pair of forceps, such as those shown in Fig. 50, are useful.

Compound
injury to
elbow-joint.

When the elbow-joint is laid open without further injury, it should be treated antiseptically (see chapter on Antiseptic Treatment). The limb should be placed on an angular splint, and, if possible, ice kept constantly applied. When the bones are severely injured, as in gunshot wounds or crushes, if the vessels are not damaged, primary excision of the joint should be performed.

For this operation the following instruments are required :—1, Scalpel ; 2, blunt hooks and retractors ; 3, lion-forceps ; 4, saw ; 5, bone-forceps. The original wound may be enlarged, if placed conveniently, or a straight incision may be made at the back of the joint, about five inches long. If possible, the triceps should not be divided, but it should be separated with the fascia from the olecranon, and held aside with a retractor. The olecranon process should then be cut off with bone-forceps. The end of the humerus is then cleared, the ulna nerve being slipped forwards over the internal condyle, and held aside with a blunt hook. It must be remembered, that in injuries to the elbow-joint the nerve is sometimes displaced, so that great care must be taken to avoid it, from the very first. The entire condyloid surface of the humerus is then removed with the saw, and the end of the ulna, including the sigmoid cavity with the head of the radius. The section of the bones must be modified, according to the injury they have sustained ; the usual mistake is the removal of too small a portion. Hæmorrhage being arrested, the wound is brought together, and the arm extended on a straight splint placed in front of the arm.

Excision of
elbow-joint.

If the injury to the soft parts and the vessels is so great that amputation is needful, it may be performed at, or in close proximity to, the joint. The best form of operation, both here and in the forearm, is the “mixed method” of skin-flap, with circular division of the muscles. The instruments required are :—1, Long bistoury ; 2, saw ; 3, lion and bone forceps. An oval flap of skin, large enough to cover in the end of the stump, is marked out with the bistoury, either in front, behind, or laterally, as the case demands ; or two short flaps are cut, to meet over the end of the stump, one being sometimes longer

Amputation
at the elbow-
joint and in
the forearm.

than the other, in order to remove the cicatrix from the face of the stump. A circular sweep is then made through the muscles, about an inch below the base of the skin-flaps. The muscles are cleared upwards from the bone, and the section through the bones completed with the saw. If the amputation is at the joint, the bones are disarticulated previous to the section of the muscles.

Compound
injury to the
wrist-joint.

In compound injury to the wrist-joint, the joint must be treated in the same manner as the elbow or shoulder-joint, the forearm and hand being placed on a straight splint. If the bones of the forearm protrude, and the tendons and blood-vessels are injured, amputation must be performed, the one just described being applicable.

Amputation
of the
thumb.

Every attempt should be made to save any portion of the phalanges or metacarpal bone of the thumb. But if the injury renders this impossible, the pha-




FIG. 51.

langes may be removed by the incisions shown in Fig. 51, or the metacarpal bone by the incision marked in Fig. 52. A long narrow-bladed bistoury is carried through the soft parts, between the metacarpal bone of the thumb and index finger, up to the trapezium; it is then turned outwards, and the joint cut through, the incision being completed in the direction of the dotted line.




FIG. 52.

The index and little fingers may be removed by incisions corresponding to the dotted line in Fig. 53. Amputation of the fingers.




FIG. 53.

The point of the knife is entered on the back of the metacarpal bone, and carried by the side, and underneath the finger, and up again to where it first entered. The soft parts are then dissected back, and the head of the metacarpal bone divided with the bone-forceps. The other fingers are removed by an incision such as that marked in Fig. 54. The adjoining fingers being held aside by an assistant, grasp the finger firmly, and place the heel of the knife on the web of the finger. It is then carried upwards to the joint, through

which it passes, and is brought out on the other side,




FIG. 54.

cutting a flap similar to the one first made. If the possessor of the hand is a labourer, leave the head of metacarpal bone, else remove it with the bone-forceps. The second or third phalanx may be removed as follows:—Bend the phalanx at right angles to the




FIG. 55.

next (Fig. 55); place the heel of a bistoury at the side of the joint midway in the thickness of the next phalanx, and draw the knife straight across the joint; straighten out the fingers, and turning the edge of the knife forwards, cut a flap from the palmar surface.

CHAPTER VI.

THE ABDOMEN.

Rupture of Muscles—Rupture of the Peritoneum—Injury to the Viscera—Non-penetrating Wounds—Penetrating Wounds—Protruded Viscera, Constriction, Distension with Flatus, Gangrene, Wounds—Protruded Omentum—Sutures in Wounded Parietes—Foreign Bodies—Obstructed, Inflamed, and Strangulated Hernia—The Taxis—Operation for Strangulated Hernia—Umbilical, Inguinal, Femoral, and Obturator Hernia.

BLOWS upon the abdomen, even of slight character, may rupture the recti muscles. The symptoms are:—1, Sudden pain and swelling; 2, sometimes a palpable sulcus between the torn ends; 3, increased pain on movement. The treatment is—rest, in a position to relax the muscles, with the shoulders raised and the thighs bent on the abdomen. A large pad should be placed over the site of rupture, and retained by a broad roller.

Rupture of muscles from blows.

The peritoneum may be ruptured by blows on the abdomen. The symptoms are:—1, Extreme collapse; 2, excessive pain, especially on movement and pressure; 3, thoracic respiration; 4, tense contraction of the abdominal muscles. If the peritoneum attached to the pregnant uterus, to the mesentery, or omentum, be torn, considerable hæmorrhage may occur, when all the symptoms of great loss of blood will present themselves. The treatment is perfect rest in the recumbent position, leeches to the abdomen, linseed-meal poultices sprinkled over with laudanum, and opium internally. All purgatives must be

Rupture of the peritoneum.

Injury to
the viscera.

avoided, and very light fluid diet administered. Ice must be applied to the abdomen, and the collapse of hæmorrhage counteracted by stimulants freely given. In all cases of severe blows, or crushing force applied to the abdominal wall, where collapse is present and injury to any of the viscera is suspected, perfect rest must be enforced, opium administered, poultices with laudanum applied, and purgation strictly avoided. If the stomach or intestines are probably ruptured, no food of any kind must be given by the mouth, but nutritive enemata, in small quantities, administered by the rectum, every four or six hours. It is also advisable to empty the bladder with a catheter, especially if damage to the kidneys is suspected. The symptoms of injury to the various abdominal viscera are as follow :—

The Liver.—Pain over its region and dulness on percussion from extravasated blood; great collapse, bilious vomiting, and white stools.

The Stomach.—Bloody vomiting and great collapse. It is, however, possible for the stomach to be severely torn without the symptom of blood vomiting.

The Spleen.—Great collapse and symptoms of internal hæmorrhage; pain over the region of the spleen, and dulness from extravasated blood.

The Kidneys.—Frequent micturition, bloody urine, pain in the lumbar region, and retraction of the testicle.

A wound or rupture of the intestine may be followed by emphysema, owing to the escape of flatus into the subperitoneal areolar tissue, and thence into the more superficial parts.

Non-penetrating
wounds of
the parietes.

In non-penetrating wounds of the parietes, especial care must be taken to search for foreign bodies, and remove them. If there is much hæmorrhage, and the vessel cannot be seen, the wound must be enlarged, and the vessel secured; but if in punctured wounds the bleeding is not severe, it may be left, but the

mouth of the wound must not be closed. In incised wounds of the epigastric region sutures must be inserted, as there is considerable tendency to gape. If the muscular tissues are wounded, care should be taken to insert the sutures through the skin only.

Penetrating wounds of the abdomen, especially punctured wounds, should be treated from the commencement with perfect rest and opium, even although, at the moment, no serious symptoms are present. If any of the viscera are wounded, the symptoms will be:—1, Collapse; 2, severe local pain, which soon becomes radiating; 3, vomiting, sometimes of blood.

Penetrating
wounds.

The various viscera are sometimes protruded through wounds in the parietes—"the small intestines most frequently, next the large, then the stomach, and lastly, the cæcum. Of the solid viscera the omentum is by far the most common, and is often associated with that of the viscera" (Poland). When protruded through small wounds, they are likely to be strangulated. If the protrusion be intestine, and it is uninjured, it must be thoroughly cleaned with a stream of tepid carbolised water and returned, care being taken to press it directly backwards, in order to avoid pushing it between the peritoneum and the muscles. If any constriction exists, the external wound must be enlarged, and the fibres of the muscle or fascia divided; but if possible the peritoneum should not be interfered with. If the bowel is so distended with flatus as to prevent its return, it may be punctured in three or four places with a grooved needle, and the air allowed to escape. If the intestine be gangrenous it must be opened, its contents allowed to escape, and then it must be fixed, with one or two sutures, to the edge of the wound. If the intestine has sustained a small punctured wound, the lips of the wound may be held together with a pair of forceps, and a fine silk

Protruding
viscera.

Constriction

Distension
with flatus.

Gangrene.

Wounds.

or catgut ligature tied firmly round the orifice. The ends are then cut off close, and the intestine returned. Incised wounds must be closed with the continuous suture either of silk or catgut, all the coats being included in the suture. The needle used should be a common round sewing needle, and care should be taken to secure perfect coaptation of the serous surfaces of the gut. The intestine is then returned. When a large portion of the circumference of the bowel is divided, particularly if the wound is lacerated, it is better to fix the divided portions by silk sutures to the edge of the external wound.

Protruded
omentum.

If omentum is protruded through the wound, it should be carefully examined to ascertain that no intestine is enclosed in its folds. If it is quite healthy, and free from laceration, it may be at once returned. But if it is dirty or lacerated, or great difficulty is found in returning it, a double ligature should be passed through the base, tied firmly, and the protruding portion cut off. The ligature may be cut short and returned with the omentum.

Sutures to
wounds of
parietes.

When there is an incised wound opening the abdominal cavity, it must be closed with interrupted sutures. The edge of the peritoneum should be included in the deep sutures, and superficial ones should be placed between them, including the skin only. The deep sutures are best inserted by transfixing the edges of the wound with a needle set in a handle, then threading the needle and withdrawing it, pulling the suture back with the needle. Wire is best for the deep, and silk for the superficial sutures.

Foreign
bodies.

When foreign bodies have passed from the œsophagus into the stomach, purgative treatment should be studiously avoided, and the diet should consist of as much solid material as can be conveniently taken, in order that the foreign body may be, if possible, embedded in solid faecal matter, and thus extruded per anum.

A patient with a hernial protrusion is liable to certain accidents other than strangulation of the bowel. An accumulation of gas and faecal matter may become pent up in the protruded bowel, and prevent its return. The symptoms are:—1, Pain; 2, distension of the tumour; 3, gurgling on manipulation; 4, generally constipation; 5, vomiting; 6, absence of the severe constitutional symptoms which indicate strangulation. The treatment consists of warm applications to the tumour, and a stimulating enema of castor-oil, turpentine, and gruel. If the vomiting is not very severe, purgatives may be given by the mouth. Manipulation of the tumour should be avoided as much as possible.

Obstructed
hernia.

The previous condition may pass on into one of inflammation, or this may be set up by a blow on the tumour, the pressure of a badly-fitting truss, or by the extension of a similar condition in the rest of the alimentary canal. The symptoms are:—1, Pain; 2, enlargement of the tumour; 3, if it be omental, increased hardness; if intestinal, increased elasticity, or fluctuation, from the presence of fluid in the sac. The treatment consists of rest, purgative enemata, and the application of ice to the tumour.

Inflamed
hernia.

Usually the earliest symptom of a strangulated hernia is vomiting, first of the food contained in the stomach, then of yellow and green bile, and lastly of brown offensive stercoraceous matter. At the same time the hernial tumour, if not recent, becomes tense, painful, and irreducible. The bowels are confined, and there is a feeling of dragging between the hernia and the bowel. Great prostration comes on, with quick respiration and feeble pulse; the surface is cold, and the countenance anxious. All these symptoms may exist when the intestine is inflamed, but not strangulated.

Strangu-
lated hernia.

The rule is, that when the intestine is inflamed

Difference
between
inflamed and
strangu-
lated hernia.

but not strangulated, the local symptoms first supervene; but when strangulation comes first, and inflammation as a secondary result, the remote and constitutional symptoms are the first to appear, and the local ones afterwards.

When the local symptoms are urgent, and come on after the remote symptoms, it is a strong argument for immediate operation. When the remote symptoms are present, but no hernia is observable, the strictest search should be made for one. If a hernia exists in which no local signs are present, but which cannot be reduced, an operation should be performed: or it should be performed at the site of any tumour which might contain a hernia; or, in the case of double irreducible herniæ, both should be operated on, if, in the first, no obstruction is found.

The taxis.

The attempt to reduce a strangulated hernia by the taxis must be made with great gentleness. The following rules should be observed:—1. It should be very sparingly tried without chloroform, and chloroform should be administered if the first attempt fails. 2. When chloroform has been administered, the manipulation must be very gentle. If gentle manipulations fail, more forcible ones will only damage the bowel, without reducing the hernia. 3. The attempt, in a small hernia, should not extend over two minutes, and in larger ones not over ten minutes, at the outside. 4. It should not be made by more than one person. 5. It should not be repeated, if it can be ascertained that it has been previously tried. 6. It is quite inadmissible, if the strangulation has been for some hours, and if the constitutional symptoms, such as vomiting, collapse, and tender abdomen, are present; or if it can be ascertained that the hernia has been habitually irreducible. If the symptoms are not very urgent, a grain of opium may be given, and the patient placed in a very hot bath, the

taxis being attempted in the bath. If the hernia be not reduced, but no bad symptoms—such as severe pain, vomiting, or abdominal tenderness, are present—the patient may be placed in bed, and kept under the influence of opium, a strict watch being kept for the onset of any bad symptom. The rule of waiting is, however, a very exceptional one, and the condition of the patient must be very favourable to permit delay.

The use of the aspirator, as an aid to taxis, has been introduced, with considerable success, by Dr. Dieulafoy. The aspirator being exhausted, one of the smallest needles is introduced into the hernial tumour. The cock connecting the needle with the vacuum is then turned, and the needle passed on slowly through the tissues. The first fluid met with will probably be in the sac; but it is not until fecal matter, mixed with bubbles of gas, passes that we can be sure the gut is punctured. When the aspiration of feces and gas is complete, the attempt to reduce the hernia may be made.

Use of the aspirator in taxis.

The description of the operation for strangulated hernia applies, with little modification, to the various forms of hernia met with. The instruments required are:—1, Scalpel; 2, dissecting forceps; 3, straight director; 4, curved hernia-director; 5, hernia knife; 6, retractors. An incision, the direction of which varies with the particular hernia, is made through the skin, and the superficial structures are divided, if needful, on the straight director, until the sac is exposed. The forefinger of the left hand is then passed up to the neck of the sac, and the nail insinuated beneath the stricture, if it be external to the sac. The hernia knife is next guided along the left forefinger and passed on the flat, by the side of the nail, beneath the stricture. The edge is then turned directly forwards, and the stricture divided

The operation for strangulated hernia.

for about a quarter of an inch by a slight sawing movement. If the nail cannot be inserted beneath the stricture the hernia-director must be used and the knife passed up in the groove of the director.

Reduction
without
opening the
sac.

The stricture being divided, an attempt must be made to return the bowel by very gentle manipulation. If the whole contents return with the characteristic gurgle, the surgeon may be satisfied that the stricture is relieved. If, however, reduction does not take place, or some portion remains behind, then the sac must be opened. In order to do this a small portion of the sac must be taken up with the forceps, and nicked through with the scalpel, cutting sideways. A small opening being made, the straight director is passed in, and the sac divided on it for a very small distance. The left forefinger is then inserted and the stricture sought for at the neck, and divided as above directed. If a director is used, great care must be taken not to pass it too far up, as it might perforate the intestine above the stricture. On using the knife, the intestine must be gently pressed back with the left forefinger to prevent its bulging over the edge of the knife and being wounded. In doing this, however, great gentleness must be used, especially if the intestine is in a bad condition, as very little pressure will rupture it.

Opening
the sac.

Examina-
tion and
treatment of
the intestine

On opening the sac the contents must be carefully examined. If the intestine is quite dead, it must be left in the open wound, the stricture above being divided and all the tissues freely laid open. If it is dark, mottled, and studded with lymph, or if there is a doubt as to its condition, it is better to return it to the abdominal cavity, especially if it is elastic and glistening. If there is a small rupture or perforation, it may be pinched up with the forceps and tied with a catgut ligature. If there are recent adhesions, they may be separated; old ones must be left with-

out interference. If the bulk of the intestine is so large from distension with flatus as to impede reduction, it may be punctured with a grooved needle and the flatus allowed to escape. If omentum is present in a recent hernia, and is non-adherent, it should be returned with the intestine; if it is very adherent, it may be left in the sac. If there is much of it, and it is in a bad condition, it may be ligatured and cut off. The greatest care must be taken to examine omental protrusions, in order to discover if they contain, as they frequently do, especially in femoral herniæ, knuckles of intestine. Intestine so found, must be carefully examined to find out if any omental bands are constricting it.

In umbilical strangulated hernia, the incision must be made in the median line, commencing two or three inches above the neck of the sac, and continued as far downwards as may be needful. Every endeavour should be made to relieve the stricture without opening the sac, which is a most fatal proceeding in this hernia. It should be remembered that this hernia often becomes obstructed, with very severe constitutional disturbance, and that aperient enemata frequently rectify this, without recourse to an operation.

Umbilical
hernia.

Inguinal hernia protrudes from the abdomen above Poupart's ligament, the spinous process of the pubes being external to the neck of the tumour. In attempting the taxis, the patient should be placed with the pelvis raised, and the thighs close together, pressure being made upwards and outwards. The operation for the relief of strangulated inguinal hernia is performed as follows:—1st, An incision, from four to five inches long, is made through the integuments, in the axis of the tumour, commencing about two inches above the centre of the external abdominal ring. 2dly, The margins of the external abdominal ring are exposed. 3dly, The remaining tissues are

Inguinal
hernia.

The taxis.

The opera-
tion.

divided down to the sac, which may be recognised by its blue translucent appearance. 4thly, The tip of the forefinger is gently insinuated beneath the external ring, but if this is impossible, a director must be passed, and the ring incised with the hernia knife. The finger is then passed on and any constriction divided, when possibly the hernia may be returned. 5thly, If this cannot be done the sac must be opened, as directed above, and the forefinger carried up to the constriction, which is then divided, if needful, on the hernia-director. All the incisions for the relief of constrictions must be made in a line parallel to the linea alba, thereby avoiding the internal epigastric artery.

Direct
inguinal
hernia.

The above rules apply equally to direct inguinal hernia. It must, however, be remembered that the spermatic cord in the male is more prominent in direct than in oblique hernia, and that the deeper coverings are often much thinner, and therefore require greater care in their division.

In the
female.

The only differences between inguinal hernia in the female and in the male are, that the round ligament is substituted for the spermatic cord, and that the hernial protrusion is found in the labium.

Femoral
hernia.

In femoral hernia the protrusion is found below Poupart's ligament, the spinous process of the pubes being internal to the neck of the tumour. The abdominal ring will also be found unoccupied by any tumour. The symptoms of strangulated femoral hernia are generally much more severe than in inguinal. For the taxis, the patient must be placed on his back, the thighs flexed towards the abdomen, and the thigh of the affected side rotated inwards, and crossed over its fellow. The tumour must be pressed first downwards, then backwards and upwards. The operation for stangulated femoral hernia is as follows:—1st, An incision three or four inches long

The taxis.

The opera-
tion.

is made through the integuments, on the pubic side of the neck of the tumour, commencing an inch above Poupart's ligament. 2dly, Poupart's ligament should be clearly defined, and Gimbernats's ligament felt on the pubic side. This may be slightly incised upwards, when, if the hernia is recent, an attempt should be made to reduce the hernia. This failing, the fascia propria must be divided on a director; upon which a layer of fat will be exposed, which must not be mistaken for omentum. On clearing this, the sac is exposed and opened; the finger must then be pushed up to Gimbernats's ligament, and that ligament divided in a line parallel to the linea alba. The incision must be very sparing. The obturator artery, as it comes off from the internal epigastric or femoral, lies close to the neck of the sac, and may be wounded. It should be felt for and avoided.

Obturator hernia appears in the thigh, in front of the capsule of the hip-joint, and below the horizontal ramus of the pubis. In front of it is the pectineus muscle, on the outer side the femoral artery and vein, and to the inner side the tendon of the adductor longus muscle. If well-marked symptoms of strangulated bowel exist, and a tumour is felt in this region, the supposition would be that it was a strangulated obturator hernia. But it frequently happens that no tumour can be felt; then, if no hernia can be found at any of the other outlets, and pain is referred to the course of the cutaneous branches of the obturator and internal cutaneous nerves, also on rotation of the hip-joint outwards, and on pressure at the site of the obturator foramen, an exploratory operation is justifiable. The operation is performed by—1st, Making an incision through the integuments, commencing a little above Poupart's ligament, midway between the spine of the pubes and the femoral artery, and parallel with it. 2dly, The fascia over the pectineus muscle

Obturator
hernia.

being exposed and divided, the fibres of the muscle are separated, and the fat over the obturator externus exposed. 3dly, If the tumour be not then found, the fibres of the obturator must be separated, and the tumour searched for with the finger at the obturator canal. If the hernia is discovered, the tissues round the neck are divided, and if not then reducible, the sac must be opened, and the constriction sought for and incised. Care must be taken to avoid the obturator nerve.

CHAPTER VII.

THE PELVIS.

Fractures and Dislocations—Injury to the Bladder—Foreign Bodies in the Urethra and Bladder—Dilating the Female Urethra—Injuries to the Urethra—Retention of Urine—Use of Catheters—Puncture of the Bladder per Rectum ; by Aspiration ; above Pubes—Cock's Operation for Perineal Section—Retention from Enlarged Prostate—Extravasation of Urine—Injuries to the Penis—Paraphimosis—Phimosis—Injuries to the Scrotum ; Testicle—Rupture of the Vas Deferens—Injuries to the Pregnant Uterus ; to the Vagina—Foreign Bodies in the Rectum—Hæmorrhage from the Rectum.

THE pelvic bones are fractured or displaced, generally by crushing violence. A crush may cause separation at the symphysis pubis, without fracture, a sulcus being felt externally. Pressure on the ilium, or any attempt to stand or walk, is attended with considerable pain. The urine passed is generally bloody. Such displacement is, however, generally accompanied by fracture of the rami, and also of the ilium. In fractures of the ischium, the injury may be discovered by passing the finger into the rectum, or the vagina in the female, or the tuber ischii being grasped and moved laterally, crepitus may be felt. Fractures of the ilium are the result of falls from a height, or crushing violence, and sometimes of muscular action alone. The spinous process may be broken off, the fragment being displaced downwards, and crepitation and motion being distinct. When portions of the wing of the ilium are broken off, movement and crepi-

Fractures
and disloca-
tions of
pubis.

Of the
ischium.

Of the
acetabulum.

tation are easily obtained by placing the hands one on either ilium, and pressing the bones inwards. The broken fragment will be thus displaced. Fractures of the floor of the acetabulum are caused, generally, by falls on the trochanter major, the head of the femur being sometimes thrust through the cavity in the acetabulum. The diagnosis is very obscure. The trochanter is driven in towards the pelvis, and pressure on it causes deep-seated pain. Crepitus is specially felt during the extension of the limb and its subsequent drawing up, and not on flexion and rotation. Symptoms of injury to the pelvic viscera are also usually present. The brim of the acetabulum is sometimes broken by the same force which causes dislocation of the femur on to the dorsum ilii. Added to the symptoms of the dislocation there is obscure crepitus, and inability to retain the head of the bone in the socket after reduction.

Treatment :—In all cases of injury to the pelvis one of the first things to be done is to pass a catheter and empty the bladder ; if the urethra or bladder is lacerated, the catheter must be retained. The patient should then be placed on his back, the knees bent and supported on pillows. In separation of the symphysis, or fracture of the ilium, the parts may be drawn together and kept in position by a pelvic band firmly buckled round the pelvis. If portions of bone are detached, as portions of the brim, or the anterior superior spine, they may be kept in position with a pad and strapping. If the injury is compound and the fragments have penetrated the abdomen, the wound should be enlarged and all the fragments removed, if possible. In fracture of the acetabulum with dislocation of the femur, the head of the bone should be replaced in the acetabulum and retained therein by the long splint and extension, as for fracture of the thigh.

Fractures of the sacrum may occur from falls on it, or by crushing violence. Simple fractures are rare, and may be detected through the rectum, the lower portion of the sacrum being carried forward. The symptom is pain at the seat of injury, especially in the attempt to defecate. The principal treatment is rest on the back, with a firm cushion under the upper fragment to take off pressure from the lower one; and opium to confine the bowels. If the displacement forwards is very great it may be reduced by the finger in the rectum, and a tube having been passed up to provide for the escape of flatus, the rectum should be plugged round it.

Fracture of
the sacrum.

The bladder may be ruptured by falls, or by blows upon it when distended. The rupture is generally found posteriorly, and urine is extravasated into the peritoneal cavity. Intense abdominal pain, rapid collapse, and urgent desire to micturate, without the ability to pass a drop of urine, and the absence of water in the bladder on the introduction of a catheter, are the prominent symptoms. Death generally ensues at an early period. Should the patient survive, the immediate treatment must be directed to keeping the bladder in a state of collapse by not allowing urine to accumulate in it. For this purpose a large flexible catheter should be passed, great care being taken not to introduce it beyond the neck of the bladder, to avoid passing it through the rent, into the peritoneal cavity. The best form of instrument is a flexible india-rubber catheter (Fig. 59, p. 105), which may be retained in the urethra, as described below. Wounds of the bladder are inflicted through the abdomen, rectum, or urethra, and must be treated as above directed. If symptoms of extravasation appear, free incisions should be made into the affected parts at the earliest moment.

Injury to
the bladder.

Foreign bodies are frequently impacted in the male

Foreign
bodies in
the urethra.

urethra. Slate-pencils, penholders, etc., may be pushed down the urethra; when an endeavour should be made to seize and extract the substance with a pair of urethra forceps, such as those made by Mayer



FIG. 56.

and Meltzer (Fig. 56). These forceps are constructed with a double joint, so that they open widely at the points, without largely distending the anterior portion of the urethra. A hair-pin, with the bent end foremost, has been passed down the urethra, and the ends springing apart, have become imbedded in the urethra. To extract it, a tube should be passed down the urethra and the points of the pin pressed together and placed in the tube, where they become fixed, when the tube and pin may be withdrawn together. Fragments of a calculus, or a small calculus, especially in children, may become impacted in the urethra; they may be removed by the forceps, or else pushed back into the bladder with a catheter. If a foreign body is impacted in the urethra, in front of the scrotum, and it cannot be removed by other means, the urethra must be incised. The incision should be made parallel with the long axis of the urethra, and must be longer externally than through the mucous membrane. There is no occasion for the use of sutures, nor for the retention of a catheter during the healing of the wound. Foreign bodies introduced into the female urethra quickly pass into the

bladder. If they cannot easily be seized with forceps and withdrawn, the urethra may be dilated by introducing the blades of a long narrow pair of forceps, and slowly opening them, the patient being placed under the influence of an anæsthetic. In a few moments the urethra becomes dilated sufficiently to admit the point of the little finger, which should be gently, but firmly passed in ; then the index finger may be introduced in the same way, and the interior of the bladder explored. A pair of forceps should then be passed by the side of the finger, the foreign body seized in a favourable position, and withdrawn.

Foreign bodies in the female bladder.

Dilating the female urethra.

Incised wounds of the urethra heal readily, when inflicted in its long axis. The principal danger arises from extravasation of urine. If, therefore, the wound is in the deeper portions of the urethra, it is better to enlarge the external wound freely, in order to give plenty of vent for the urine.

Injuries to the urethra. Incised wounds.

The urethra is frequently lacerated, especially in youth, by such accidents as a fall on the perinæum, astride a rail. The immediate external signs are generally very slight: the chief symptoms are—bleeding from the urethra, retention of urine, pain at the point of injury, and extravasation of blood into the perinæum. An attempt should at once be made to pass a catheter into the bladder. It should be a flexible one, either such as that referred to at p. 97, or one of the French probe-pointed flexible catheters. If it can be introduced, and the water drawn off, it must be retained in the bladder. If, however, the catheter cannot be passed, it is better at once to incise the perinæum through the median raphé. It may then be possible to discover the divided ends of the urethra, and to bring them together with a suture, after which a catheter may be passed into the bladder. If much hæmorrhage occurs after the incision in the perinæum, the wound must not be

Lacerated wounds.

plugged, but the bleeding must be controlled by digital pressure, provided the ends of the vessels cannot be secured.

Retention
of urine.




FIG. 57.

employ a metallic instrument. A probe-pointed

When a patient is suffering from retention of urine, it is needful, as far as possible, to determine the cause of the obstruction. It may be (1) inflammatory, when there will be a history of purulent discharge from the urethra, which has probably suddenly ceased; or (2) it may be the result of organic stricture; or (3) of enlarged prostate gland. Whatever may be the cause, an attempt should be made at once to pass an instrument into the bladder. Before using the instrument it should be ascertained that it is perfectly clean and pervious. Carbolic oil should be used as a lubricant. As a rule, the flexible probe-pointed French catheter is the instrument which will be the most easily introduced — certainly it is the one which produces the least pain in its use, and which is likely to inflict the least injury. It is essentially useful in strictures of the inflammatory type, and in enlarged prostate. In old organic strictures, it will generally be found needful to

catheter, invented by Sir Henry Thompson, and made by Coxeter, is a very useful instrument (Fig. 57). If the retention arises from enlarged prostate, and the French catheter fails, the usual prostatic catheter must be used, or a full-sized gum-elastic, which should be kept for the purpose on an over-curved stilette (see p. 104). If the first attempt to introduce an instrument fails, recourse must then be had to the hot bath, and a full dose of opium ; or else—and this is the better course—chloroform should be administered, and the attempt again made. The introduction of pieces of ice into the rectum has been recommended, as a means of relief, when all others have failed. Supposing failure again takes place, one of the following plans may be adopted :—Puncture of the bladder through the rectum, which is thus performed :—Place the patient in the lithotomy position ; then pass the left forefinger up the rectum, which must have been previously well cleared out by an enema of water. The point of the finger must be placed just behind the prostate, and as near the centre of the trigone as possible. A tap with the other hand on the hypogastric region will establish the fact of fluctuation. An assistant then supports the bladder externally, by placing the palms of his hands firmly on the lower part of the abdomen ; then pass the rectum trocar, withdrawn within the canula, along the concavity of the forefinger in the rectum. When it is placed at the point where you intend to puncture, press home the handle of the trocar with the palm of the hand, at the same time depressing it. Thus the point will be carried into the bladder, the direction being towards the umbilicus. Then withdraw the trocar, and fix the canula in with tapes attached to the shield, and carried, two in front and two behind, and tied to a bandage round the loins. Puncture by the rectum is not admissible, when fluctuation

Puncture of
the bladder
per rectum.

When
puncture by
the rectum
is not
admissible.

cannot be felt, as will frequently be the case in old-standing cases of stricture, with greatly thickened bladder; neither is it to be used in very great enlargement of the prostate gland. It is also a questionable proceeding in cases where the circumstances lead to the conclusion that the canula will have to remain for a longer period than two or three days.

Aspiration
above the
pubes.

The bladder may be punctured with an aspirating needle above the pubes. One of the finest needles must be passed into the bladder, just above the symphysis; and when a drop of urine appears, the needle is connected with the exhausted aspirator and the tap turned, when the urine will flow into the receiver. When this is full, it is emptied by the discharge-tap, again exhausted, and the process repeated until the bladder is empty. The needle is then withdrawn. The plan is admirably adapted for the immediate relief of a distended bladder; but unless there is a prospect of getting an instrument through the stricture, as may be frequently accomplished after the bladder is emptied, it is better to have recourse to an operation which will secure a permanent exit for the urine. This may be secured by puncturing the bladder above the pubes with a trocar. The operation is thus performed:—The pubes being shaved, place the patient in a semi-recumbent position; make a vertical incision two inches long through the integuments directly above the symphysis; dissect down through the fat, until the top of the forefinger can be placed upon the distended bladder; puncture the bladder with a straight, or slightly curved trocar, the direction being a little downwards; withdraw the trocar, and pass in through the canula either a gum-elastic catheter, or else one of the india-rubber ones, which must be secured by tapes round the loins. In all cases where it is probable that the puncture will be made the exit for the

Puncture
above the
pubes.

urine for some period, this operation is the preferable one. It cannot, however, be performed in cases of greatly thickened bladder, as the bladder does not rise sufficiently above the pubes to permit of its being punctured with safety.

When none of the above proceedings are admissible, it is needful to open the bladder through the perinæum; and for this purpose Cock's operation is the easiest and safest. It is devised in order to open the urethra behind the point of stricture. The instruments required are:—1, A broad double-edged knife; 2, a large silver probe-pointed director with a handle (Brodie's flexible one, used for fistula in ano, will

Opening the
urethra
through the
perinæum.
Cock's
operation.




FIG. 58.

answer the purpose); 3, a perineal catheter (Fig. 58). The operation is thus performed:—Place the patient in the lithotomy position, taking care to keep the pelvis perfectly even. Introduce the left forefinger into the rectum, and place the tip on the apex of the prostate. Plunge the knife into the median line of the perinæum, and carry it forward steadily, *without any withdrawal*, towards the tip of the left forefinger in the rectum; at the same time, by an upward and downward sawing motion, enlarge the external wound vertically in the median line. When the point of the knife is clearly felt by the left forefinger to be in close proximity to the apex of the prostate, carry it forward with a slightly oblique movement into the urethra. Withdraw the knife, and introduce the probe-pointed director, guiding it, with the left forefinger still in the rectum, into the


bladder. Remove the forefinger from the rectum, and holding the director with the left hand, introduce the perineal catheter along the groove, and secure it with tapes in the ordinary way. A flexible catheter may be used instead of the metal one.

Retention
of urine
from
enlarged
prostate.

When the bladder is distended from enlarged prostate, instruments specially adapted for use in such cases must be employed. A large-sized French probe-pointed catheter; the ordinary silver prostatic catheter, from twelve to fourteen inches long, and about No. 10; a gum-elastic catheter, kept for the purpose on an over-curved stilette, and a gum-elastic curved at the beak, like a sound for stone, or Hutchinson's flexible catheter described below, are all instruments which will be found useful in their turn. The first attempt should be made with the French probe-pointed catheter, which in a vast number of cases will succeed, and which is by far the easiest to pass. To pass the silver instrument, place the patient on his back, and standing on his left side introduce the point of the catheter into the urethra, the shaft being held parallel to the left groin. Carry the instrument on as far as it will go without elevating the handle, then bring it gently to the median line of the body, and at the same time raise it a little. By degrees bring it to the perpendicular, and then depress it between the patient's thighs, keeping the beak well against the roof of the urethra. If the point fails to enter the bladder, withdraw it a few inches, and then incline first to one side, and then to the other. If a silver catheter cannot be passed, the over-curved gum-elastic must be tried. Withdraw the stilette, and do not warm the catheter. Pass it over the left groin, pulling the penis well down over the groin, and proceed to pass it in the same direction as the silver instrument. This failing, try the gum-elastic curved at the beak. To obtain this curve,

put the point of the catheter into warm water, curve it to the desired extent, and then plunge it into cold water, which will fix the curve. In all these attempts, the left forefinger in the rectum will be found a great adjunct in tilting up the point of the instrument over the prostatic enlargement, and thus enabling it to enter the bladder.

Hutchinson's flexible catheter has been already referred to. Its method of use is here explained. Fig. 59 represents a silver prostatic catheter, with this peculiarity—that instead of having two eyes, one on either side, there is only one, placed in front. This catheter having been passed into the bladder, a flexible india-rubber catheter or tube, represented in Fig. 60 (it is represented as tied in knots in order to show its perfect flexibility), is passed through the silver catheter, its point emerging through the eye. Then, as the silver catheter is withdrawn, the india-rub-



Hutchinson's india-rubber catheter.

FIG. 59.

ber one is pushed down the tube with the rod (Fig. 61),




FIG. 60.




FIG. 61.

and thus it is prevented from being withdrawn with the silver catheter. In order to retain the india-rubber catheter in the bladder, the metal nozzle with




FIG. 62.

two rings (Fig. 62) is introduced, and fastened to

the penis by tapes and plaster. The stiletted plug (Fig. 62) is then placed in the nozzle. It is found that a flexible catheter thus fixed will be retained in the urethra without the least fear of its being extruded. In very many cases the india-rubber catheter can be introduced without the intervention of the silver catheter. These catheters, and the apparatus complete, are manufactured by Messrs. Krohne and Sesemann, of Duke Street, Manchester Square.

A change in the position of the patient will sometimes enable you to pass an instrument; thus if your endeavours have failed with the patient recumbent, it is possible that you may succeed with him in the erect posture, or *vice versa*. As a rule it is well to at once administer a draught containing quinine gr. ii. and tinct. of opium ℥ x., to be repeated every four hours.

Extravasation of urine generally shows itself first in the tissues of the scrotum, then on the penis, pubes, and abdomen. Free incisions must be at once made, especially in the scrotum, on either side of the middle line, and the patient should be well supported with alcohol, ammonia, and bark. The perinæum ought to be carefully examined, for extravasation sometimes takes place between the layers of the deep perineal fascia. A deep-seated, obscure, brawny swelling may be detected, which must be at once very freely incised in the median line.

Extravasation of urine.

Incised wounds of the penis gape considerably, and must be drawn together with sutures. Effusions of blood from blows may cause prolonged priapism. They must be treated by rest and evaporating lotions. Children have a trick of tying a string round the penis. This causes great swelling, and at a later period ulceration. A careful search must be made for the ligature, which will be found sometimes deeply imbedded in the tissues. The removal of the ligature, and water-dressing, soon brings about a cure.

Injuries to the penis.

Paraphimosis.

Paraphimosis, both in adults and children, should be reduced without delay. The best way is to grasp the penis firmly in the palm of the left hand, whilst with the fingers and thumb of the right hand you steadily compress the glans penis, previously well oiled. The prepuce should be pushed forwards with the left hand, and the glans penis pressed backwards with the right at the same time. Steady and prolonged pressure in this manner will seldom fail. In children chloroform should be given. If the attempt fails, the constricting band must be freely divided with a sharp-pointed curved bistoury, when reduction will be accomplished.

Phimosis.

Phimosis, when acute, must be remedied at once. If, after a short fomentation and syringing out with warm water, the prepuce cannot be drawn back, the following plan will enable you to do so:—Take a blunt-pointed tenotomy knife, pass it down on the flat between the glans penis and the prepuce, then turning the edge towards the prepuce freely incise the mucous membrane. Do this in three places, and you will then be able to draw back the prepuce. This is not to be a substitute for circumcision at some future period, should it be needful.

Injuries to the scrotum.

Incised or lacerated wounds of the scrotum are sometimes attended with extrusion of the testicle. Warm fomentations must be applied for a time to relax the tissues, and the wound must then be carefully brought together with sutures, over the testicle. A cold lotion should then be applied, the patient kept perfectly at rest on his back, and the scrotum raised on a small pillow or sandbag.

Injuries to the testicle.

A blow or a squeeze will frequently cause the testicle to swell, and give excessive pain, both in the testicle itself, and in the inguinal region. Rest, leeches, hot fomentations, and opiates are the remedies. In some cases, if there is much fever, active pur-

gatives, with small doses of tartar emetic, will be found useful.

A violent straining exertion will occasionally cause rupture of the vas deferens. The symptoms are:—
1, A sense of something giving way in the groin, and great pain; 2, swelling of the testicle on the same side; 3, discharge of arterial blood from the penis; 4, clear and bloodless urine on use of catheter. The treatment consists of rest, leeching the testis and lower part of the abdomen, and opiates.

Rupture of
the vas
deferens.

In wounds involving the pregnant uterus, if the fœtus has escaped, an endeavour should be made to bring it through the os uteri. If, however, it presents at the wound, it should be removed and the placenta also. (See Cæsarean Section.) If a limb only protrudes, place it in the uterus, dilate the os uteri, and deliver as soon as possible.

Injuries to
the pregnant
uterus.

Falls on broken crockery may cause severe hæmorrhage from wounds in the vagina. The internal pudic is the most likely source of hæmorrhage, and the cut ends of the vessel must be secured if possible, else digital pressure must be employed, or the wound plugged with lint dipped in perchloride of iron.

Injuries to
the vagina.

Foreign bodies, such as fish-bones, or the husks of apple-pips, cause considerable pain by sticking within the verge of the anus. Frequently, by getting the patient to strain well down, the offending body may be seen, and extracted with a pair of dressing forceps; remembering that in the case of bones or other sharp bodies descending from above, the traction should be made in an upward direction.

Foreign
bodies in the
rectum.

Severe hæmorrhage from the rectum sometimes occurs after the operation for piles, just when the ligatures are separating. The following is Allingham's plan for plugging the rectum:—Take a bell-shaped sponge of fair size. Pass a strong silk ligature up through the apex and back again. Having wetted

Hæmorrhage
from the
rectum.

the sponge and wrung it dry, cover it with powdered alum or persulphate of iron, and pass it five inches up the rectum, apex first, with a penholder or bougie. Fill up the cavity with cotton-wool impregnated with alum or iron. Then, pressing up the wool with one hand, pull down the apex of the sponge by the silk ligature, with the other hand. This inverts the sponge, and fixes it in position. Introduce a catheter or tube by the side to allow flatus to escape. This plug may remain in a week or more.

CHAPTER VIII.

THE LOWER EXTREMITY.

Contusions—Rupture of Muscles—Sprains—Wounds of the Knee-Joint—Ligature of Superficial Femoral ; of Posterior Tibial ; of Anterior Tibial—Fractures and Dislocations of the Femur—Compound Injuries to the Shaft of the Femur and Knee-Joint—Excision of Knee-Joint—Amputation of Thigh—Fractures and Dislocations of Patella ; of Tibia ; of Fibula ; of both Bones—Compound Comminuted Fractures—Compound Dislocation of Ankle-Joint—Fractures and Dislocations of Tarsal Bones ; of Metatarsal ; of Phalanges—Amputation of Leg—Chopart's Amputation—Hey's Amputation—Amputation of Toes.

CONTUSIONS about the hip-joint sometimes simulate impacted fractures of the neck of the femur. In a doubtful case, it is better to treat it from the first as a fracture, with Liston's splint. This will provide the perfect rest necessary for the case, if it be only a contusion. Contusions.

Certain muscles of the lower extremity are subject to rupture, generally by a sudden and violent attempt to recover a lost step, or by some unexpected force being suddenly applied to a certain muscle. The symptoms are :—1, Sudden pain, like a sharp blow from a whip, and sometimes a distinct sound like snapping ; 2, loss of motion in the part, or motion accompanied by very severe pain and spasm ; 3, if the muscle is superficial, a distinct sulcus between the ruptured ends. The ligamentum patellæ, biceps, gastrocnemius, plantaris, and tendo Achillis are the most frequent sites of rupture. Rupture of muscles.

Of liga-
mentum
patellæ.

Of the
biceps.

Of the
gastroc-
nemius and
plantaris.

Of the
tendo
Achillis.

Sprains of
the ankle-
joint.

In rupture of the ligamentum patellæ, which is caused by such movements as an attempt to prevent falling backwards, the patella is drawn up some inches. It must be treated in the same manner as fractured patella (see below), and the effusion of blood dispersed by evaporating lotions. Rupture of the biceps may be caused by severe wrenches, or attempts to lift heavy weights from the ground. The symptoms are severe pain on the outer and posterior part of the thigh, and inability to bear weight on the thigh. If the tendon is ruptured near its insertion, a depression will be felt near the head of the fibula. The treatment is to flex the leg on the thigh and fix it to an outside splint. If flexion cannot be accomplished, a straight Liston's splint must be substituted. The fibres of the gastrocnemius are sometimes ruptured, causing severe pain at the moment and swelling. The plantaris is ruptured in the act of raising the heel from the ground, especially when a heavy burthen is being carried on the back. The tendon sometimes snaps audibly. One of the principal symptoms is dragging and eversion of the foot. The heel must be raised and the knee bent, the treatment being the same as for rupture of the tendo Achillis.

The tendo Achillis is ruptured in the same manner as the plantaris. The treatment is to flex the knee and draw the heel up by attaching a strap to the heel of a slipper, as in Fig. 63. A simple apparatus for treating this accident may be formed, by using a broad dog-collar for the strap above the knee, and attaching it with a bit of string to a loop sewn on to the heel of a slipper.

Sprains of the ankle-joint are of very frequent occurrence, and arise from "turning the foot over." Severe pain and rapid swelling supervene, and it is very difficult to decide in many cases whether the



FIG. 63.

fibula is not broken. In cases of doubt, it is better to act as if it were. In simple sprain of the ankle, the best treatment, if you see the case early, is at once to put on a dozen leeches, and let the bites bleed into a hot bran poultice for an hour or two. Then strap the joint firmly with soap plaster, and bandage it. The next day remove the strapping, replace it more firmly, and make the patient move about on the foot. If the ankle is too painful to bear this treatment, cold evaporating lotions must be used, or hot fomentations if more agreeable to the patient. In gouty subjects an acute attack of gout in the joint and foot is very likely to supervene. This, of course, must be treated by the ordinary constitutional remedies suitable for this condition.

The knee-joint is peculiarly liable to incised and punctured wounds. The principal symptom of wound

Wounds of
the knee-
joint.

into the joint is the flow of synovia. The synovia, if rubbed between the finger and thumb, feels oily and sticky. If any doubt exists as to the joint being opened, the treatment should be the same as if it were wounded; but probing the wound in order to solve doubts should be used as sparingly as possible.

The treatment consists in syringing out the joint with carbolic oil (1 part to 6). This must be squeezed out, and the lips of the wound, if it be incised, accurately brought together with catgut sutures, and covered over with lint steeped in carbolic oil. The limb must then be placed on a MacIntyre splint and swung in a Salter's swing. A bag of ice should be kept constantly applied. If a portion of a needle is run into the joint, it should be dissected out, if possible, as its presence in the joint is sure to set up destructive inflammation. The after-treatment should be that just described.

Of the
femoral
artery.

Application
of tourni-
quet.

A simple
tourniquet.

Digital
pressure.

The femoral artery, from its superficial position, is not unfrequently wounded. It is better at once to apply a tourniquet. The usual method of placing the pad of the tourniquet in Scarpa's triangle is a bad one. The proper place to compress the femoral artery is at the spot where it passes over the pubes, and the best instrument for the purpose is Signoroni's horseshoe tourniquet. The lower pad of the tourniquet is placed on the tuberosity of the ischium, while the upper one is screwed down upon the artery. A tourniquet may be readily made as follows:—Roll a bandage on the middle of a stick a foot long. Place the bandage on the artery, the stick being at right angles to the thigh; then pass a strap and buckle behind the limb and over each end of the stick and tighten. A piece of india-rubber tubing two feet long passed tightly round the limb twice and tied is a good temporary method of arresting bleeding. In making digital pressure, the best plan

is to lay the three middle fingers of the left hand over the artery, and then, with the right hand, to press down on them a large door-key with the handle well padded with lint or cotton-wool. In this way pressure may be kept up for a considerable time without tiring the fingers.

The immediate hæmorrhage being restrained, the wound must, if needful, be enlarged, and the wounded ends tied or twisted. The passage of a probe down to the bottom of the wound will be found a useful guide. The more freely the wound is enlarged, the greater will be the facility with which the vessel will be secured.

Enlarge the wound in search for cut ends.

Supposing all other means fail to arrest the hæmorrhage, the artery must be tied, and Scarpa's triangle will be the most probable spot for the operation. For instruments required, see p. 42. The operation is thus performed:—The patient must be laid on his back, with the thigh slightly abducted and rotated outwards, and the knee bent and resting on a pillow. A line from the middle of Poupart's ligament to the inner side of the knee indicates the position of the vessel. 1st, Make an incision through the skin four inches long, commencing two or three inches below Poupart's ligament, and carried obliquely across the line of the artery from within outwards. 2dly, Divide the fascia lata to the same extent, avoiding the saphena vein, which may appear on the inner side. 3dly, With the handle of the scalpel, or better with the two forefingers, define the edge of the sartorius, an assistant drawing it outwards with a retractor. 4thly, Take up a bit of the sheath of the artery which now appears, with the forceps, and nick it with the scalpel, cutting on the flat. 5thly, Still holding the sheath with the forceps, and making it tense, insert the aneurism needle into the opening, and pass it round the vessel from within outwards,

Ligature of superficial femoral.

making the point emerge at the same aperture in the sheath ; or, if any tissue appears on the point of the needle as it turns forwards, scratch it through with the nail of the forefinger, or gently nick it with the knife. 6thly, Pass the ligature through the eye of the needle, and withdrawing the needle pull the ligature back with it. 7thly, First ascertain, by gently pulling up the ligature and pressing with your forefinger on the vessel, that you have the artery between your finger and the ligature, and then tie in the usual manner. If the catgut ligature be used, the ends are cut short. If the ordinary hempen ligature, one end is cut short, and the other drawn out through the most dependent part of the wound. 8thly, Bring the edges of the wound together with several sutures, and apply a pad of lint and a bandage.

In wounds of the leg, especially when inflicted by thin sharp instruments, like a chisel, and between the tibia and fibula, it is sometimes impossible to secure the bleeding vessel. Ligature of the posterior or anterior tibial then becomes needful.

The operation on the posterior tibial is thus performed :—The patient must be placed on his back, the limb being laid on its outer side, with the knee bent and the heel raised. A line from the centre of the popliteal space to midway between the internal malleolus and the tendo Achillis indicates the course of the artery. 1st, Make an incision through the skin four inches long, one inch from the inner edge of the tibia and parallel with it, commencing two inches above the junction of the upper and middle third of the leg. 2dly, Divide the deep fascia, and having exposed the edge of the gastrocnemius, let an assistant draw it aside with a retractor. 3dly, The soleus being thus exposed, divide its tibial origin along the whole length of the wound on a director. The glistening anterior surface of the muscle must

Ligature of
posterior
tibial.

not be mistaken for the fascia between the superficial and deep layers of muscles, which must be, 4thly, now divided, thus exposing the artery lying on the tibialis posticus. It will be found with its two veins, one on either side, and the nerve somewhat superficial and to the outer side. It is to be secured in the usual way.

The course of the anterior tibial is indicated by a line drawn from the head of the fibula to the base of the great toe.

The operation for ligaturing the vessel in the upper part of the leg is as follows:—1st, Make an incision four inches long through the skin, midway between the spine of the tibia and fibula, commencing two inches above the junction of the upper and middle third of the leg. 2dly, The deep fascia being thus exposed, a white line external to the tibia will indicate the septum between the tibialis anticus and the extensor longus digitorum; these muscles must be separated. 3dly, The anterior tibial nerve will now be seen in front of the artery, which lies on the interosseous membrane between its two veins, and may here be ligatured.

Ligature of
anterior
tibial.

Fractures of the neck of the femur are either intra-capsular, extra-capsular, or extra-capsular impacted. The table on page 119, from Mr. Holthouse's paper in Holmes's "System of Surgery," gives the differential diagnosis in each case. In order the better to diagnose injuries about the hip-joint, Mr. Bryant has adopted a measurement which is illustrated in Fig. 64, taken from Heath's "Minor Surgery." "The patient being recumbent, a vertical line (A C) is allowed to fall from the anterior superior iliac spine, from which the distance (C D) to the top of the trochanter can be measured at a right angle to it. If the neck of the femur be broken, and the trochanter drawn up, this line will necessarily be shorter than on the sound

Fracture of
the femur
at the neck.




FIG. 64.

side.” Fig. 64 shows the usual appearance of fracture of the neck of the femur.

Conditions likely to be mistaken for fracture of the neck.




FIG. 65.

There are some conditions likely to be mistaken for fractures of the neck of the femur.

1. Fractures of the pelvis (see p. 95).
2. Dislocation of the head of the bone on to the pubes (see p. 128).
3. Contusion of hip, where you may have every symptom of fractured neck, with the exception of shortening and crepitus; and as in some cases of fracture there is hardly any appreciable shortening and no crepitus, the difficulty of diagnosis is very great.

Chronic rheumatic arthritis will give rise to symptoms of fractured neck. After a fall on the hip the limb may be found shortened, and pain on move-

TABLE to assist in the Diagnosis of Intra-and Extra-capsular Fractures of the Femur.

INTRA-CAPSULAR FRACTURE.		EXTRA-CAPSULAR FRACTURE.	EXTRA-CAPSULAR IMPACTED FRACTURE.
1. Cause :—	Often slight or indirect.	Falls on the trochanter with great force.	Falls on the trochanter with moderate force.
2. Age :—	Rare before fifty.	Not peculiar to, though more common in the aged.	Ditto.
3. Sex :—	More frequent in females.	Relative frequency in males and females not determined.	Ditto.
4. Shortening :—	Not exceeding an inch. Can be removed by moderate extension, but recurs when this is remitted.	An inch and a half to two and a half. Ditto.	Not exceeding an inch. Cannot be overcome without using great force.
5. Position of the limb :—	It rests on its outer side.	Ditto.	Inclination outwards less.
6. Position of trochanter :—	Nearer the iliac crest than normal; moves in a small arc; head of the bone does not move with it.	Nearer the iliac crest; often detached; does not move with the shaft.	Nearer the iliac crest; moves with the shaft, and the head of the bone moves with it.
7. Crepitus :—	Difficult to detect.	Crepitus readily detected.	None.
8. Helplessness :—	Great.	Ditto.	Limb less helpless.
9. Pain :—	Generally moderate.	Often severe.	Generally slight.
10. Bruising :—	Generally none.	Frequently evident.	Not generally evident.

ment very great. Further examination will show that extension does not remove the shortening, nor can crepitus be produced. A previous history of rheumatic arthritis will also be elicited.

Treatment.

In all cases of injury about the hip-joint, the greatest gentleness must be exercised in making the examination. All violent attempts to procure crepitus are most injurious. For if the fracture is impacted, such attempts will probably disengage the fragments from one another; or, if it be intra-capsular, the capsule will very likely be torn across, and the fragments separated. If there are sufficient signs present to make you even suspect fracture, it is better to act on the supposition that one exists. The treatment will vary with the condition of the patient. If the patient is very advanced in years, infirm, and has any tendency to bronchitic mischief, the complete recumbent posture is not applicable, and the limb had better be put up on a double-inclined plane. This can be easily formed with the aid of a round sofa-cushion and some ordinary pillows, the knee being bent over the pillows, and secured by strips of bandage, or kept in place by sand-bags. Frequently it will be found useful and more comfortable to the patient to place both limbs on the double-incline, and to fix the injured limb to the sound one. The patient should, if possible, be placed on a water pillow, with the shoulders well raised. Great care must be taken to prevent any pressure on the heel. For this, and any similar purposes, the felt-plaster, made by Ewen, of Jermyn Street, is admirably adapted. Or a small circular air-cushion, made of india-rubber, with a hole in the middle to receive the point of the heel, will be found serviceable.

If the patient will bear complete recumbency, then Liston's long splint may be used; and if it is a case of non-impacted fracture, extension should be made.

If it is an impacted fracture, extension must not be used. Sometimes a weight and pulley will be found a useful substitute for the long splint. When required for extension, the weight must be increased; when only to steady the limb, it must be just sufficient for that purpose, sand-bags being placed on either side of the thigh and the leg.

The method of applying the weight is described in a future page.

Fractures of the femur in any portions of the shaft are generally caused by direct violence. The following are the symptoms:—1, Shortening from half an inch to an inch and a half, and deformity; 2, eversion; 3, mobility; 4, crepitus on rotation; 5, pain and loss of power. If crepitus is produced on rotation without making extension, the fracture may be presumed to be oblique; if, on the contrary, crepitus cannot be produced without previous extension, the fracture is probably transverse. The ordinary treatment for fractures of the shaft is with Liston's long splint, and the perineal bandage. The method of application is as follows:—The patient must be placed on a firm mattress, resting, if possible, on a small iron bedstead. A set of fracture-boards intervening between the mattress and the bedstead are a great adjunct to successful treatment. Each fracture-board should be about four or five inches broad, and long enough to reach from side to side of the bed. There must be a sufficient number of them to reach from the foot of the bed to within a few inches of the head. The outer and inner ankles and the tendo Achillis should then be guarded with felt-plaster (p. 120), and the foot and ankle bandaged with a few turns of the roller. The splint, well padded, is then to be applied to the outer aspect of the limb, and must extend from a point opposite the nipple to four inches below the foot, care being taken to make the outer ankle fit into

Fractures of
the shaft of
the femur.

Treatment.

Perineal
band.

the oval hole provided for it. The roller is then carried round the foot and splint, passing through the notches at the extremity of the splint, and being carried up a little above the knee. The perineal band should be made of a piece of bandaging sewn into a long narrow bag, and stuffed with cotton-wool. It should reach from the middle of the groin to a corresponding point behind, and terminate in long tapes. The padded portion rests on the perinæum, and the tapes are passed from within outwards through the two holes at the upper end of the splint. This being adjusted, extension is made until the length of the limb is ascertained to equal its fellow, when the perineal band is to be tightened, and the tapes tied in a bow on the outside of the splint; the upper part of the splint is then confined to the side by a few turns of a broad roller (Fig. 66). In fractures of the lower third, Sir William Fergusson recommends the following ready method:—A long outer splint, and a shorter one for the inner side, rolled up on each side




FIG. 66.

of a double tablecloth until there is just space enough in the middle to contain the limb. The bones being placed in apposition, the limb is then laid between the splints, and retained there by broad tapes or straps.

Fractures of
the shaft in
children.

Fractures of the shaft of the femur in children are best treated with the extension apparatus, described hereafter. Children are very intolerant of the long

splint, and the perineal bandage, being continually wetted, is sure sooner or later to cause excoriation.

Fractures of the condyles of the femur or of the head of the tibia necessarily involve the knee-joint. They are attended with considerable effusion into the joint, and unless the case is seen immediately after the accident, the swollen joint must be treated by evaporating lotions, ice, and leeching if pain and swelling are very great. The limb should be laid on a pillow, and kept in place with sand-bags. If the case is seen early, and there is little swelling, it had better be put up in a MacIntyre splint, and swung in a Salter's swing (Fig. 67).

Fractures
into the
knee-joint.




FIG. 67.

Dislocation of the head of the femur upwards and backwards on the dorsum ilii (Fig. 68) is caused by extreme violence, such as a fall from a height upon the outer side of the knee, a fall on the foot or knee when the thigh is adducted, or the fall of a weight on the pelvis when the body is bent forward. Symptoms:—1, Shortening from one inch and a half to three inches ;

Dislocation
of the femur
on the
dorsum ilii.

2, adduction, rotation inwards, and slight flexion of the thigh, the great toe of the dislocated limb resting on the instep of the opposite foot ; 3, approximation of the great trochanter to the anterior superior spine ; 4, inability to abduct the limb ; 5, in thin subjects the discovery of the head of the bone in its new position, where it may be felt moving in flexion and extension. The following table shows the points of difference between this injury and fracture of the neck of the femur :—

<i>Fracture of the Neck of Femur.</i>	<i>Dislocation on Dorsum Ilii.</i>
1. Crepitus generally felt.	1. No crepitus.
2. Limb mobile.	2. Limb fixed.
3. Toes everted.	3. Toes inverted.
4. Limb slightly shortened.	4. Limb shortened $1\frac{1}{2}$ to 3 ins.
5. Ability to walk on the limb.	5. Inability to bear any weight on the limb.

A test of
dislocation.

The following test applies to all dislocations of the thigh :—Draw a line from the anterior superior spine to the most prominent part of the tuber ischii. When the head of the bone is in place, the trochanter in every position of the limb touches the lower border of this line. When the thigh is dislocated the trochanter is found above this line. The line A B, Fig. 64, p. 118, illustrates this test.

Dislocation
into great
sciatic
notch.

Dislocation upwards and backwards into the great sciatic notch is caused in a similar manner to the last, only the limb at the time of the accident is more nearly at a right angle with the body. The symptoms are very similar to dislocation on the dorsum, but the shortening is not more than half an inch, the flexion is generally less, and the tip of the great toe of the dislocated limb touches the ball of the great toe of the opposite side. In Fig. 69, showing the deformity in this dislocation, the right great toe is not drawn sufficiently forward.




FIG. 68.



FIG. 69.

The treatment in both these forms of dislocation is very much the same. It is directed, not so much to overcome muscular rigidity, which has little or nothing to do with the difficulty in reduction, as to relax the ilio-femoral ligament. To accomplish this an attempt should be made to reduce the dislocation by manipulation. This should be tried at first without chloroform. The operation is thus performed :—

Place the patient on his back on a mattress laid on the floor. Supposing it to be a right limb, grasp the ankle with the right hand and the knee with the left, flex the leg on the thigh, and the thigh on the pelvis

Treatment.

in a direction towards the umbilicus. The flexion of the thigh on the pelvis should cease as soon as resistance is felt. Then move the knee outwards, at the same time rotating the limb slightly outwards, and when the thigh is brought nearly at right angles with the body, bring it down suddenly to its natural



FIG. 70.

position in a straight line with the body (Fig. 70). Should the first attempt fail, it may be renewed under chloroform. Occasionally the head of the bone ruptures the ilio-femoral ligament and passes through a slit like a button-hole in the ligament. The bone being grasped in this slit, the reduction is prevented. To overcome this, before the manipulations above described are commenced the limb should be freely circumducted, by which means the rent in the capsule is enlarged, and the head of the bone permitted to return through it. In dislocations into the sciatic notch the extent of the circuit outwards must be diminished, as it sometimes happens that the head of the bone may be tilted into the foramen ovale instead of into the acetabulum. Should this happen, the

reverse movement must be made, as directed at p. 128. After reduction has been accomplished the two limbs must be tied together, no splints being required.

Dislocation downwards and forwards into the

Dislocation
into the
foramen
ovale.




FIG. 71.

foramen ovale (Fig. 71) is caused by such violence as the fall of a weight on the back of the pelvis, the thigh being at the moment abducted and the body bent. Symptoms:—1, Lengthening of the limb by from one or two inches; 2, bending forward of the body on the thigh; 3, advance of the dislocated limb in front of its fellow, the toes pointing directly forwards;

4, flattening of the hip, and approximation of the trochanter towards the mesial line ; 5, tension of the adductors, the head of the bone being sometimes felt beneath them.

Treatment.

The treatment by manipulation is thus performed :—Grasping the limb as before described, the knee is carried upwards towards the abdomen until resistance is felt. The limb must then be made to describe an arc inwards, and be suddenly brought



FIG. 72.

down to the straight position (Fig. 72). When the knee is just opposite the pubes the thigh should be gently rotated inwards, which helps to tilt the head of the bone into its acetabulum. It may further be assisted into the place by a band placed round the upper part of the thigh and pulled on by an assistant.

Dislocation
on to the
pubes.

Dislocation upwards and forwards on to the pubes (Fig. 73) is caused by falls on the foot, when the leg is thrown backwards behind the centre of gravity. Symptoms :—1, Shortening, abduction, slight flexion and rotation outwards of the thigh ; 2, rotation inwards of the trochanter and flattening at its usual situation ; 3, the discovery of the head of the bone in its abnormal position. The treatment by manipu-

lation is as follows:—Seize the limb by the foot and knee, and rotate the thigh well outwards. Flex the




FIG. 73.

leg on the thigh and bend the knee well over the opposite knee, keeping the heel elevated and the knee firmly pressed down. Continue the motion by carrying the thigh over the sound one as far up as the upper part of the middle third. Then carry the limb directly upwards by elevating the knee, the foot being held firmly, at the same time making gentle oscillations with the knee, when the head of the bone will drop into its socket.

Should manipulation fail in any of these cases, it

Reduction
by extension

will then be needful to have recourse to the pulleys. The patient being laid on a mattress, must be placed




FIG. 74.

under the influence of chloroform, and the parts likely to be galled well protected. In cases of dislocation




FIG. 75.

on to the dorsum or into the sciatic notch, the pulleys will be arranged as in Fig. 74, care being taken to pad the perinaeum well and to bandage the lower part of the thigh above the knee. The traction must be very gradual, and the trochanter must be watched as it descends. Reduction may be assisted by traction with a long towel placed round the upper part of the thigh in a transverse direction. In dislocation into the foramen ovale,

the pulleys must be arranged as in Fig. 75. Less force is required, and the reduction is facilitated by drawing the foot of the dislocated limb towards its fellow, using the strap of the pulley as a fulcrum on which to lever the bone into its socket.

Compound fractures of the thigh are the frequent accompaniments of gunshot wounds. The result of Hamilton's experience may be summed up as follows:—Gunshot fractures of the head and neck of the femur always terminate fatally under amputation, excision, or when an attempt is made to save the limb without recourse to the knife. In the upper third the results are nearly the same, but if the main artery and the principal nerves are uninjured, there is a slight balance in favour of attempting to save the limb. In the middle third the chances between amputation and the endeavour to save the limb are equal. In the lower third the chances are in favour of amputation. If an attempt is made to save the limb, the following rules should be observed:—1. Remove all projecting pieces of bone which cannot be replaced, with the bone-forceps or saw. The lion-forceps and the small saw figured at p. 68 will facilitate this. Remove also all detached portions of bone and all foreign bodies. 2. All portions of muscle or fascia and tendon blocking the wound must be freely divided. 3. Every possible facility must be given for free drainage by counter-openings and the insertion of drainage tubes if necessary. 4. The limb should be put up on an extension apparatus.

Compound
injuries to
shaft of
femur.

Compound injuries of the knee-joint are frequently the result of gunshot wounds. The treatment must be entirely guided by the damage to the soft tissues, by the extent of the fracture, and by the presence or otherwise of injuries to other parts. If the soft parts are not much lacerated, and the fracture is confined to the joint ends, the patient being of fit age, it is

Compound
injuries to
knee-joint.

quite admissible to excise the joint (see below). But if the soft tissues are much lacerated, and the fracture extends up into the shaft of the femur, amputation must be performed (see below).

Operation
for excision
of the knee-
joint.

In performing the operation of excision of the knee-joint the following instruments and appliances are required :—1, A long, strong bistoury ; 2, a saw with a blade at least three and a half inches broad ; 3, lion-forceps ; 4, retractors ; 5, stout scissors, cutting on the flat ; 6, an excision splint ; 7, a Salter's swing ; 8, four strips of adhesive plaster, three inches broad, and long enough to go twice round the limb and splint. The operation is thus performed :—Make an incision with the bistoury across the joint, which must be modified according to the condition of the integuments, but must be carried through the ligamentum patellæ into the joint ; the joint being well flexed, divide the lateral and crucial ligaments. Having cleared the end of the femur, lower the thigh on to the table, the leg hanging over the edge, and being supported by an assistant, and proceed to remove the condyles of the femur with the saw, another assistant steadying the bone with the lion-forceps. The section through the bone must be modified by the direction of the fracture ; thus, if one condyle alone is involved, remove it by an oblique section, sawing off only the articular surface of the other. The thigh must then be raised from the table, and the head of the tibia thrust well upwards, cleared with the knife, and a section removed with the saw. The patella must then be dissected out, the vessels secured, and the limb adjusted on the splint, and retained there by straps of plaster and bandages. The wound is then closed with sutures.

Amputation
of the thigh.

When amputation of the thigh becomes needful, it should be performed as low as possible. It is better to risk making a poor stump than for the sake of get-

ting perfectly sound tissue for flaps to amputate higher up in the thigh.

For amputation at the hip-joint the following instruments are required :—1, A Lister's abdominal tourniquet ; 2, a long straight bistoury ; 3, an amputating knife, twelve inches in the blade ; 4, lion-forceps ; 5, tenacula. The operation is thus performed :—Place the patient with the buttocks well over the edge of the table, and compress the abdominal aorta with Lister's tourniquet. 1st, Make an external semicircular skin-flap, commencing the incision with the bistoury at the centre of the groin just external to the femoral vessels, and carrying it down in a circular direction about four inches below the great trochanter, and up to the tuberosity of the ischium ; retract this flap, and expose the joint. 2dly, The limb being, if possible, forcibly adducted by an assistant, open and disarticulate the joint. When, as is usually the case in traumatic amputations, the bone is shattered very high up, the neck of the femur must be firmly laid hold of with the lion-forceps, and pulled inwards. 3dly, Lay aside the bistoury, and make the internal flap by transfixing with the amputating knife, entering the point in front at the upper angle of the external flap, and bringing it out at the lower angle in front of the tuberosity of the ischium. Cut out through the adductors, and thus form the internal flap ; with a few touches of the knife divide any tissues remaining uncut, and secure the vessels as rapidly as possible.

Amputation
at the hip-
joint.

In amputation of the thigh in the upper third, anterior and posterior skin-flaps, with a circular sweep through the muscles, form the best amputation. In the lower third a long anterior skin-flap and a short posterior are best.

Amputation
in the upper
and lower
third.

In some cases of severe injury of the leg, amputation at the knee-joint, or through the condyles, is

Amputation
at the knee-
joint.

useful. The operation is thus performed :—The instruments required are, 1, a bistoury ; 2, a catling, or amputating knife ; 3, an excision saw ; 4, lion-forceps ; 5, tenacula. Standing on the right of the limb to be amputated (supposing it to be a right limb), 1st, enter the point of the bistoury at the inner side of the knee-joint just below the internal condyle of the femur, and bring it round through skin and fat in a circular direction below the tuberosity of the tibia, to a corresponding point at the outer side of the joint below the external condyle ; 2dly, dissect up the skin-flap, taking as much cellular tissue as possible ; 3dly, flex the knee, and open the joint ; 4thly, lay aside the bistoury, and with the long knife cut a short posterior flap, passing the blade of the knife close behind the head of the tibia, and cutting outwards through all the tissues ; secure the popliteal and other vessels. If it is needful to amputate a little higher, the condyles of the femur may be removed with the saw, an assistant steadying the bone with the lion-forceps. In either case the patella need not be removed.

Fracture of
patella.

Fractures of the patella are caused either by muscular action or by direct blows. Symptoms :—1, Inability to stand or extend the leg on the thigh ; 2, in transverse fractures, which are the most usual, separation of the fragments ; 3, rapid effusion into the knee-joint. Blows on the patella frequently cause longitudinal or comminuted fractures. The immediate treatment of fractured patella must be guided by circumstances. If the effusion is very great, the limb must simply be placed in the horizontal position, and leeches, ice, or hot fomentations applied. The removal of the effused fluid by aspiration will enable the fragments to be at once placed in apposition, and the limb may be immediately placed on Wood's splint (Fig. 76). The fragments are approximated by a

Aspiration
of effused
fluid.




FIG. 76.

figure-of-eight bandage carried round the hooks on the under surface of the splint, and the entire limb carefully bandaged from the toes upwards. A bag of ice may be kept constantly applied to the joint. Compound fractures should be treated in a similar manner.

Dislocations of the patella are caused by muscular action, and occur especially in knock-kneed persons and in women, by twisting the thigh inwards, the foot being fixed. Sometimes the accident is caused by direct violence.

Dislocations
of patella.

Dislocation outwards is the most common form. Symptoms:—1, Limb slightly bent but immovable; 2, breadth of knee increased; 3, projection of internal condyle; 4, patella can be felt on the outer side; 5, great pain. Treatment:—Seat the patient on a chair, and, kneeling before him, place his foot on your shoulder. In the act of raising the limb the patella will frequently slip into place; if not, a little manipulation with the fingers will cause it to do so. Dislocation inwards, except that the patella is reversed in position, presents similar symptoms, and requires the same treatment.

Dislocation
outwards.

Dislocation
inwards.

Dislocation of the patella on to its edge is rare; the projecting edge of the bone is sufficient to show the nature of the injury. Treatment:—If the above

Dislocation
of the
patella on
its edge.

plan does not answer, forcible flexion and extension must be tried, with rotation of the tibia on the femur. At the moment of extension the lower edge of the bone should be pressed on with the handle of a door-key, well protected with lint, by which means the bone is tilted back into its place.

Fracture of
the tibia.

Fractures of the tibia alone are generally the result of direct violence. The nearer the joint, the more transverse is the fracture. The tendency to displacement is very slight, and the treatment simple. A plaster-of-Paris bandage is perhaps the most effective (see Chapter XII.)

Fracture of
the fibula.




FIG. 77.

Fractures of the fibula alone are caused either by direct violence, or by turns of the foot. In the upper two-thirds it is generally fractured by direct violence, in the lower third by indirect violence. Fracture in the lower third is usually accompanied with eversion of the foot, and dislocation inwards of the tibia. Fig. 77, taken from a cast in King's College Museum, shows the deformity. Treatment:—If the external malleolus is alone fractured, or if the fracture is in the middle or upper third,

simple rest on a pillow, or a plaster-of-Paris bandage, will suffice. If, however, the foot is twisted, with displacement of the tibia, indicative of fracture in the lower third, the case had better be treated with Dupuytren's splint (Fig. 78).

Dupuytren's
splint.

This splint requires some care to adjust properly.

It should reach from a little above the knee down to an inch or so below the toes, and must be applied to





FIG. 78.

the inner side of the leg. A pad, two or three inches thicker at its lower than its upper extremity, must be fitted to the splint, so that the lower end rests on the tibia just above the internal malleolus. Fix the upper extremity of the splint with a few turns of a roller, and then inverting the foot, secure it to the lower extremity with a figure-of-eight bandage, taking especial care not to carry the bandage above the external malleolus. Bend the knee, and lay the leg on its outer side on a pillow.

Fractures of both bones of the leg are generally the result of direct violence. Muscular action and falls on the feet will also cause fracture of both bones. The lower third is the most frequent site for fracture of the tibia, the fibula being fractured at the same time frequently somewhat higher. The fractures are almost invari-



Fracture of
tibia and
fibula.

FIG. 79.

ably more or less oblique, the upper fragment of the tibia riding in front of the lower, and more upon the inner than the outer side. When the fracture is just above the ankle, it presents the appearance in Fig. 79 (taken from a cast in King's College Museum).

Treatment :—If there is not much displacement, a MacIntyre splint will be of use. The foot should first be bandaged to the footpiece, which should then be drawn down a little, and fixed with the screw in order to make slight extension. The upper part of the leg and the thigh may then be fixed, and the limb swung in a Salter's swing. In oblique fractures with much displacement, good apposition may be secured by flexing the leg at right angles to the thigh, and keeping it in that position by a light wooden angular splint applied on the outer side, and extending some inches above the knee. The limb must then be laid on its outer side, on a pillow, and the patient's body inclined to that side.

Compound
fracture.

In cases of compound fracture, if there is difficulty in returning the point of bone beneath the skin, endeavour, with a probe, to raise the inverted edges of the skin ; and if the bone cannot then be returned make an incision through the skin sufficiently large to admit of the return of the bone. It may be needful to remove a portion of the protruding bone with the bone-forceps or saw. The limb had better be put up on a back splint, with a footpiece at right angles, and two side splints, which can be secured by straps and buckles. The splint made by Messrs. Arnold (Fig. 80), and used at St. Bartholomew's, is a very useful one. It is thus applied :—Place the leg on the back splint of iron, which must be well padded, carefully fitting the heel into the opening made for it. Secure the foot to the footpiece with a thick piece of gutta-percha moulded to the dorsum, coming round

the edges of the footpiece, and fastened with tapes passed through holes bored in the gutta-percha.




FIG. 80.

Some flannel should be placed between the foot and the gutta-percha, and the latter should be punched with a number of holes for ventilation. Take care that the ball of the heel and the ball of the great toe are well against the footpiece; and, as in all other methods of setting a fractured leg, that the ball of the great toe is in a line with the inner border of the patella. Then apply the side splints, securing them to the knee with a mould of gutta-percha, and further fixing them in place with straps and buckles. Swing the limb to a cradle with leathern straps passed through the two cross-bars of the back splint. The




FIG. 81.

wire splint, invented by Surgeon McNalty, will be

found useful, especially in country practice, as it is very portable. Fig. 81 shows this splint folded up ;




FIG. 82.

Fig. 82 shows it open : and Fig. 83 represents the limb placed in it.




FIG. 83.

In comminuted fractures remove all detached pieces of bone, but any fragments firmly adherent to periosteum, and covered with it, should if possible be preserved. (For antiseptic treatment of compound fractures, see Chapter XI.) As a general rule, if the temperature and sensation of the foot are natural, an attempt should be made to save the limb. If the contrary conditions prevail, amputation had better at once be performed. (See below.)

Dislocations at the knee-joint are generally the

result of severe violence. The symptoms are so pronounced as to require no description. The reduction is accomplished by forced flexion and extension, combined with a rocking motion of the joint. Sometimes it may be needful to apply extension by pulleys, and to make firm pressure on the prominent dislocated bone. Chloroform or ether should be administered if possible.

Dislocations of the tibia at the knee-joint.

Displacement of the semilunar cartilages of the knee-joint may be caused by catching the point of the toe in a stone during walking, and thus wrenching the knee-joint; or by a patient turning in bed, the weight of the bedclothes on the toes twisting the knee. Symptoms:—1, Sudden severe pain; 2, the joint fixed in the semiflexed position; 3, sometimes the patient falls suddenly to the ground. Treatment:—Flex the leg as much as possible, and then suddenly extend it. If the first attempt fails, repeat it with a rocking motion of the joint. Reduction is attended by immediate cessation of pain and return of free motion to the joint.

Dislocations of the semilunar cartilages.

Dislocation of the tibia at the ankle-joint inwards, combined with fracture of the fibula, has been described at p. 136. Dislocation outwards is frequently associated with fracture of the internal malleolus. Fig. 84, taken from a cast in King's College Museum, shows this deformity. Treatment:—Grasp the foot firmly with both hands, and draw on it gently, at the same time rotating the foot outwards.

Dislocations at the ankle-joint.
Outwards.

Dislocation forwards (Fig. 85) is caused by violent extension of the foot. The fibula is frequently fractured. Treatment:—Flex the leg on the thigh; and whilst an assistant makes extension from the foot, press the tibia backwards and the heel forwards. As the bones begin to slide into place, forcibly flex the foot on the leg. Great difficulty is found in keeping the

Forwards.

bones in place, and it may be needful to divide the tendo Achillis, after which the limb should be fixed




FIG. 84.



FIG. 85.

to an outside splint, and laid on its side on a pillow in the semiflexed position.



FIG. 86.

Dislocation backwards is very rare (Fig. 86). It is

usually accompanied by fracture of one or both malleoli. It is caused by any force which produces extreme flexion of the foot. Treatment:—The tendo Achillis will probably have to be divided before reduction can be accomplished. Steady extension, with alternate flexion and extension of the foot on the leg, must be employed.

Dislocation
backwards.

In compound injuries to the ankle-joint every attempt should be made to save the limb, and unless the soft parts are very severely damaged, amputation should not be performed. If the protruding ends of the bone cannot be replaced, they should be removed, the extremities being firmly held with the lion-forceps, and the section made with the small saw figured at p. 68. For the antiseptic treatment see Chapter XI.

Compound
dislocations
of the ankle-
joint.

The astragalus and os calcis are sometimes broken by falls on the feet, and occasionally the latter is broken by muscular action, as in jumping on the toes, the posterior portion of the bone being separated by the tendo Achillis. The other bones of the tarsus are generally broken by crushing violence. Symptoms:—The only prominent symptom in fracture of the tarsal bones is crepitus, and this is not always present. When the posterior part of the os calcis is separated, the fragment is drawn upwards by the tendo Achillis, and the heel is shortened. Treatment:—There is little to be done beyond rest and evaporating lotions. If the os calcis is fractured posteriorly, the case should be treated as in rupture of the tendo Achillis (Fig. 63, p. 113).

Fractures of
the tarsal
bones.

The astragalus may be dislocated in various directions by falls on the foot from a height, accompanied by violent twisting of the foot; or it may be dislocated by a direct blow. Symptoms:—1, Great prominence of the bone in the direction of the dislocation; 2, lateral deviation of the foot; 3, shortening of the leg; 4, in the backward dislocation the position of the

Dislocations
of the tarsal
bones.

foot is not much changed, but its dorsal aspect is shortened. So great is the force needful to dislocate this bone, that it is frequently a compound injury. Treatment:—In simple luxation place the patient under the influence of chloroform, and firm extension being made, endeavour to press back the bone into its place. This failing, divide the tendo Achillis, when reduction may be accomplished. Should it, however, be still impossible, it is better at once to divide the skin over the bone and remove it. In compound dislocation this latter course is invariably the best.

Displacement of the other tarsal bones may be easily reduced by simple pressure.

Fractures of
the meta-
tarsal bones
and
phalanges.

In fractures of the metatarsal bones the usual displacement of the fragments is upwards. They must, if possible, be pressed backwards into position, and retained by a gutta-percha splint, moulded to the dorsum. In fractures of the phalanges, it is very often better to amputate, as the attempt to save the end of the toe is frequently the cause of considerable irritation. If the fracture is to be treated, a broad gutta-percha splint, moulded to the sole of the foot, and supporting all the toes, is better than a narrow one for the broken toe alone.

Dislocations
of meta-
tarsal bones.

One or more or the entire row of metatarsal bones may be dislocated from the tarsus. The direction is generally upwards and backwards. Treatment:—To reduce a single bone, pressure on the head is the most effectual method; but if the whole row be displaced, extension must then be made, in addition to pressure on their heads.

Dislocations
of the
phalanges.

The phalanges are generally displaced backwards. The reduction may be accomplished by combined extension and pressure on the head of the bone.

Amputation
of the leg.


In amputation of the leg the best operation is by lateral skin-flaps, with a circular amputation through

the muscles. Make two lateral flaps of skin sufficiently long to cover the end of the stump, retract the flaps, and divide the muscles one inch below their base by a circular sweep down to the bone; make the section of the bones with the saw on a level with the base of the skin-flaps.

The various amputations through the foot are indicated in the adjoining diagram (Fig. 87), from Bryant's "Surgery," which will be referred to in the following descriptions.

Chopart's amputation is thus performed on the right foot:—Grasp the fore part of the foot with the left hand, placing the forefinger on a point one inch behind H, the prominence of the metatarsal bone of the little toe, and the thumb on A, the tubercle of the scaphoid. Make a curved incision between these two points through the skin, and a second incision through the tendons down to the bone; forcibly bend down the foot, and open the articulations between the os calcis and the cuboid and between the astragalus and scaphoid. Keeping the knife close to the bones, cut the plantar flap as far down as the roots of the toes.

When all the toes require removal, Hey's amputation must be performed. Operation on right foot:—Grasp the foot with the left hand; commence the



Amputation of the foot.

Chopart's amputation.

FIG. 87.

Hey's amputation.

incision on the dorsum, half an inch in front of the point H, the projection of the metatarsal bone of the little toe, and carry it with the convexity downwards to a point half an inch in front of C, the base of the metacarpal bone of the great toe ; make the plantar flap by two lateral incisions, carried from the base of the dorsal flap along the metatarsal bones, and joined by a curved incision a little beyond the ball of the great toe ; reflect this flap, and with a saw divide the metatarsal bones just in front of their articulations with the tarsus.

Amputation
of toes.

The rules for amputation of the toes are precisely similar to those given for the fingers at pp. 81, 82. The head of the metatarsal bone should never be removed if it can be saved.

CHAPTER IX.

EMERGENCIES CONNECTED WITH PARTURITION.

Signs of Pregnancy—Accidental Hæmorrhage—Placenta Prævia—Retained Placenta—Flooding—Induction of Premature Labour—Cæsarean Section—Craniotomy—Cephalotripsy—Embryulcia—Decapitation—Displacements of the Gravid Uterus—Mal-presentations—Turning—Prolapse of the Cord—Puerperal Convulsions—Rupture of the Uterus ; of the Bladder ; of the Perinæum—Inversion of the Uterus.

THE following are the signs of pregnancy during the successive months :—

Signs of pregnancy.

First Month.—Cessation of menses ; morning sickness ; sometimes profuse salivation ; soft, cushiony condition of cervix ; circular form of os ; vagina hotter and secreting more mucus.

Second Month.—Breasts become larger and tender ; veins slightly enlarged ; nipples swollen ; sebaceous matter found in the follicles round the nipple ; secretion of milk ; abdomen becomes flat.

Third Month.—Slight abdominal enlargement ; areola round nipples darker ; os raised higher in vagina ; in thin people the placental bruit may be detected by pressing the stethoscope over the brim of the pelvis on to the fundus of the uterus.

Fourth Month.—Enlargement of abdomen more marked ; uterus felt two inches above the pelvis ; mother feels the sensation of quickening ; foetal heart may perhaps be detected beating from 130 to 150 per minute.

Fifth Month.—The patient being placed in the upright position and the finger introduced into the vagina and pushed against the uterus, ballottement may be obtained.

Sixth Month.—The fundus reaches the umbilicus.

Seventh Month.—Umbilical depression removed; the os is close against the sacrum, and the cervix drawn up much higher.

Eighth Month.—The fundus reaches midway between the umbilicus and the ensiform cartilage; foetal movements plainly felt by the hand on the abdomen.

Ninth Month.—The fundus reaches the ensiform cartilage; the cervix is obliterated.

Accidental
hæmorrhage

Accidental hæmorrhage may occur before delivery from detachment of a portion of the placenta. It is caused generally by falls, blows, violent mental emotions, jars in driving over a rough road, or by railway travelling. The symptoms are, pain at the seat of injury, sense of distension, feeble pulse, fainting, and all the ordinary signs of loss of blood. It is possible that no blood may escape externally, but often the quantity which escapes is enormous. If labour is in progress, the flow will be arrested during a pain, or when faintness comes on. The following are the distinguishing points between accidental hæmorrhage and placenta prævia :—

Accidental Hæmorrhage.

1. Discharge arrested by labour-pain.
2. Membranes felt, with a portion of the child presenting.
3. The os thin.
4. History of injury.

Placenta Prævia.

1. Discharge increased by labour-pain.
2. The placenta felt as a soft substance.
3. The os very thick and pulsating.
4. No history of injury; bleeding coming on of itself.

Treatment :—If slight rest and astringents, such as the Acetate of Lead two grains, with half a drachm of dilute acetic acid in water every four or six hours, fails, plug the vagina with soft linen, tow, or cotton-wool, introduced gradually, and forced well up to the os, a firm binder being applied externally. The pulse should be carefully watched, and internal hæmorrhage carefully looked out for. If the hæmorrhage is very excessive, delivery must be expedited. Puncture the membranes, and give ergot. If needful, turn, or apply the forceps. If the patient is apparently dying from loss of blood, attempt transfusion (see Chap. XII.)

Plugging
the vagina.

In placenta prævia hæmorrhage generally comes on about the seventh or eighth month. It is a sudden discharge, and stops suddenly, to be repeated again probably in a week or fortnight. The other symptoms are given above. Treatment :—If the hæmorrhage is slight, keep the patient cool and in the recumbent posture: give the acetate of lead, and opiates, if there be uterine action. If the hæmorrhage is severe, the sooner labour is commenced and terminated the better. If the os is not dilated, plug the vagina and wait for uterine action. If this is tardy, remove the plug, puncture the membranes with a stilet, and plug again if hæmorrhage continues. If the os does not dilate, the dilating india-rubber bags may be employed (see below), and when the os is sufficiently dilated either turn or apply the forceps. If the patient is too exhausted to bear turning or the forceps, pass the fingers into the uterus, completely separating the placenta, and leave the case to nature.

Placenta
prævia.

Retained placenta from adhesions to the uterine walls may be diagnosed by the fact that uterine action continues after the birth of the child, without expelling the placenta, the uterine contractions being generally accompanied by gushes of blood. On digital examination the placenta will be found closely ad-

Retained
placenta
from adhe-
sion to the
womb.

herent to the walls of the uterus. Treatment:—Introduce the hand into the cavity of the womb, the other being placed on the fundus; feel along the cord for the centre, and thence for the edge of the placenta, and gently peel it off. If it is impossible to remove some portions without the risk of damage to the womb, it is better to leave them.

Retained
placenta
from
uterine
atony.
Retained
placenta
from hour-
glass
contraction.

When the placenta is retained from uterine atony, give ergot, apply pressure with the hand on the fundus, and dash cold water on the vulva and abdomen. In cases of hour-glass contraction, introduce the fingers, cone-shaped, into the womb, and endeavour to dilate the contraction: then grasping the placenta, gently withdraw it. Chloroform is of great assistance.

Flooding.

Hæmorrhage into the cavity of the womb after the expulsion of the placenta is the result of uterine atony. Treatment:—Place the patient on her back, and give stimulants freely; grasp the uterus with one hand externally, and with the other in the cavity of the womb clear out all clots; pour cold water on the abdomen from a height, and, if at hand, introduce ice into the cavity of the womb. Inject two minims of solution of ergotine under the skin (see p. 11). If these means fail, remove all clots and inject four ounces of *Liquor Ferri Perchloridi Fortior* (B. P.), diluted with twelve ounces of water. A Higginson's syringe, with a uterine tube attached, may be used; and the injection must be gently sent well up to the fundus, great care being taken to inject no air. A simpler and perhaps a safer method is to inject hot water at a temperature from 110° to 115° . In either case the hand should be introduced into the uterus and the point of the syringe carried well up to the fundus. As a last resource, transfusion may be had recourse to (see Chapter XII.)

The induction of premature labour is required as

an emergency, in cases of placenta prævia, of pulmonary or cardiac disease, or of convulsions. The best plan is to introduce one of Barnes's india-rubber bags. They are of various sizes, the smallest the size of the finger, with a constriction in the middle, by which it is grasped by the os and kept in situ. Being well oiled, the bag is passed through the cervix as far as the constriction; it is then distended with air or water through the elastic tube attached to the lower end. If the bag be not at hand, an elastic male catheter with the top cut off must be passed through the os, and the membranes ruptured by passing the wire stilet through the catheter. Labour will come on after the lapse of several hours.

Induction
of prema-
ture labour.

Dilating
bags.

Cæsarean section may be called for as an immediate proceeding in cases of sudden death of the mother, provided it can be ascertained that the child is viable, and the stethoscope determines that it is living. If the mother has died of hæmorrhage or asphyxia, it is probable that the child too is dead. Ten minutes after the mother's death is the longest period that can elapse with the chance of saving the child. In opening the uterus care must be taken not to wound the child, and in removing the child it should be grasped so as to bring out the head or the breech first. When Cæsarean section is performed with a view of saving the mother's life as well as that of the child, the following are the directions for the operation. The instruments required are:—1, Large, strong bistoury; 2, director; 3, retractors; 4, dissecting forceps; 5, large curved needle set in a handle; 6, silver wire sutures; 7, catgut ligatures. The rectum and bladder being emptied, place the patient on her back with the shoulders raised and the pelvis elevated. Do not give chloroform except under very exceptional circumstances, such as high nervous excitement or fear of the operation. Use the ether spray

Cæsarean
section.

for the first incision, which should extend for six or eight inches from the umbilicus to within a short distance of the pubes. Make the incision exactly in the linea alba. Dissect down to the peritoneum and divide that upon the director. The parts being held aside with retractors, the uterus is now exposed. Feel on its surface with your hand for the thrill indicative of the position of the placenta, and if possible avoid that spot. Make an incision into the body of the uterus, five inches long, avoiding the fundus and the neck. As soon as the incision is made, let an assistant, with his two forefingers in the angles of the wound, hook up the uterus towards the external wound. Should the placenta be met with, thrust it aside, extract the child, and after it the membranes and placenta. If the uterus does not contract, press it with the hand, or insert a lump of ice into its cavity, or apply galvanism. Clear out all clots, and see that the os is perfectly free. Do not apply sutures to the uterine wound. Bring the parietal wound together with silver sutures passed with the curved needle, and including the cut edges of the peritoneum. During all the proceedings keep hot flannels on the abdomen and over the intestines, should they protrude. Apply broad straps of plaster across the abdomen, put on a large wad of cotton-wool, and a binder or flannel roller.

Craniotomy. Craniotomy may be required immediately, in convulsions, rupture of the uterus, or hæmorrhage, when the forceps cannot be used, or when they have failed to deliver. As little delay as possible should be made. The operation is thus performed :—Empty the bladder and rectum, and place the patient in the usual obstetric position. An assistant pressing on the fundus, pass two fingers of the left hand up to the head, and upon them guide the perforator (Fig. 88). Do not perforate through a suture, or fontanelle, but placing the point

of the instrument perpendicularly to the bone at the most dependent point of the head, push it by a gentle



FIG. 88.

semi-rotating movement, through up to the shoulder or cutting notch. Open the blades to the full extent, thus cutting an opening in the skull. Repeat this proceeding in the opposite direction, thus making a crucial opening. Thrust in the perforator, and break up the brain. Wait to see if uterine action will expel the head. If it does not, then apply the craniotomy




FIG. 89.

forceps (Fig. 89), one blade being placed within the skull and the other external to it.

In cases of arm presentation, the liquor amnii having escaped, and the child being jammed down into the pelvis, so that turning is impossible, the operation of embryulcia may be used. This consists in thrusting the perforator through an intercostal space, and cutting through the ribs, if needful, with a pair of long pointed scissors. The contents of the thorax are then evacuated. Decapitation is best performed with a long pair of sharp-pointed scissors. These are

Embryulcia.

Decapitation.

thrust into the neck, and the spinal column divided. The head being separated from the trunk, the latter is brought down, the former being afterwards extracted with the forceps.

Cephalo-
tripsy.

Cephalotripsy is an operation which may be substituted for craniotomy, but is probably easier of performance. The cephalotribe (Fig. 90) is the




FIG. 90.

instrument used. It is applied in the same way in which the long forceps are put on—the head being first perforated with the perforator. When the blades of the cephalotribe are applied, the crushing force is then commenced by slowly turning the screw. It may be needful after the first crushing to re-apply the instrument in another direction, and again crush the head. The head is then withdrawn

by the instrument, care being taken to preserve the soft parts from injury by spiculæ of bone.

The gravid uterus is subject to sudden displacements, generally retroversion or anteversion. In retroversion there is difficulty in passing water, sometimes complete retention—a frequent desire to empty the bowels, and violent bearing-down pains. On examination the cervix is found tilted forward behind the pubis, the vagina being drawn up with it out of its usual axis. Posteriorly, filling the sacrum, the globular fundus will be felt. In order to reduce the deformity, the rectum and bladder must be emptied. Place the patient on her elbows and knees, with the pelvis higher than the rest of the body, and then passing the fingers into the vagina, press the fundus upwards and forwards into its natural position. The patient must then be placed in the recumbent position and kept so for a few days, care being taken to keep the bowels freely open.

Displacements of gravid uterus.

In anteversion the os uteri is found high up towards the promontory of the sacrum, the fundus lying behind the pubes. The symptoms are much the same as in retroversion. To reduce the misplacement, empty the bladder and rectum, place the patient on her back, and then, introducing the fingers into the vagina, press the fundus upwards and backwards, at the same time hooking the cervix downwards and forwards with the fingers of the other hand.

Face presentations may be mistaken for the breech. The distinctive mark is the bridge of the nose. The malar bone and the eye may also be felt. The treatment required is not special, except that in some cases of delay it may be needful to apply the forceps.

Mal-presentations. Face.

A breech presentation is diagnosed by the plump, round feel of the buttocks, the anal aperture, and the sharp point of the coccyx. The tuberosities of the ischium may be mistaken for the malar bones, but

Breech.

Treatment.

the absence of the bridge of the nose will correct this error. The presence of meconium is no sure test, as it may occur in head presentations. The treatment is simple. The nates being born, it is better to retard rather than assist the birth of the trunk, in order that the contraction of the uterus may keep the chin well pressed down on the chest. This is the best safeguard against pressure on the cord. If a portion of the cord is seen, a loop should be pulled down to prevent stretching. If the head remains unborn, and pulsation in the cord ceases, place two fingers of the left hand on the upper jaw on either side of the nose, and press the head down, at the same time making traction on the shoulders with the fingers of the right hand. If this fail, apply the forceps without delay.

Shoulder.

A shoulder presentation has the following peculiarities:—1, The membranes project in a conical form; 2, the shoulder feels smaller and softer than the head, but larger than the elbow, and it has not the two prominences with the groove between, characteristic of the knee; 3, the scapula with its spine, the neck, and the ribs, may be felt. The shoulder is generally the first presenting part of the upper extremity, but it may be succeeded by the arm and hand. The hand is distinguished from the foot by the greater length of the fingers, the irregularity of their length, the distance the thumb may be stretched away from the fingers, the being able to close them on the palm, and the absence of the heel. The treatment is to turn as soon as the os is sufficiently dilated. If the membranes rupture before the os is dilated, this must be accomplished by steadily introducing the hand, at the same time preventing the presenting part from descending. When the os is rigid, chloroform will be of great assistance.

Arm and hand.

Hands and feet.

Hands and feet, and with them the cord, sometimes present during the progress of labour. The presenta-

tion will become one either of hands or feet alone. If possible draw down the feet, and terminate the labour as in a breech.

When the hand and head present, an attempt should be made to return the hand. If there is then delay in the descent of the head, the forceps must be applied, or turning or craniotomy resorted to.

Hand and head.

The operation of turning is thus performed:—If the right hand and arm present, pass up the right hand on the left of the pelvis, along the right side of the child until the fundus is reached. Then, if the membranes are not previously ruptured, break through them with the finger, and search for a foot. Be sure and make no mistake between a foot and a hand (see above). Seize the foot, and by gentle traction bring it down, when the arm will recede within the os. During the whole of this process, steady the uterus with the other hand on the fundus. If the left arm presents, introduce the left hand on the right side of the pelvis. The feet must be brought down over the abdomen, and not over the back. Protect the cord from pressure by pushing it aside to one or other sacro-iliac synchondrosis. If the head is long in being born, apply the forceps.

Turning.

Prolapse of the cord requires either its immediate replacement within the uterus, or the delivery of the child. The first may be accomplished by raising the patient's hips very high on pillows, the waist being on a much lower level. The cord may then by the force of gravity slip back into the uterus, and a pain will force down the head, and prevent the re-prolapse. If this does not succeed, either turning or the forceps must be used. No steps need be taken if the pulsation in the cord has ceased, and the stethoscope still further confirms the death of the child.

Prolapse of the cord.

Puerperal convulsions present all the ordinary features of epileptic convulsions. The treatment is:—

Puerperal convulsions.

1, Evacuate the contents of the bowels freely with scammony and jalap; 2, shave the head, and apply ice or evaporating lotions; 3, empty the bladder, or at any rate pass a catheter to be sure that it is empty; 4, wait until the os is sufficiently dilated, and then apply the forceps and deliver the child as soon as possible. Chloroform should be given, both to allay the violence of the convulsions, and also to facilitate the delivery; 5, give chloral hydrate in thirty-grain doses. If the patient is unable to swallow, administer it by the rectum. If the uterus is rigid, and the labour does not advance, the convulsions being constant, perform craniotomy.

Rupture of
the uterus.

The symptoms of rupture of the uterus are:—1, Violent and continued uterine action without corresponding descent of the child; 2, sudden agonising pain of a crampy character; 3, a sensation of something having given way; 4, cessation of all uterine action; 5, sickness, clammy sweats, small pulse, and anxious expression; 6, recession of the child beyond the reach of the hand, its limbs being felt through the abdominal walls; 7, hæmorrhage from the vagina; 8, vomiting of a coffee-ground appearance. The treatment is at once to deliver the child. If the head can be reached, the forceps may be applied; if not, the hand and arm being passed up into the uterine cavity, an attempt to turn must be made. If this cannot be done, it is better at once to perform the Cæsarean section as the best chance of saving the life of the child. Very free administration of stimulants throughout is needful.

Treatment.

Rupture of
the bladder.

Rupture of the bladder presents many of the symptoms of rupture of the uterus; but, 1, There is no hæmorrhage from the vagina; 2, no recession of the child, or indication of its presence in the abdomen; 3, uterine action is not arrested. The belly, however, becomes rapidly swollen and tender. The treatment

Treatment.

is, first of all, as rapidly as possible, to deliver the child by the forceps or turning. A gum-elastic catheter must then be introduced into the bladder, and retained. One of the soft india-rubber catheters would be the best, if at hand. The patient must be placed under the influence of opium as rapidly as possible, and turpentine stupes applied to the abdomen.

Rupture of the perinæum is often the result of too speedy delivery, especially when the forceps are used. It is of little consequence, unless the sphincter ani is torn through. If this be the case, the better plan is at once to bring the parts together with interrupted sutures. It is not needful to use the quilled suture. Pass three or four sutures of thick silk through the deep parts with a curved needle set in a handle, and some superficial ones of finer silk, with an ordinary needle, should be inserted between. The patient should be kept on her side, with the thighs tied together. The water must be drawn off with a catheter, and the bowels confined for a few days.

Rupture
of the
perinæum.

Inversion of the uterus is sometimes the result of precipitate labour, and sometimes it is spontaneous, occurring after labour, or as the result of violent coughing, or straining of any kind. The symptoms are:—1, Great pain, of a dragging, bearing-down character, with a sense of nausea and sinking; 2, a round, hard tumour is felt per vaginam, whilst, on examining the abdomen, the uterus cannot be felt. The treatment is to return the womb to its proper position as soon as possible. To accomplish this, firmly grasp the uterus, so as to empty it of blood, at the same time pressing firmly on the fundus with the palm of the hand. As the organ returns, follow it with the hand, which should not be withdrawn until the uterus firmly contracts upon it. If the placenta is attached, it must be removed before reduction is attempted.

Inversion
of the
uterus.

Treatment.

CHAPTER X.

POISONING.

Poisoning by Sulphuric Acid—Nitric Acid—Oxalic Acid—Salt of Sorrel—Hartshorn—Phosphorus—Arsenic—Scheele's Green—Orpiment—Sugar of Lead—Blue Vitriol—Tartar Emetic—Chloride of Zinc—Cantharides—Mussels—Opium—Godfrey's Cordial—Dalby's Carminative—Paregoric Elixir—Prussic Acid—Cyanide of Potassium—Essential Oil of Bitter Almonds—Chloroform—Tobacco—Strychnia—Hemlock—Water-Hemlock—Monkshood—Deadly Nightshade—Fungi—The Use of the Stomach-pump.

Sulphuric
acid.

POISONING by strong sulphuric acid is accompanied by intense burning pain in the throat, gullet, and stomach, vomiting of coffee-grounds mixed with shreds of mucus and blood. Treatment:—The administration of any alkali at hand, such as lime or magnesia, and of demulcent drinks.

Nitric acid.

The symptoms of poisoning by nitric acid are the same as those above, and the remedies are also the same.

Oxalic acid.

Oxalic acid is sometimes taken in mistake for Epsom salts. The symptoms are, burning sensation in the throat, great collapse, and vomiting of dark green matter. Treatment:—The administration of copious draughts of warm water, and of chalk or magnesia, or for promptitude the scrapings from the ceilings or white-washed walls, made into a creamy paste. The stomach-pump should be applied as early as possible. Salt of Sorrel, or essential salt of lemons, is used for bleaching straw and removing

Salt of
Sorrel.

ink stains ; it consists of acid oxalate of potash, and, if taken, produces all the symptoms of oxalic acid poisoning, and must be treated in the same way.

Poisoning by "hartshorn and oil" is not uncommon. The symptoms are, intense burning sensation in the throat, gullet, and stomach, and, when vomiting occurs, which is not always present, the vomited matter is mixed with blood of a dark brown colour. The immediate remedy is to give vinegar, lemon-juice, and demulcents.

Ammonia
or harts-
horn.

Phosphorus paste is sometimes taken as a poison. The symptoms of phosphorus poisoning are those of an irritant poison. There is intense thirst, nausea, severe pain, and an odour of garlic in the breath and vomited matter. The treatment is to administer emetics, and give magnesia.

Phosphorus.

Various powders for the destruction of vermin, etc., contain arsenic, and are frequent sources of poisoning. The symptoms come on about half an hour after the dose has been taken, and are—nausea and burning pain in the stomach, violent vomiting, intense thirst, purging, and pain in the epigastrium on pressure. The treatment is to excite vomiting as much as possible by the administration of a scruple of Sulphate of Zinc and Ipecacuanha and plenty of warm water, and to apply the stomach-pump as soon as possible. The Hydrated Peroxide of Iron, given in large quantities, is also useful, and it may be readily prepared by mixing two or three ounces of Liquor Ferri Perchloridi B.P., with about a quarter of an ounce of Magnesia B.P. A mixture of linseed-meal, castor-oil, and water, made to the consistency of treacle, may also be given to protect the stomach.

Arsenic.

Scheele's Green, or arsenite of copper, is met with in a variety of pigments and paper-hangings. Symptoms of poisoning by it are those of arsenical poisoning, and must be treated in the same way.

Scheele's
Green.

Orpiment. Orpiment, or sulphide of arsenic, is also employed in the arts as a yellow colouring, and may produce arsenical poisoning.

Corrosive sublimate. Corrosive sublimate, or Perchloride of Mercury, when taken in poisonous doses, produces a metallic taste in the mouth, pain, vomiting of white matter, and profuse diarrhœa. Treatment:—Give white of eggs, followed by infusion of galls or catechu, and use the stomach-pump as soon as possible.

Sugar of lead. Sugar of Lead, or Acetate of Lead, is frequently taken as a poison, although it is seldom fatal except when given in very large doses. The symptoms are:—Pricking sensation in the throat, dryness, thirst, vomiting, and colic. Treatment:—Emetics of Sulphate of Zinc, diluents, and Sulphate of Soda and Magnesia, and Phosphate of Soda.

Blue vitriol. Blue vitriol, or Sulphate of Copper, when taken, causes such severe vomiting as to be an antidote to itself. When this is not effectual, the symptoms are, headache, abdominal pain, purging, and sometimes convulsions. Treatment:—Encourage vomiting with warm water, and give milk, flour and water, and coffee; white of egg also is useful.

Tartar emetic. Tartar emetic has been taken in mistake for Epsom salts. The symptoms are, collapse, with livid countenance, violent pain in the stomach, spasmodic contraction of the muscles, particularly of the arms and abdomen, violent vomiting and purging. Treatment:—Excite vomiting, and use the stomach-pump; give infusions of gall-nuts or cinchona, or strong tea.

Chloride of zinc. "Sir W. Burnett's Fluid," which is a solution of chloride of zinc, is not unfrequently the cause of poisoning. The symptoms are, immediate pain and vomiting, burning pain in the gullet, and collapse. Diluents must be freely given, and milk, white of egg, and Sulphate of Soda and Magnesia. The fatal effects are generally of a secondary character.

Cantharides, either in powder or tincture, is sometimes administered in poisonous doses for criminal purposes ; one or two drachms of the powder will cause burning in the throat, pain in the abdomen, vomiting of bloody mucus, incessant desire to pass water. The treatment must consist of emetics and demulcents. Cantharides.

Poisonous mussels produce uneasiness and weight in the stomach, numbness in the extremities, dryness in the throat, cramps in the limbs, swelling of the eyelids, and eruption of nettle-rash, with great exhaustion. Emetics must be freely given, and diffusible stimulants, with opiates to allay pain. Mussels.

In poisoning by opium the symptoms are :—Stupor, from which the patient may be roused, but at once relapses ; pulse small and irregular ; skin warm and livid ; pupils contracted in the early stage, but dilated at a later period ; occasionally the odour of opium in the breath. The symptoms usually commence about half an hour after the poison has been taken. Treatment :—Immediate use of the stomach-pump, Zinc emetics, animal charcoal, infusion of galls or Tannic Acid, cold affusion to the head, and strong coffee. The subcutaneous injection of atropine is also useful. One drop of the *Liquor Atropiæ B.P.* contains the $\frac{1}{160}$ of a grain. At least two drops of this solution may be injected. The patient must be kept in continual motion, the surface of the body being flapped with wet towels. Lastly, when all else fails, artificial respiration should be kept up for a considerable time. Opium.

Godfrey's Cordial is made of infusion of Sassafras, Treacle, and Tincture of Opium. It contains about one drachm of the latter in six ounces, or half a grain of Opium in an ounce. Half a teaspoonful has been known to cause the death of an infant. Godfrey's
Cordial.

Dalby's Carminative is composed of essential oils, aromatic tinctures, Carbonate of Magnesia, and Tinc- Dalby's
Carminative.

ture of Opium. It contains one-eighth of a grain of Opium in every ounce. Forty drops have been known to kill an infant.

Paregoric
elixir.

Paregoric Elixir is a preparation of Alcohol, Opium, Benzoic Acid, and Oil of Anise and Camphor. It contains one grain of opium in every half ounce. Half a teaspoonful has been fatal to an infant.

Prussic acid.

Prussic acid when taken in a large dose is generally immediately fatal. In smaller doses (about thirty minims of the weak acid) the symptoms are, weight and pain in the head, giddiness, nausea, quick pulse, loss of muscular power, foaming at the mouth, strong odour of bitter almonds. Treatment:—Cold affusion to the head and spine, Carbonate of Ammonia internally, inhalation of chlorine gas, or small doses of Chloride of Lime or Soda. The chemical antidote, if there is time for its administration, is the moist Peroxide of Iron.

Cyanide of
potassium
and essen-
tial oil of
bitter
almonds.

Cyanide of potassium, much used by photographers, when taken produces all the symptoms of poisoning by prussic acid; and the treatment is the same. So also the essential oil of bitter almonds, used as a flavouring, has been known to destroy life in a dose of seventeen drops.

Alcohol.

The symptoms of poisoning from Alcohol come on in the course of a few minutes, and soon proceed to coma. Insensibility may, however, not come on for some time, and then suddenly. Intoxication, concussion of the brain, and poisoning by Opium are likely to be confounded. Usually in intoxication the smell of the breath is decisive. If this is absent, it may be presumed that the case is not one of intoxication. In alcoholic poisoning the face is generally flushed and the pupils dilated; in opium poisoning the face is pale and the pupils generally contracted. The temperature in alcoholic poisoning is generally below normal. The contents of the stomach will usually

reveal the true state of affairs. Treatment :—Immediate use of the stomach-pump, cold affusions to the head, and artificial respiration if the breathing is affected.

In threatened death from chloroform, the tongue should be immediately pulled forwards with hooked forceps and the epigastrium forcibly pressed up; artificial respiration by Sylvester's method (see p. 43) should be resorted to without delay, and galvanism: the patient should be placed in a current of cold air, and the chest and extremities flapped with a wet towel. The head, if the face is pale, should be lowered; if turgid, should be raised. Artificial respiration should be kept up for a long period, even after all appearance of life has ceased.

Chloroform.

The symptoms of poisoning with Tobacco are, faintness, vomiting, giddiness, delirium, coldness of the surface, and convulsions. The treatment is the administration of stimulants, strong coffee, and the application of warmth to the surface.

Tobacco.

Poisoning with Strychnia shows itself at variable periods after the dose has been taken. The sixteenth of a grain has been known to kill a child, and half a grain an adult. The symptoms are, twitchings of the head and limbs, tetanic convulsions, opisthotonos, livid face, with complete retention of the intellectual faculties. The patient either dies or recovers within two hours of the commencement of the attack. The treatment is to inject into the stomach large quantities of powdered charcoal suspended in water. If the spasms have set in, chloroform should be given, or large doses of chloral.

Strychnia.

Hemlock, or *Conium maculatum*, has been taken in mistake for parsley. The symptoms are, loss of power in the upper and lower extremities and of deglutition, partial paralysis of sensation, and fixed pupils. The two plates, Figs. 91, 92, exhibit the

Hemlock.



FIG. 91—HEMLOCK.



FIG. 92—PARSLEY.

difference between the leaves of hemlock and parsley.


Water-hemlock.

The water-hemlock has given rise to fatal accidents,

its roots having been mistaken for parsnips. The symptoms are, dimness of sight, difficulty of breathing, burning pain in the stomach, vomiting and convulsions.

Monkshood, or *Aconitum Napellus*, has frequently been the cause of poisoning, the root being mistaken

Monkshood.



ROOT OF HORSERADISH.

ROOT OF ACONITE.

FIG. 93.

for horseradish. Fig. 93 shows the difference between

the two roots. The symptoms of poisoning are, tingling and numbness in the throat and limbs, frothing at the mouth, severe pain in the stomach, vomiting and purging, dimness of sight, dilated pupils, and livid skin.

Deadly
Nightshade.

Death has sometimes been caused by partaking of the berries or the leaves of Deadly Nightshade, or *Atropa belladonna*. The symptoms are, hot and dry throat, flushed face, dilated pupils, staggering gait, loss of hearing and power of speech. The treatment in all these cases consists in the administration of large quantities of powdered animal charcoal, suspended in water, and after that a good Zinc emetic.


Fungi.

The treatment of poisoning by Fungi is to produce vomiting by tickling the fauces with a feather, and to inject one drop of the solution of Atropine hypodermically, repeating the injection in short intervals if required.

The use of
the stomach-
pump.

The stomach-pump should be used as follows:— Bend the extremity of the long tube rather abruptly at its point, and oil it; pass it to the back of the throat, and when the tube touches the back of the pharynx, bend the patient's head a little forward; slight pressure will then make the tube pass on to the back of the larynx, when it will be arrested, but gentle pressure will overcome the difficulty. When the tube has reached the stomach, inject some water; in order to do this, fill the pump with water from a basin by steadily drawing out the piston (Fig. 94), the lever connected with the stop-cock being not pressed down. The pump being filled with water, press down the lever with the thumb of the left hand, and gently press in the piston with the right, thus propelling the water into the stomach. When enough has been injected reverse the action. If the tube becomes blocked, again reverse, and so eject the fragment blocking the eye of the tube. If the

patient is unconscious or unruly, the mouth must be opened with a screw-gag, and kept open with a



wooden gag supplied for the purpose with the stomach-pump.

A far simpler and less expensive form of stomach-pump is that introduced to the profession by Mr. Tossell of Exeter.

The long flexible india-rubber tube is passed to the

back of the mouth, and then its passage down the œsophagus is effected by twisting it round and round




FIG. 95.

in the hand until it reaches the stomach. One end of the tube is then fixed to the apparatus, which consists of a Higginson's syringe without the valve, and the other end is placed in a jug of water. Then the tube is pinched up by the finger and thumb at a point between the ball and the mouth, the ball being at the same time compressed by the other hand and the air in it expelled. The ball then becomes filled with water, and on raising the jug above the patient's head and discontinuing the pinching of the tube, the water flows in a continuous stream into the stomach. When enough has been introduced, the tube must again be pinched between the ball and the mouth, the end of the tube taken out of the jug and placed in a basin, below the level of the stomach, when upon discontinuing the pinching of the tube, the fluid will flow in a continuous stream from the stomach into the basin. Should the stomach tube become choked, it can be freed by pinching the tube between the ball and the jug, and compressing the ball with the other hand, when the fluid contained in it will be pressed out and the tube freed.

CHAPTER XI.

ANTISEPTIC TREATMENT.

General Remarks—Carbolic Acid—Antiseptic Gauze—Gauze Bandage—Antiseptic Atmosphere—Steam Spray Apparatus—Boric Acid—Salicylic Acid—Chloride of Zinc—Iodoform—Oil-Silk Protective—Drainage Tube—Catgut Ligatures—Carbolized Silk Sutures—Abscesses—Precautions before and during an Operation—Dressing after Operation—Compound Fractures—Dressing in Compound Fractures—Operation on Parts affected with Sinuses—Abscess by the Side of Rectum—Superficial Sores.

THE mere use of antiseptic dressings will not of itself yield successful results to the surgeon. He must give unremitting attention to all the minutiae of application which have been deduced from the germ-theory of putrefaction, upon which the treatment is based. The results of true antiseptic management will then far more than repay the effort required for their attainment; to say nothing as to the average saving of time, and the relief derived from diminished anxiety.

The following is a short account of the antiseptic treatment employed by Professor Lister in 1880:—

The five antiseptics which have been found most useful are Carbolic Acid, Boric Acid, Salicylic Acid, Chloride of Zinc, and Iodoform.

Pure carbolic acid, remarkable for its power of destroying low forms of life, is, on the whole, the most efficient antiseptic for general purposes. Its energetic action on the epidermis makes it the best

Carbolic
acid.

agent for cleansing purposes. In the form of one to twenty watery solution, it is used for purifying the integument of the part to be operated upon, and the sponges, instruments, etc. As a one to forty solution, it is used for washing sponges during an operation, for the hands of the operator and assistants, and for the changing of dressings.

Antiseptic
gauze.

The volatility of carbolic acid renders it invaluable for dressing hollow wounds and abscesses. It is the active constituent of the ordinary dressing—antiseptic gauze¹—which is applied in eight layers, of size proportioned to the expected amount of discharge. A piece of thin mackintosh cloth (technically called hat-lining) is interposed beneath the outer (eighth) layer of gauze, to prevent the discharge from soaking directly through the centre of the dressing, washing out the portion of acid there stored up, and thus giving putrefactive organisms direct access to the cavity. Further, as carbolic acid is given off very slowly by the gauze at the ordinary temperature of the air, it is quite possible that septic organisms, accidentally adherent, might be conveyed to the wound by the surface of the dressing itself. To guard against this mishap, a small piece of gauze wrung out of the promptly-acting one to forty solution is applied before the ordinary eight-fold dressing.

Gauze
bandage.

The antiseptic gauze is convenient in the form of bandage, which not only applies itself, and retains its position better than a calico bandage, but is often a valuable addition to the antiseptic quality of the dressing. In the case of stumps, in which there is a tendency to retraction of the soft parts, the antiseptic bandage enables the surgeon to overcome what would otherwise be a considerable difficulty in the treatment of the case.

¹ The gauze is now prepared with one part of pure carbolic acid to four parts of resin and four parts of pure paraffin.

It is not sufficient to have a reliable dressing, and to be able to purify the skin and the instruments. There must also be an antiseptic state of the atmosphere, which cannot fail to gain access to the wound or abscess during the operation, or the changing of dressings. This is provided in the form of a spray of about one to thirty carbolic acid solution, for which a steam spray apparatus is the most easily managed and efficient instrument at present available.¹

Carbolic
spray.

The steam spray apparatus should produce a large and finely-divided cloud of vapour, and there should be an arrangement for altering the incidence of the spray and for regulating the size of the flame. The boiler should not have the aperture for filling placed at the summit of the dome, but at a lower level; otherwise the boiler may be over-filled, with the result that when the steam is turned on there will be a jet of scalding water ejected, or there may be blocking of the steam point by solid particles washed into it by the water. After use the boiler should be carefully emptied, and during the emptying it should be so held that the water cannot flow into the steam-pipe.

Steam
spray
apparatus.

When the spray is suspended during an operation or the changing of a dressing, the wound should be covered with a piece of sound calico moistened with one to forty carbolic acid solution, which, for convenience, is termed the guard. Of course, neither spray nor guard is required for superficial sores.

The guard.

Boric acid is a valuable antiseptic, but its non-volatility prevents its being used for the dressing of hollow wounds, and for the spray. It is bland and unirritating as compared with carbolic acid, and is

Boric acid.

¹ The bottle of the spray apparatus is filled with filtered one to twenty carbolic acid solution, which, mingling with the steam, yields a vapour consisting of one part of carbolic acid to about thirty parts of water.

therefore particularly useful as a dressing for superficial wounds and sores. It is employed in the form of a saturated watery solution ; as boric lint, which contains about half its weight of the crystals of boric acid ; and as an ointment which may be prepared by mixing one part of powdered boric acid with five or ten parts of vaseline, according to the strength desired.

Salicylic
acid.

Salicylic acid, which was introduced into surgical practice by Professor Thiersch, of Leipzig, at the suggestion of Professor Kolbe, has not been adopted by Mr. Lister as an ordinary dressing, having been proved by him to be inferior to carbolic acid in its power of destroying bacteria, though very efficient in preventing chemical fermentations. He has, however, found salicylic acid to be of great use in certain exceptional cases, where irritation of the skin occurs after a gauze dressing has been left on for several days. The form of eczema thus arising was formerly attributed by Mr. Lister to the irritating effect of carbolic acid on sensitive skins, but the invariable disappearance of the affection after one or at most a few applications to the inner layer of the gauze dressing of a small quantity of salicylic acid suspended in one to forty carbolic acid solution has shown the irritation to be due to an alteration in the effused serum brought about by some other agency.

Chloride of
zinc.

Chloride of zinc has the remarkable peculiarity among antiseptics of producing such an effect upon the tissues of a recent wound, when applied in a watery solution of about forty grains to the ounce, that, as the result of a single application, the cut surface, though not presenting any visible slough, is rendered incapable of putrefaction for two or three days, notwithstanding its exposure to the influence of septic material. The patient is thus tided over the dangerous period preceding granulation, during which

the divided tissues are most prone to inflammation and the absorption of septic products. Hence this salt, though very irritating and non-volatile, and therefore not adapted for general use, is of the highest value in circumstances which render it impossible to exclude septic organisms in the after-treatment; as, for example, after the removal of tumours of the jaws, or operations about the anus, and after amputations or excisions in parts affected with putrid sinuses. The solution is freely applied to all textures, including bones, by means of a piece of lint held in dressing forceps. Sinuses are injected at the conclusion of the operation by means of an ordinary brass syringe.

Iodoform, which has the advantage over chloride of zinc of not causing pain, has lately been sprinkled over the surface of putrid ulcers, in order to purify them prior to the application of the antiseptic dressing. Iodoform.

In order to obtain the best results from the antiseptic treatment, it is desirable to protect the healing parts from the irritating influence of the antiseptic itself, which, in the case of carbolic acid, would often prevent cicatrisation altogether, and which, even with the much blander boric acid, is better avoided. This protection is effected by interposing a layer, consisting of some material unirritating in itself, and as far as possible impervious to the antiseptic agent.

The protective commonly used is composed of thin oil-silk, varnished with copal and then coated with a layer of dextrine. The dextrine allows the oil-silk to become uniformly wetted by the antiseptic solution, into which it is dipped at the moment of application. It is obvious that the antiseptic dressing proper must extend a considerable distance beyond the protective, or putrefaction may spread under the Oil-silk protective.

latter in spite of the dressing which is above it. A thoroughly aseptic sponge wrung out of the one to twenty carbolic acid solution is often a useful addition to the first dressing when uniform pressure is desired after an operation. The sponge should be placed above the protective and beneath the large eight-fold (or outer) dressing.

The
drainage
tube.

The application of antiseptics to the raw surfaces of a wound causes a certain amount of irritation, and leads to an unusually abundant discharge of serum; so, in order that this may not be pent up, and occasion inflammation from tension, a very free outlet must be provided. This is effected by means of Chassaignac's drainage tubing, of which the red is preferable to the black variety. Four sizes are used, having an external diameter of $\frac{1}{3}$, $\frac{1}{4}$, $\frac{1}{5}$, and $\frac{1}{8}$ inch respectively. Large holes are cut in the walls to allow the discharge to reach the interior of the tube; and two loops of silk thread, knotted at the ends, are attached to opposite sides of the margin of the outer extremity of the tube to prevent its displacement. The outer end is cut either transversely or obliquely, according to the direction of the cavity, so that the orifice of the tube may be flush with the integument.

When abscesses are opened, a drainage tube should be used, of a length equal to the depth of the cavity, and of a calibre proportioned to the quantity and thickness of the pus discharged. The tube should be shortened from time to time as the cavity becomes smaller.

Horsehair which has been soaked for a few minutes in one to twenty carbolic acid solution constitutes an efficient drain for operation wounds etc. As the discharge of serum diminishes, the drain may be lessened by removing a few hairs at each dressing. The drain may be partially drawn out of the wound

and shortened when expedient. In large operation wounds it is often desirable to use both drainage tubes and drains of horsehair. In these circumstances the tubes are discontinued before the horsehair.

For arresting arterial hæmorrhage, the prepared catgut used antiseptically combines the security and universal applicability of the ligature with the virtual absence of any foreign body from the wound. The vessel is tied in the usual way, but the ends of the catgut are cut off near the knot.

The catgut may be conveniently carried in the pocket-case in one of Mr. Lister's catgut-holders. As the right preparation of the catgut is a matter of very great importance, it is best to obtain it ready prepared along with the other antiseptic materials.

As metallic sutures are liable to catch in the dressings, carbolized silk sutures are used with advantage. They are prepared by passing ordinary ligature silk through melted beeswax, containing ten per cent of carbolic acid. The superfluous wax is wiped off the thread as it is drawn out of the mixture. The antiseptic property of the silk is conveniently retained by keeping it in a stoppered bottle or wrapped in antiseptic gauze.

Metallic sutures are used with lead buttons to constitute "stitches of relaxation" when the ordinary sutures are subjected to much tension. Horsehair is very useful as an intermediary suture when accurate apposition of the edges of a wound is desired. Horsehair sutures are serviceable for stitching the mucous membrane to the skin in plastic operations on the lips, etc. Long hairs from the tail of a stallion are the best for surgical work. The prepared catgut may be used for sutures or drains in cases where its rapid absorption is not likely to be disadvantageous.

DIRECTIONS FOR SPECIAL CASES.

I. *Abscess.*

Abscess.

This is the simplest case for antiseptic treatment, the antiseptic not being applied at all to the affected tissues, but being used merely to prevent the access of septic mischief from without.

Carefully select the spot for incision with special attention to its distance from a source of putrefaction,¹ so that the dressing may extend freely beyond the incision in every direction. The skin, the knife, and the operator's fingers having been cleansed with one to twenty carbolic acid solution, after shaving off hairs from the integument, if necessary, let the spray play effectively on the part, then make an incision in the selected position, and of such a size as will easily admit the drainage tube; press out the pus;² insert the drainage tube, and extend the loops of silk in opposite directions; apply over the incision a small bit of gauze wetted with carbolic acid solution; cover this with a large dressing consisting of eight folds of gauze and the enclosed piece of mackintosh, as above described; and lastly, secure this with a well-adjusted gauze bandage and with safety-pins, which must not penetrate the centre of the mackintosh, or its efficiency will be destroyed. When the abscess is situated in a part where the dressing is particularly liable to disturbance from the movements of the patient (as at the groin, in case of psoas abscess), great additional security is afforded by the employment of a turn or two of elastic webbing applied over the edges of the dressing, but not so tightly

¹ So important is this that it might lead to the refusal to operate, *e.g.*, in a case of spinal abscess pointing near the anus.

² In strumous abscesses it is often advisable to apply a sharp spoon to the walls of the cavity.

as to occasion uneasiness or œdema of the limb. This elastic bandage follows the movements of the patient, and insures the constant application of the dressing to the skin. If it is desirable to explore the abscess cavity with the finger, special care must be taken that the latter be well cleansed about the nail and elsewhere with carbolic acid solution, immediately before introduction.

Next day let a hand be placed so as to keep the dressing in position whilst the pins are being removed and the bandage is being cut. Direct the spray towards one side of the dressing, which must then be cautiously raised, so that the spray may pass into the angle between the dressing and the skin. Remove the drainage tube under the spray;¹ cover the incision with a guard; wash the drainage tube and the skin with one to forty carbolic acid solution; again direct the spray to the incision, remove the guard, insert the tube, and dress as above described. Re-dressing.

As the discharge diminishes, dress the case every two, three, four, five, or six days as required, the general rule being that the dressing is changed, if discharge is found to have made its appearance beyond it at the usual time of the surgeon's visit. Give the same amount of care to every dressing until cicatrization is complete; this being the only trustworthy evidence that the sinus is closed. In cases of chronic abscess, due to caries of vertebræ, or to morbus coxæ, keep the patient in bed six weeks after the abscess is closed. In the latter cases, keep on the long splint during the whole period.

II. *Operations on parts where the skin is unbroken.*

These cases come next in the order of simplicity

¹ Usually it is best to leave the drainage tubes undisturbed during the first three or four days, otherwise there may be difficulty in reinserting them.

of treatment, the essential object being, as in abscesses, to prevent the introduction of septic organisms during the operation; although the divided tissues are necessarily more or less exposed to the influence of the antiseptic in the spray and the sponges, etc.

Prepare the skin, the hands of the operator, and the instruments as above; conduct the operation as usual, but keep the spray at work until the dressing has been applied, of course using the guard if the spray be suspended.

When the hand of the operator or assistant has been withdrawn from the spray, and when a sponge or an instrument has been laid aside during the performance of an operation, it is an essential point, very apt to be overlooked, that before being reintroduced into the wound, the hand or sponge or instrument should be again thoroughly cleansed with the carbolic acid solution.

After securing the vessels, adjust a drain, and stitch up the wound; then dip into the solution a piece of protective, large enough just to cover the incision, and lay it upon the wound; over this apply a piece of gauze wetted with the solution, a sponge if required,¹ and then the usual well-overlapping dry gauze dressing. Change the dressing next day, and afterwards dress daily, or less frequently, according to the amount of the discharge. As a rule, the dressing should not be left on longer than a week, or putrefaction might occur after the evaporation of the carbolic acid from the gauze.

Amputa-
tion. -

In amputations apply a gauze dressing large enough to wrap round the limb, and to turn up over the face of the stump, like the wrapping at the end of a parcel.

In primary amputations, and in amputations through limbs above the seat of putrid discharge,

¹ It is often useful to secure the deep dressing by means of one or two turns of gauze bandage.

cover the contused or putrid parts with a towel wrung out of one to twenty carbolic acid solution, so that the sound parts may not become contaminated during the operation.

III. *Accidental wounds, including compound fractures and dislocations which do not require amputation.*¹

Here, opportunity having been afforded for the entrance of septic material either in the infliction of the injury, or subsequently before the case is seen by the surgeon, the antiseptic must be applied to all the recesses of the wound, in order to counteract the mischief which has so entered. For this purpose a strong antiseptic solution must be used, but while it must be introduced among the blood-clots in the recesses of the wound, it is very undesirable that it should pass unnecessarily into the cellular tissue. In ordinary circumstances the one to twenty carbolic acid solution is employed, but instead of forcing it into the wound generally, by means of a syringe, a better plan is to apply it to the various irregularities of the injured part by a flexible catheter adapted to the syringe with caoutchouc tubing. When many hours have passed since the infliction of the injury, Mr. Lister has for several years used a still more powerful solution, viz. one part of carbolic acid to five parts of methylated spirit.

Wounds and compound fractures.

These operations are conducted under the spray, after cleansing the skin with the one to twenty carbolic acid solution. If a fragment or an end of a bone is projecting, it is well washed with the same solution and returned, in some cases after sawing off its extremity; but reduction is never effected until the depths of the

¹ Experience will soon teach the surgeon that he may save, under the antiseptic treatment, many cases that would require amputation without it.

wound have been injected, and, of course, loose fragments and foreign bodies extracted.

It is difficult to assign the precise limit of time when antiseptic treatment must be regarded as hopeless in such injuries. Success has sometimes been obtained as late as thirty-six hours after the accident.

The introduction of stitches should always be avoided in this class of injuries, whilst in some cases a drainage tube should be inserted; and if the wound is very small, it should be enlarged to prevent tension from the accumulation of serum.

The dressing is then applied, as in the last class of cases, the folded gauze being made to envelop the limb for a considerable extent above and below the injured part, in accordance with the large amount of bloody and serous discharge to be anticipated. A gauze padding for the splints in the vicinity of the injury is also often valuable as an additional precaution.

The dressing should always be changed on the following day; but afterwards, if all proceeds satisfactorily, the intervals between the dressings are to be increased as the discharge diminishes. It must, however, be borne in mind, that from the very nature of these cases, as some organisms in the recesses of the wound may occasionally elude the action of the antiseptic, success cannot be reckoned on with certainty as in the two former classes.

IV. *Operations through parts affected with putrid sinuses.*

Putrid
sinuses.

These cases present the greatest degree of complexity in the antiseptic treatment. It is hardly to be expected that the putrefaction which exists in the recesses of the sinuses can always be completely eradicated; but in a certain proportion of cases this is effected by the practice recently adopted of scrap-

ing away all granulation material from the tracks of the sinuses and elsewhere by means of the sharp spoon, after the example of Professor Volkmann, of Halle, and injecting the solution of chloride of zinc at the conclusion of the operation.¹ The chloride of zinc solution is at the same time applied to the cut surfaces, and by this means the patient is saved a great deal of pain and danger, as above alluded to, even should putrefaction ultimately show itself.

The usual external antiseptic dressing is employed in the hope of an aseptic result, while it also protects the atmosphere of a hospital ward from foul emanations in case putrefaction should occur.

In these cases of the recurrence of putrefaction, a dressing of boric lint, or of boric ointment, may often be advantageously substituted for the gauze dressing.

V. *Abscess threatening fistula in ano.*

This particular kind of abscess is of such importance as to deserve special notice here. In consequence of the vicinity of the anus preventing the skin around the wound from being overlapped by the dressing to a sufficient extent to insure absence of putrefaction, the gauze cannot be employed. But the difficulty may be overcome by applying to the anal orifice and surrounding parts a mass of lint soaked with a solution of carbolic acid in ten parts of olive oil, to be retained in position by a T-bandage; the patient being directed to draw the pad of lint slightly towards the affected side before defæcation,

Abscess
near anus.

¹ At one time Mr. Lister recommended the injection of the sinuses before the operation was commenced; but though this is the most effective mode of introducing the solution, it has been abandoned for several years in consequence of inconveniences having in some cases followed from the solution becoming diffused into the cellular tissue of the part.

so that the evacuation may be effected without exposing the wound, and to cleanse the anal outlet with the oily solution before readjusting the pad. The oiled lint is changed about every four hours during the first day ; but when the discharge becomes very slight, every twelve hours is sufficiently often. The abscess is opened under the spray as usual, after washing well with the one to twenty watery solution of carbolic acid ; and, instead of a drainage tube, a bit of lint soaked in the carbolized oil is inserted to prevent primary union. Fœtor of the pus on evacuation of the abscess must not discourage the surgeon from continuing the antiseptic measures, for experience has shown that in such circumstances, as well as when the pus is odourless, the usual results of the antiseptic system applied to abscesses may be attained, all further suppuration ceasing, and fistula being averted.

VI. *Superficial Sores.*

If there are sloughs, boric lint moistened with the boric acid solution is often employed, after the manner of the water dressing. When the sloughs have separated, the granulating surface is dressed with protective and boric lint, or with boric ointment spread on thin calico. In either case the dressing is to be dipped into the solution before it is applied. In cases of putrid chronic ulcer, the integument is cleansed with one to twenty carbolic acid solution, iodoform is sprinkled over the surface of the ulcer, to which is then applied a dressing of protective and boric lint. The application of iodoform may be repeated at the next dressing should it be found that putrefaction has not been eradicated by the first application.

CHAPTER XII.

APPARATUS AND DRESSINGS.

Extemporary Apparatus—Conveyance of injured persons—Esmarch's Cloth—Swing for patients in bed—Pasteboard Splints—Plaster of Paris—Gum, Chalk, and Starch—Glue—Gutta-percha Splints—Extension by Weight—The Use of the Aspirator—Transfusion of Blood—Esmarch's Bloodless Operation—The Elastic Tourniquet—Hunter's Tourniquet—Method of Opening Deep-seated Abscesses—Counter-Openings—Burns and Scalds.

EXTEMPORARY splints may be manufactured on the moment from various substances nearly always to be found in the neighbourhood of the accident. For fractures of the lower extremities two bundles of reed cut the length of the limb and rolled up in a coat or rug, one on either side of the limb and secured by two or three straps, make an admirable splint. On the injured limb being secured to its fellow both are placed on a back-board padded with some material.

For a fractured thigh place a long splint (the bar of a gate) on the outer side from the hip to the ankle and another splint on the inner side, a short one being placed in front of the thigh, and secure all with straps or handkerchiefs. The injured limb should also be secured to its fellow. For fractures of the upper extremity, splints of reed, of pasteboard, or made from an old hat or bonnet-box, will serve the temporary purpose. Stirrup leathers, bridles, and traces may be utilised in various ways.

Conveyance
of injured
persons.

Various stretchers may be extemporised for conveying an injured person home or to hospital. Take a blanket and have a loop sewn on at each of the four corners. Bring the two loops at each end together and pass a long pole through the four loops. A ladder may sometimes be found useful, or the stretcher here figured from Surgeon-Major Porter's "Surgeon's Pocket Book" may easily be extemporised if the materials are at hand (Fig. 96).




FIG. 96.

The bearers of a stretcher should not walk in step. That is to say, that if the front man steps off with the left foot the rear man should step off with the right. They should march with a steady and equal step, and should avoid a springy gait, planting the foot firmly on the ground, and not rising on the toe. The patient should be carried feet foremost and as nearly as possible on a level. The stretcher should never be carried on the shoulders.

Esmarch's
triangular
cloth.

The use of a triangular cloth introduced by Professor Esmarch for the immediate dressing of wounds or the support of fractured limbs, is as follows:—A piece of unbleached calico is cut into a triangular shape (Fig. 97), and in the following description the border from *b* to *c* is called the lower and the two others the side borders. The corner *a* is called the point, and the two others, *b* and *c*, the ends.




FIG. 97.

To apply it to the head place the lower border across the forehead, the point hanging down over the nape of the neck. Carry the two ends back above the ears, crossing them behind, and bringing them for-

Applied to
the head.



FIG. 98.

wards tie in front of the forehead. Turn the point upwards, and fasten with safety-pin (Fig. 98).

To form a sling for the arm, throw one end over the sound shoulder, bend the arm across the middle of the cloth and bring the other end up to meet its fellow, and tie. Draw the point forward round the elbow, and pin (Fig. 99).

Sling for
arm.




FIG. 99.

Applied to
shoulder.

For wounds of the shoulder the cloth must be cut in halves. One half is used to support the forearm as a sling, the other half is applied to cover the shoulder thus :—Place lower border on the middle of the outside of the upper arm, the point lying upon the side of the neck. Carry the two ends round the inside of the arm, cross, bring forward, and tie on the outer side. Slip the point under the sling, fold it back on itself, and fasten with a pin (Fig. 100).

Applied to
chest.

To apply the cloth to the chest, place the lower border across the lower part of the chest, throwing the point over one shoulder. Carry the two ends backwards and pin them together over the spine. Then draw the point down, pass it under the two




FIG. 100.

ends, fold it on itself, and pin (Fig. 101). It is obviously better to pin than tie as shown in the figure.

In injuries to the hip the whole cloth must be used. Place the middle of the lower border in front of the upper part of the thigh, the point lying upwards on the abdomen. Carry the ends backwards behind the thigh, cross, and bringing them forward pin them together. A waist-belt must be put on and the point passed under it, folded on itself, and pinned (Fig. 102).

Applied to
the hip.

In dressing the hand or foot only half a cloth is required. To apply it to the hand, place the centre

Applied to
the hand
and foot.




FIG. 101.

of the lower border across the wrist. Turn the point up over the fingers on the back of the forearm, and tie the ends over it (Fig. 103).

To apply to the foot, place the sole of the foot in the middle of the cloth, turn the point up over the toes, and passing the ends round the ankle carry them down and tie on the sole (Fig. 104).

wing for
patient in
bed.

In severe compound injuries of the lower extremities or of the spine or pelvis the swing here figured (Fig. 105) is invaluable for raising the patient from the bed with the smallest possible disturbance. It consists of two poles, which are slipped through the broad hems of the three pieces of canvas represented in the woodcut, and which are placed under the patient as he lies in bed. The ends of the poles are



FIG. 102.



FIG. 103.




FIG. 104.




FIG. 105.

then connected together and kept asunder by a light rod of iron at the upper and lower end. Two ropes, each having a loop at either end and one in the middle, are then attached to the poles by slipping the loops at either extremity over their ends. The central loops are then attached by a hook to a multiplying pulley, which in its turn is connected by ropes to a hook fixed in the ceiling or to a framework erected over the bed. By this means the patient can be readily lifted up from the bed, or can even lift himself. The swing is the invention of Mr. Thom, late House Surgeon to the Royal Albert Hospital, and is used there especially for the after-treatment of cases of excision of the hip-joint.

In treating a fracture with pasteboard, plaster-of-Paris, or gutta-percha splints, it must be remembered that it is necessary to secure the immobility of the joints above and below the fracture. For instance, in a fracture of the upper part of the humerus the shoulder and elbow must be secured, or in a fracture of the thigh the hip and knee. For pasteboard splints rough unglazed millboard, one-eighth of an inch thick, is to be used. It should not be cut, but bent over the edge of a table, and then torn, by which a soft edge is obtained. After the splint is properly shaped it should be soaked for a few minutes in boiling water. The limb to be encased must be carefully covered with cotton-wool, and the pasteboard moulded so as to fit accurately to the wool, and secured with a bandage, which should be smeared with starch a little thicker than that ordinarily used in the laundry. An interval of a quarter of an inch to an inch should intervene between the edges of the splint.

Pasteboard
splints.

The following directions, condensed from Mr. Gamgee's book on "Fractures of the Limbs," will serve

for the treatment of special fractures with the pasteboard apparatus :—

Fractures of the Humerus in the upper part.—Bend the elbow at right angles and secure it with an outer and inner pasteboard splint, extending from the tip of the fingers to half-way up the humerus. Reduce the fracture, and apply two right-angled pasteboard splints, one from the sternal end of the clavicle to the elbow, and the other from the inner edge of the scapula to the same point. Secure with a figure-of-eight starch bandage, thoroughly covering the shoulder, and passing under the opposite axilla.

Fractures of the Leg.—Three splints are required : one behind, extending from half an inch above the heel to four inches above the knee : the other two, each made with a foot-piece, which is moulded round under the sole of the foot, are placed, the one on the outer, and the other on the inner side of the limb, and carried up to four inches above the knee. During the application of the starch bandage, extension must be kept up by two assistants, one at the foot and the other at or above the knee.

Fractures of the Femur.—Extension being kept up by two assistants, one at the foot and the other with a jack-towel round the perinæum, three splints are applied : an inner one from a little below the knee to the ramus of the ischium, an outer one from the same point to the crest of the ilium, and a posterior one also from the same point below to the crest of the ilium behind. The lateral splints must be of uniform width, whilst the posterior one should increase in breadth at the upper part so as to cover the whole buttock. To facilitate the application of the splints and bandages, the patient should be raised a little by passing a jack-towel under the loins, two assistants by this means lifting the pelvis from the bed. A figure-of-eight starch bandage is then to be applied

round the pelvis and upper part of the thigh, taking care so to spread the bandage out behind as to cover the entire buttock. The pelvis and thigh being thus encased, the apparatus described above for fracture of the leg is to be applied below and made continuous with the splints and bandage above. It is advisable to keep up extension during the drying process, by fixing a perineal band to the head of the bed whilst extension is kept up from the foot by means of a weight.

It is necessary in all these cases, after the lapse of a day or so, to slit up the bandage with cutting pliers, and pare off the edges, in order to bring the apparatus in close contact with the limb, which, owing to the subsidence of swelling, will probably have receded from the splints. A good pair of pliers, made by Coxeter for the purpose, are here figured (Fig. 106).




FIG. 106.

Fractures of the Fingers may be treated with paper and gum. Ordinary writing-paper, cut in strips long enough to extend from the tip of the finger to a couple of inches above the head of the metacarpal bone, should be placed in front and

behind, and eight or ten layers, well gummed together, will form a sufficient splint, which must be secured by a narrow bandage. The finger should first be cased in cotton-wool.

Plaster-of-Paris.

The plaster-of-Paris bandage is best applied as follows:—Pour a small quantity of water into a basin, and sprinkle into it perfectly fresh and dry plaster-of-Paris until a thin paste is formed. Take an ordinary unbleached calico bandage of the required breadth, previously saturated with water, and unroll and re-roll it under the paste, so as to impregnate the whole bandage with the plaster paste. The bandage must then be rapidly applied to the limb, which should be first covered with a flannel bandage. If possible no turns should be made in the bandage, and some of the paste will be laid on with the hand between the layers. If the plaster is fresh and dry, the bandage will set in five minutes, when it may be painted over with white of egg, to prevent its crumbling. The addition of a little gum to the plaster-of-Paris renders the bandage less brittle when dry. The following is a good formula:—Gum mucilage (B.P.) $\mathfrak{z}\text{j}$, water $\mathfrak{z}\text{viii}$, plaster-of-Paris lbj . The plaster-of-Paris is sprinkled into the mixture of gum and water.

Gutta-percha splints.

Gutta-percha splints are made, as a rule, about one-eighth of an inch in thickness. The gutta-percha is obtained in sheets, and the best method for cutting out the splint is to trace the outline with the point of a knife. Then bending the gutta-percha, and drawing the point of the knife successively along the original line, the gutta-percha will give before it, and a clean cut will be effected in the exact direction required. It is then to be placed flat in a tub of boiling water, and retained long enough to make it quite soft. The limb must be well oiled, and the gutta-percha being moulded to it should be retained

in place for a few minutes by a bandage. When it is sufficiently hard it may be removed, the edges pared, the inner edge being bevelled off, and the inside covered with flannel stuck on with paste. The splint must then be well perforated with small holes made with a punch. If the splint requires strengthening at any point, this may be accomplished by putting on to the parts requiring extra support strips of gutta-percha softened in boiling water.

A very excellent and stiff bandage is made with gum and starch, of each ℥iv , chalk ℥viii , water ℥viii . These ingredients are mixed well in a mortar and run through a hair-sieve, to make the mixture perfectly smooth. The addition of methylated spirits ℥iv prevents the mixture spoiling, and it can be kept in a bottle ready for use. The bandage should be rolled in the mixture immediately before application. Its removal is very easily effected by immersing the limb in hot water for a short time, which softens the bandage, when it can be simply removed by un-rolling.

Gum, starch,
and chalk.

Glue is sometimes used instead of starch. A strong solution of the best French glue is made, to which is added one-eighth part of methylated spirits of wine. The limb being cased in wool is then bandaged, the glue being painted on with a stiff brush. Successive layers of bandage and glue may thus be applied.

Glue.

The method of applying extension to the lower extremities by weight is as follows:—

Extension
by weight.

A thin piece of board, from three to four inches long and two and a half broad, is used as a foot-piece, to prevent the strips of plaster pressing on the malleoli. This foot-piece is bored in the centre with a hole, through which is passed the cord attached to the weight (Fig. 107). The plaster is to be cut in the shape shown in Fig. 108, and of the following dimen-

sions:—Five and a half inches wide in the centre,




FIG. 107.

and two and a half inches wide at the narrowest points, gradually widening again to four inches at the extremities. The two ends are slit down for about two-thirds of their length, as shown in the figures. The total length of the strip for an adult is about four feet eight inches, sixteen inches for the central portion, and twenty inches for each tail. The foot-piece is laid on the central portion of the strip, and the edges are folded over it (Fig. 109). The plaster is applied to the leg, one tail on the outer side, and the other on the inner side of the limb, the foot-piece being kept about two inches below the sole of the foot. The strip is further




FIG. 108. FIG. 109.

secured to the limb by a bandage, which should be applied from the toes upwards, as shown in Fig. 110. It is advisable to protect the tendo Achillis and the instep with cotton-wool. The amount of extension required for an adult is about eighteen or twenty pounds, and this may be applied at once. A pillow placed under the




FIG. 110.

popliteal space, raising the knee, relieves the pain produced by the dragging on the hamstring tendons. In very muscular patients it is sometimes advisable to apply the extension when the patient is under the influence of chloroform, as the muscles are then perfectly relaxed. For children, the rule as to weight is to add a pound for each year up to twenty.

The use of the aspirator has been alluded to in the previous pages. The following description of the aspirator, and its method of use, is published by Messrs. Weiss, whose aspirator is here figured (Fig. 111):—

The aspirator.

The aspirator consists of—

1. A glass bottle or reservoir, A, mounted with a two-way stop-cock, B, and having an opening at the bottom for the insertion of the tube, C.
2. An exhausting syringe with elastic connecting tube, D H.
3. A series of tubular needles to be attached to the reservoir by an elastic tube, as shown at E F.

Directions for Use.—Adjust the aspirator as figured in the diagram with the stop-cock B turned vertically—that is, open to the bottle; close the stop-cock in the tube C, and form a vacuum by a few upward and downward movements of the piston of the exhausting syringe D.




FIG. 111.

Insert one of the needles beyond the two eyes, attach tube F to it, turn the stop-cock B towards the needle—namely, horizontally, and continue the insertion of the needle until fluid is seen to flow through the short glass tube G into the reservoir.

To empty the latter turn stop-cock B vertically, detach the syringe tube, and open the stop-cock in tube C.

The presence of fluid having been established by the use of one of the fine needles, it is recommended for more quickly emptying the cyst to use one of the larger needles or trocars.

The introduction of the needle into the tissues requires some precautions. In place of endeavouring to penetrate by pressure, as with an ordinary trocar, it is preferable to combine pressure with rotation, by taking the needle in the forefinger and thumb, and rolling it between them. Such a manœuvre is rendered necessary by the extreme fineness of the needle, which would be liable to bend or twist if driven in by direct pressure. Before using a needle it is well to be assured of its permeability.

In cases requiring transfusion of blood, the best method is that described by Dr. Aveling. His instrument, made by Messrs. Krohne and Sesemann, is here figured (Fig. 112), and the mode of operating is thus described :—

Transfusion.

First place the apparatus in a basin of tepid water, and while completely under the water, to fill it and insure its cleanliness, compress the bulb until all the air is expelled.

The patient having been brought to the side of the bed, and the arm bared, a fold of skin over a vein at the bend of the arm should be raised, transfixed, and divided. The flattened vein now brought into view should be seized with a pair of fine forceps, raised while an incision is made in it, and the bevel-pointed

silver tube inserted. In taking this tube out of the




FIG. 112.

basin, it should be kept full of water, by placing the

tip of the thumb over its larger opening. While the operator is doing this, an assistant should prepare the arm of the blood-donor, as in ordinary bleeding, making an incision directly into the vein, and passing the round-pointed tube into it, with its point towards the fingers. This person should then be brought to the bedside of the patient, and seated in a chair. It is better not to secure the tubes in the veins by ligatures. B represents the hand of an assistant holding the efferent tube and the lips of the small wound together, and A shows the afferent tube secured in the same manner. The india-rubber portion of the apparatus, filled with water, and kept so by turning the cock at each end of it, is now fitted into the two tubes. The cocks are then turned straight, and the operation commenced by compressing the india-rubber tube on the efferent side D, and squeezing the bulb C; this forces two drachms of water into the afferent vein. Next shift the hand D to D' and compress the tube on the afferent side, then allow the bulb to expand slowly, when blood will be drawn into it from the efferent vein. By repeating this process, any quantity of blood can, at any rate, be transmitted, the amount being measured by counting the number of times the bulb is emptied.

The method of bloodless operations introduced by Esmarch may be carried out by two methods. One consists of an elastic bandage, which is applied firmly from the extremity of the limb to be operated on to a point above the site of the operation. A thick india-rubber cord is then wound very tightly round the limb just above the edge of the elastic bandage, which is then removed. The limb below the india-rubber cord will then be found perfectly blanched and empty of blood. A second method is described by Mr. Cripps in the *Lancet* of October 11th, 1873:—

Esmarch's
bloodless
operation.

"The following is a simple modification of Es-march's arrangement, by which many yards of elastic bandage may be dispensed with, and it can be easily and quickly applied.

"To apply this to the arm, three or four complete turns of the elastic ring (Fig. 113) are wound tightly




FIG. 113.

round the hand in such a manner as to include the fingers and thumb, care being taken that the turns lie even and do not cross one another. The reel is then put up under the free portion of the ring connecting the upper and lower coil. The reel is passed round and round the limb in an upward direction: thus each coil is unwound from below as another is added above. In this way four tight coils of india-rubber are carried up the limb to any distance required. The degree of tightness can be regulated with the greatest nicety by the distance the reel is drawn from the limb by the bandager.

"This method of driving blood from the limb answers perfectly in the arm and in the lower part of the leg; but in carrying the bandage over the popliteal space the flexor tendons prevent the artery

being effectually compressed. A firm pad in the space would probably answer the purpose.

"It may be unrolled by reversing the action of the wheel to remove the bandage."

As a substitute for the usual elastic cord the following very admirable and simple instrument may be had recourse to, or it may be used simply as a

The elastic
tourniquet.




FIG. 114.

tourniquet. It consists of a thin piece of elastic tubing and of a boxwood clip. To apply it, embrace the limb twice with the cord, and whilst the cord is in a state of tension slip both ends into the groove in the clip (Fig. 114). In relaxing the tension they become firmly fixed. To remove the tourniquet stretch the ends and lift them out of the groove. Fig. 115 represents the tourniquet in position with a roller bandage slipped under the clip and resting over the artery. This tourniquet is made by Mr. Coxeter, of Grafton Street, London.




FIG. 115.

Hunter's
tourniquet.

A very simple and efficient tourniquet, invented by the late Dr. Hunter, R.N., is here figured (Figs. 116, 117). It consists of a piece of beech-wood shaped like




FIG. 116.

the letter U, $2\frac{1}{4}$ inches long by $1\frac{3}{4}$ inches broad, which narrows down to form an oval-shaped pad, $1\frac{1}{2}$ inches




FIG. 117.

by 1 inch, with a slightly concave surface covered with chamois leather. Two slits half an inch wide and half an inch from one another are cut through the top, in a direction outwards and downwards, terminating on either side at a point opposite the centre of the pad. Beneath one slit a piece of webbing four inches long and one inch broad

is nailed, a buckle being attached to the end. Beneath the other slit another piece of webbing is nailed, eighteen inches long. A piece of "French sennit,"¹ eight inches long, is then passed through the slits, which it must fit loosely, and sewn by either extremity to the upper surface of the webbing. The centre of the sennit is firmly sewn round a bit of tin tubing, in which runs a three-inch piece of iron wire, No. 9 gauge, with a knob at each end to prevent its slipping out of the tube. To apply this tourniquet, buckle the webbing firmly round the limb, the pad being placed over the artery. Then, with the bit of iron acting as a lever, twist the sennit round and round until the required pressure is obtained, fixing the iron by slipping it through the small ring screwed into the top of the wooden pad. Fig. 116 shows the tourniquet unfitted, and Fig. 117 shows it buckled, with the sennit on the stretch. The webbing on the second figure appears twisted, but it is only so drawn to illustrate the fact that the whole strain of the pressure is on the sennit. As a matter of fact the webbing lies quite flat upon the limb.

In opening a deep-seated abscess in the neighbourhood of important parts the best plan is to make an incision through the skin only, and then thrust in a director until the cavity of the abscess is reached, which will be known by the pus welling up along the groove of the director. Then pass the blades of a pair of dressing-forceps along the director, and on opening them a free exit will be given to the matter. A drainage tube may be then inserted. To make a counter-opening in dependent parts is frequently a difficult and dangerous proceeding. I have found great help from the use of a rectum trocar and canula. Pass the canula, with the trocar withdrawn within

Opening
deep-seated
abscesses.

Counter-
opening.

¹ French sennit is made by plaiting about nine strands of soft thickish twine into a flat band, half an inch in width.

it, down to the point where the counter-opening is desired ; make it project underneath the skin, and then rapidly push the trocar through ; withdraw the trocar, and pass a drainage tube through the canula, or, if desirable, enlarge the counter-opening by passing a blunt-pointed bistoury down by the side of the projecting end of the canula. (To open an abscess by the antiseptic method, see p. 178.)

Dressing for
burns and
scalds.

By far the best local application for burns is the Collodion Flexile of the Pharmacopœia. This should be painted on smoothly with a large brush. It will frequently prevent vesication, if it has not already taken place. If there are vesications the serum should be let out through small openings, and the surface painted over with the collodion. A solution of soda applied to the injured surface immediately abates pain. In extensive burns, which are generally accompanied by considerable shock, the heat of the body must be maintained, and stimulants freely given, with opiates to relieve pain.

INDEX.

	PAGE
ABDOMEN, injuries to	83
Abscesses, opening, under carbolic acid	176
" " deep-seated	207
Accidental hæmorrhage in parturition	148
Acetabulum, fracture of	96
Acids, injury of the eye by	27
" " larynx by	35
Aconitum Napellus, poisoning by	167
Acute glaucoma	28
Adams on fracture of nasal bone	11
Aitkin's Crimean arm splints	59
Alcohol, poisoning by	164
Amputation of arm	68
" " at elbow-joint	79
" " fingers	81
" " forearm	79
" " foot	145
" " at knee-joint	133
" " of leg	144
" " at shoulder-joint	68
" " at the hip-joint	133
" " of thigh in upper and lower third	133
" " of thumb	80
" " of toes	146
Ankle-joint, dislocation of	141
" " sprains of	112
Anterior chamber, effusion of blood into	117
" " tibial artery, ligature of	117
Anteversion of gravid uterus	155
Antiseptic atmosphere	175
" " gauze	176
" " " bandage	176
" " " treatment	171
Antrum, fractures of	13
Apparatus, extemporary	208

	PAGE
Arm and hand presentation	156
Arnold's splint	139
Arsenic, poisoning by	161
Artificial eye	30
" respiration	43
Aspirator, description of	199
" use of, in hernia	88
" " retention of urine	103
Atropa belladonna, poisoning by	168
Atropine, hypodermic injection of, in poisoning by fungi	168
" " " " " opium	163
 BANDAGE for fracture of lower jaw	15
Base of skull, fracture of	5
Bladder, foreign bodies in the female	99
" injury to	97
" puncture of, by rectum	101
" rupture of	158
Blue vitriol, poisoning by	162
Boric acid in dressings	173
Brain, compression of	2
" concussion of	1
Breech presentation	155
Burns	208
 CÆSAREAN section	151
Cantharides, poisoning by	163
Canula for tracheotomy	38
Carbolic acid	171
" dressing	173
Carbolized silk sutures	76
Carpal bones, dislocation of	77
" fracture of	75
Catarrhal ophthalmia	19
Catgut ligature	177
Catheter, india-rubber	105
" probe-pointed	104
" prostatic	104
Cephalotripsy	154
Cerebral nerves, injury to	6
Cerebro-spinal fluid, discharge of	6
Chloride of zinc dressing	174
" " poisoning by	162
Chloroform, death from	165
Choroid, injuries to	27
Clavicle, dislocation of	54
" fracture of	52
Clove-hitch	78
Cock's operation for retention of urine	103

INDEX.

211

	PAGE
Coins in the throat	33
Colles' fracture of radius	72
Common carotid, ligature of	41
Compound fractures	138
" " treatment by carbolic acid	181
" " of the leg	136
Compression of the brain	2
Concussion " "	1
" of the spine	17
Conjunctiva, acute diseases of	19
" effusion of blood under	19
" lacerations of	19
Contusions of lung	4
" lower extremities	111
Conveyance of injured persons	186
Cord, prolapse of	159
Cornea, abrasions of	22
" acute diseases of	23
" crescentic ulceration of	24
" foreign bodies in	20
" " instrument for removal of	21
" paracentesis of	23
" sloughing ulcer of	24
" wounds of	22
Coronoid process, fracture of	69
Corrosive sublimate, poisoning by	162
Counter-openings	207
Craniotomy	152
" forceps	153
Cut-throat	40
Cutting pliers for starch bandages	195
Cyanide of potassium, poisoning by	164
 DALBY's carminative, poisoning by	 163
Decapitation of fœtus	153
Detachment of iris	25
Diffuse suppurative corneitis	23
Dilating bags	151
" female urethra	99
Dislocation of ankle-joint "	141
" clavicle	54
" femur on to the pubes	128
" " on dorsum ilii	123
" " into foramen ovale	127
" " " sciatic notch	124
" " reduction of, by extension	130
" " " manipulation	125
" " humerus, sub-coracoid	62
" " " sub-glenoid	62

	PAGE
Dislocation of humerus, sub-spinous	63
" " reduction of, by extension †	64
" " " manipulation	64
" lens	26
" lower jaw	16
" metatarsal bones	144
" patella	135
" pubis	95
" radius	74
" " and ulna	75
" scapula	55
" semilunar cartilages	141
" spine	17
" tarsal bones	145
" carpal bones	77
" thumb	77
" tibia	141
Displacements of the gravid uterus	155
Drainage tube	176
Drowning	42
Dupuytren's splint	136
EARS, foreign bodies in	7
" forceps for	9
Elbow-joint, amputation at	79
" compound injuries to	78
" excision of	78
" injuries to	68
Elevator	5
Embryulcia	153
Emphysema	49
Epistaxis	8
Ergotine, subcutaneous injection of	11
Esmarch's cloth	186
" bloodless operations	203
Essential oil of bitter almonds, poisoning by	164
Extemporary apparatus	185
Extension by weight	197
Extirpation of eyeball	29
Eye, artificial	30
" extirpation of	29
" bandage	23
" foreign bodies within	28
" " under lid	29
" injuries to, from chemicals, etc.	27
" " gunpowder	28
" " strong acids	27
" needles for operations on the	21
" shade for	26

	PAGE
Eye, speculum	29
„ lids, wounds of	6
FACE, injuries to	6
„ presentation	155
Female bladder, foreign bodies in	99
Femoral hernia	92
„ artery, application of tourniquet to	114
„ „ digital pressure on	114
„ „ ligature of	115
„ „ wounds of	114
Fingers, amputation of	81
Fish bones in the throat	32
Flooding	150
Fœtus, decapitation of	153
Foot, amputations of	145
Forceps for the ear	9
Foreign bodies in œsophagus	33
„ cornea	20
„ ear	7
„ under eyelid	22
„ in the larynx	34
„ „ nose	7
„ „ pharynx	32
„ „ rectum	100
„ „ trachea	34
„ „ urethra	98
Fracture of acetabulum	96
„ antrum	13
„ base of skull	5
„ carpal and metacarpal bones	75
„ clavicle	52
„ coronoid process of ulna	69
„ at elbow-joint	60
„ the femur, neck of	117
„ „ table of diagnosis in	124
„ „ shaft of	121
„ „ „ in children	122
„ fibula	136
„ humerus, neck of	58
„ „ shaft of	59
„ „ lower extremity of	60
„ ischium	95
„ into the knee-joint	123
„ lower jaw	14
„ „ treatment of	14
„ metatarsal bones and phalanges	144
„ nasal bones	11

	PAGE
Fracture of olecranon	69
" patella	134
" pelvis	95
" pubis	95
" radius, neck of	71
" " shaft of	71
" " and ulna	74
" ribs	48
" sacrum	97
" scapula	55
" skull	2
" " vault of	2
" spine	17
" tarsal bones	143
" tibia	136
" " and fibula	137
" ulna, shaft of	70
" upper jaw	12
" zygoma	14
Fungi, poisoning by	168
GASES, respiration of noxious	45
Gastrocnemius, rupture of	112
Gauze bandage	176
Glaucoma, traumatic	26
" acute	28
Glue bandages	197
Godfrey's cordial, poisoning by	163
Gonorrhœal ophthalmia	20
Gordon's splint for Colles' fracture	72
Gum, chalk, and starch bandages	197
Gunpowder, injuries of the eye from	28
Gunshot wound of chest	50
Gutta-percha splints	196
HEMOTHORAX	49
Hamilton's tables of injuries to the shoulder	56
Hands and feet presentation	156
" head "	157
Hanging	45
Hartshorn and oil, poisoning by	161
Hemlock, poisoning by	164
Hernia, the aspirator in	48
" femoral	92
" inflamed	87
" inguinal	91
" obstructed	87
" obturator	93

INDEX.

215

	PAGE
Hernia, operation for	89
„ strangulated	87
„ the taxis in	88
„ umbilical	91
„ of lung	48
Hey's saw	5
Hip-joint, contusion of	111
„ amputation at	133
Horsehair probang	33
Humerus, dislocation of	58
„ fracture of	59
Hypopion	23
INDUCTION of premature labour	151
Inguinal hernia	91
Injuries to cerebral nerves	6
„ optic „	31
Instrument for plugging nose	8
„ removing foreign bodies from cornea	21
Intestines, constriction of	85
„ foreign bodies in	86
„ gangrene of	85
„ protrusion of	85
„ wounds of	85
Iris, acute diseases of	25
„ injury to	24
„ detachment of	25
„ prolapse of	25
Iritis, rheumatic	25
„ syphilitic	25
„ traumatic	25
Iron in cornea	25
JUGULAR vein, venesection from	44
KNEE-JOINT, amputation at	133
„ compound injuries to	131
„ excision of	132
„ fractures into	123
„ needle in	114
„ wounds of	113
LARYNGITIS	35
Laryngotomy	35, 36
Laryngo-tracheotomy	37
Larynx, foreign bodies in	35
„ injury by acids	35
„ scalds of	34

	PAGE
Leg, amputation of the	144
Lens, dislocation of	26
„ injury to	26
„ partial dislocation of	27
Liebreich's eye bandage	22
Ligamentum patellæ, rupture of	112
Lime, injury of the eye by	27
Lips, wounds of	6
Lower jaw, dislocations of	16
„ fractures of	14
Lung, contusion of	47
„ hernia of	48
„ wound of	48
MACINTYRE's splint	123
Mayer's instrument for plugging nares	10
McNalty's splint for fracture of leg	9
Malar bone, fracture of	13
„ „ depression of	13
Mal-presentations	155
Metacarpal bones, fracture of	75
Money probang	33
Muscles, rupture of abdominal	83
„ „ of lower extremity	111
Mussels, poisoning by	163
NASAL bones, fracture of	11
Needle-holder	6
„ in the hand	52
Nerves, injury to	6
Nitric acid, poisoning by	168
Nose, bleeding from	9
„ foreign bodies in	7
„ plugging the	8
OBTURATOR hernia	93
Œsophagus, foreign bodies in	33
Olecranon, fracture of	69
Omentum, protrusion of	86
Ophthalmia, catarrhal	19
„ purulent	20
„ gonorrhœal	20
Opium, poisoning by	163
Optic nerve, injury to	31
Orbit, injuries to	30
Orpiment, poisoning by	162
Oxalic acid, poisoning by	160
PALMAR arch, wounds of	51

	PAGE
Paracentesis thoracis	50
" corneæ	28
Paraphimosis	108
Paregoric elixir, poisoning by	164
Parturition, accidental hæmorrhage in	148
" emergencies of	147
Pasteboard splints	193
Patella, dislocation of	135
" fracture of	134
Pelvis, fracture of bones of the	95
Penis, injuries to	107
Perforator	153
Perinæum, rupture of	159
Perineal band	122
" catheter	159
Peritoneum, rupture of the	83
Pharynx, foreign bodies in	32
Phosphorus, poisoning by	141
Phimosis	108
Pins in the throat	33
Placenta prævia	149
" retained	149
Plantaris, rupture of	112
Plaster-of-Paris bandage	196
Platinum wire suture in wounds of sclerotic	22
Plugging posterior nares	10
Pneumothorax	49
Poisoning by alcohol	164
" arsenic	161
" atropine	168
" blue vitriol	162
" cantharides	163
" chloroform	164
" chloride of zinc	162
" corrosive sublimate	162
" cyanide of potassium	164
" Dalby's carminative	163
" deadly nightshade	168
" essential oil of bitter almonds	164
" fungi	168
" Godfrey's cordial	163
" hartshorn and oil	161
" hemlock	164
" monkshood	167
" mussels	163
" nitric acid	160
" opium	163
" orpiment	162
" oxalic acid	160

	PAGE
Poisoning by paregoric elixir	164
" phosphorus	161
" prussic acid	164
" salt of sorrel	160
" Scheele's green	161
" strychnia	164
" sugar of lead	162
" sulphuric acid	160
" tartar emetic	162
" tobacco	164
Posterior tibial artery, ligature of	116
Pregnancy, signs of	147
Premature labour, induction of	151
Prolapse of cord	157
" iris	25
Prostate, enlarged	104
Protective oil-silk	175
Prussic acid, poisoning by	164
Pubis, dislocation and fracture of	95
Puerperal convulsions	157
Puncture of bladder by rectum	101
Punctured wounds of scalp	1
Purulent ophthalmia	20
Pus in anterior chamber	20
 RADIUS, dislocation of	 74
" fracture of the neck of	75
" " " shaft of	75
Rectum, foreign bodies in	109
" hæmorrhage from	109
Respiration, artificial	43
" of noxious gases	45
Retained placenta	149
Retention of urine	100
" " probe-pointed catheter for	100
" " from enlarged prostate	104
Retroversion of gravid uterus	155
Ribs, fracture of	46
Rupture of muscles	83
" peritoneum	83
 SACRUM, fracture of	 97
Salt of sorrel, poisoning by	160
Saw for resection of shoulder-joint	68
Scalds	208
Scapula, dislocations of	55
" fractures of	55
Scheele's green, poisoning by	161
Scissors for extirpation of eyeball	29

INDEX.

219

	PAGE
Sclerotic, wounds of	22
Scrotum, injuries of	108
Semilunar cartilages, dislocation of	141
Shade for eye	26
Shoulder-joint, amputation at	68
" " compound injuries to	65
" " injuries to	55
" " resection of	67
" presentation	156
Signs of pregnancy	147
Skull, fracture of	2
Solution of ergotine for subcutaneous injection	11
Speculum for eye	29
Spine, injury to	16
Splint for fractures of the neck of humerus	59
" interdental	15
" pasteboard	193
Sprains of ankle-joint	15
" spine	16
Steam-spray apparatus	173
Stomach-pump, use of	168
Strapping in fractured ribs	46
Strychnia, poisoning by	164
Sugar of lead, poisoning by	162
Sulphuric acid, poisoning by	160
Swing for raising patients in bed	190
Sylvester's method of artificial respiration	43
Syphon tube	7
 TARSAL bones, dislocation of	 143
" " fracture of	143
Tartar emetic, poisoning by	162
Tendo Achillis, rupture of	112
Testicle, injuries to	108
Thigh, amputation of	133
Throat, stabs in	41
Thumb, amputation of	80
" dislocation of	77
Tibia, fracture of	136
Tobacco, poisoning by	164
Toes, amputation of	146
Tongue, wounds of	32
Tosswell's stomach-pump	169
Tourniquet, elastic	205
" Hunter's	206
" temporary	206
Trachea, foreign bodies in	204
Tracheotomy	37
Transfusion	201

	PAGE
Trephining the skull	3
Turning	157
ULNA, dislocation of	75
,, fracture of the shaft of	74
Umbilical hernia	91
Upper jaw, fracture of	12
Urethra, forceps for	98
,, foreign bodies in	98
,, injuries to	99
,, wounds of	99
Urine, extravasation of	107
,, retention of	100
Uterus, displacements of gravid	155
,, injuries to pregnant	109
,, inversion of	159
,, rupture of	158
VAGINA, injuries to	109
,, plugging the	149
Vas deferens, rupture of	109
Venesection from jugular vein	44
Viscera, injury to abdominal	84
,, protrusion of	85
WOUNDS of abdominal parietes	84
,, chest	48
,, cornea	22
,, eyelids	6
,, face	6
,, heart	50
,, intercostal arteries	47
,, intestines	85
,, knee-joint	113
,, lips	71
,, lungs	48
,, palmar arch	51
,, scalp	1
,, sclerotic	22
,, tongue	32
,, urethra	99
,, near wrist-joint	51
Wrist-joint, compound injury to	80
ZYGOMA, fracture of	14



LONDON, *January*, 1880.

J. & A. CHURCHILL'S
MEDICAL CLASS BOOKS.

ANATOMY.

BRAUNE.—An Atlas of Topographical Anatomy, after Plane Sections of Frozen Bodies. By WILHELM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., Surgeon to Charing Cross Hospital, and Lecturer on Anatomy in its School. With 34 Photo-lithographic Plates and 46 Woodcuts. Large imp. 8vo, 40s.

FLOWER.—Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S., Conservator of the Museum of the Royal College of Surgeons. Second Edition, containing 6 Plates. Royal 4to, 12s.

GODLEE.—An Atlas of Human Anatomy: illustrating most of the ordinary Dissections; and many not usually practised by the Student. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant-Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. To be completed in 12 Parts, each containing 4 Coloured Plates, with Explanatory Text. Parts I to X. Imp. 4to, 7s. 6d. each.

HEATH.—Practical Anatomy: a Manual of Dissections. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Fourth Edition. With 16 Coloured Plates and 264 Engravings. Crown 8vo, 14s.

NEW BURLINGTON STREET.

ANATOMY—*continued.*

HOLDEN.—**Human Osteology:** comprising a

Description of the Bones, with Delineations of the Attachments of the Muscles, the General and Microscopical Structure of Bone and its Development. By LUTHER HOLDEN, F.R.C.S., Senior Surgeon to St. Bartholomew's and the Foundling Hospitals, and ALBAN DORAN, F.R.C.S., late Anatomical, now Pathological, Assistant to the Museum of the Royal College of Surgeons. Fifth Edition. With 61 Lithographic Plates and 89 Engravings. Royal 8vo, 16s.

By the same Author.

A Manual of the Dissection of the

Human Body. Fourth Edition. Revised by the Author and JOHN LANGTON, F.R.C.S., Assistant Surgeon and Lecturer on Anatomy at St. Bartholomew's Hospital. With Engravings. 8vo, 16s.

ALSO,

Landmarks, Medical and Surgical. Second

Edition. 8vo, 3s. 6d.

MORRIS.—**The Anatomy of the Joints of Man.**

By HENRY MORRIS, M.A., F.R.C.S., Surgeon to, and Lecturer on Anatomy and Practical Surgery at, the Middlesex Hospital. With 44 Plates (19 Coloured) and Engravings. 8vo, 16s.

WAGSTAFFE.—**The Student's Guide to Human**

Osteology. By WM. WARWICK WAGSTAFFE, F.R.C.S., Assistant-Surgeon to, and Lecturer on Anatomy at, St. Thomas's Hospital. With 23 Plates and 66 Engravings. Fcap. 8vo, 10s. 6d.

WILSON.—**The Anatomist's Vade-Mecum: a**

System of Human Anatomy. By ERASMUS WILSON, F.R.C.S., F.R.S., late Professor of Dermatology to the Royal College of Surgeons. Ninth Edition, by G. BUCHANAN, M.A., M.D., late Professor of Clinical Surgery in the University of Glasgow, and HENRY E. CLARK, F.F.P.S., Lecturer on Anatomy in the Glasgow Royal Infirmary School of Medicine. With 371 Engravings. Crown 8vo, 14s.

Anatomical Remembrancer (the); or, Complete Pocket Anatomist. Eighth Edition. 32mo, 3s. 6d.

NEW BURLINGTON STREET.

BOTANY.

BENTLEY.—*A Manual of Botany.* By ROBERT BENTLEY, F.L.S., Professor of Botany in King's College and to the Pharmaceutical Society. With 1138 Engravings. Third Edition. Crown 8vo, 14s.

BENTLEY AND TRIMEN.—*Medicinal Plants:* being descriptions, with original Figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By ROBERT BENTLEY, F.L.S., and HENRY TRIMEN, M.B., F.L.S., British Museum, and Lecturer on Botany at St. Mary's Hospital Medical School. In 42 Monthly Parts, each containing 8 Coloured Plates. Large 8vo, 5s. each part.

CHEMISTRY.

BERNAYS.—*Notes for Students in Chemistry;* being a Syllabus of Chemistry compiled mainly from the Manuals of Fownes-Watts, Miller, Wurz, and Schorlemmer. By ALBERT J. BERNAYS, Ph.D., Professor of Chemistry at St. Thomas's Hospital, Examiner in Chemistry at the Royal College of Physicians of London. Sixth Edition. Fcap. 8vo, 3s. 6d.

By the same Author.

Skeleton Notes on Analytical Chemistry,
for Students in Medicine. Fcap. 8vo, 2s. 6d.

BLOXAM.—*Chemistry, Inorganic and Organic;* with Experiments. By CHARLES L. BLOXAM, Professor of Chemistry in King's College. Fourth Edition. With nearly 300 Engravings. 8vo, 16s.

By the same Author.

Laboratory Teaching; or, Progressive
Exercises in Practical Chemistry. Fourth Edition. With 83 Engravings. Crown 8vo, 5s. 6d.

BOWMAN AND BLOXAM.—*Practical Chemistry,* including Analysis. By JOHN E. BOWMAN, Formerly Professor of Practical Chemistry in King's College, and CHARLES L. BLOXAM, Professor of Chemistry in King's College. With 93 Engravings. Seventh Edition. Fcap. 8vo, 6s. 6d.

NEW BURLINGTON STREET.

CHEMISTRY—*continued.*

CLOWES.—**Practical Chemistry and Qualitative Inorganic Analysis.** An Elementary Treatise specially adapted for use in the Laboratories of Schools and Colleges, and by Beginners. By FRANK CLOWES, D.SC., Senior Science Master at the High School, Newcastle-under-Lyme. Second Edition. With 47 Engravings. Post 8vo, 7s. 6d.

FOWNES AND WATTS.—**Physical and Inorganic Chemistry.** Twelfth Edition. By GEORGE FOWNES, F.R.S., and HENRY WATTS, B.A., F.R.S. With 154 Engravings, and Coloured Plate of Spectra. Crown 8vo, 8s. 6d.

By the same Authors.

Chemistry of Carbon - Compounds, or Organic Chemistry. Twelfth Edition. With Engravings. Crown 8vo, 10s.

GALLOWAY.—**A Manual of Qualitative Analysis.** By ROBERT GALLOWAY, Professor of Applied Chemistry in the Royal College of Science for Ireland. Fifth Edition. With Engravings. Post 8vo, 8s. 6d.

VACHER.—**A Primer of Chemistry, including Analysis.** By ARTHUR VACHER. 18mo, 1s.

VALENTIN.—**Introduction to Inorganic Chemistry.** By WILLIAM G. VALENTIN, F.C.S., Late Principal Demonstrator of Practical Chemistry in the Science Training Schools. Third Edition. With 82 Engravings. 8vo, 6s. 6d.

By the same Author.

A Course of Qualitative Chemical Analysis. Fourth Edition. With 19 Engravings. 8vo, 7s. 6d.

ALSO,

Chemical Tables for the Lecture-room and Laboratory. In Five large Sheets, 5s. 6d.

NEW BURLINGTON STREET.

CHILDREN, DISEASES OF.

ELLIS.—A Practical Manual of the Diseases of Children. By EDWARD ELLIS, M.D., late Senior Physician to the Victoria Hospital for Sick Children. With a Formulary. Third Edition. Crown 8vo, 7s. 6d.

SMITH.—Clinical Studies of Disease in Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to H.M. the King of the Belgians, and to the East London Hospital for Children. Post 8vo, 7s. 6d.

By the same Author.

On the Wasting Diseases of Infants and Children. Third Edition. Post 8vo, 8s. 6d.

STEINER.—Compendium of Children's Diseases; a Handbook for Practitioners and Students. By JOHANN STEINER, M.D. Translated from the Second German Edition, by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women, &c. 8vo, 12s. 6d.

DENTISTRY.

SEWILL.—The Student's Guide to Dental Anatomy and Surgery. By HENRY E. SEWILL, M.R.C.S., L.D.S., late Dental Surgeon to the West London Hospital. With 77 Engravings. Fcap. 8vo, 5s. 6d.

SMITH.—Handbook of Dental Anatomy and Surgery. For the Use of Students and Practitioners. By JOHN SMITH, M.D., F.R.S.E., Dental Surgeon to the Royal Infirmary, Edinburgh. Second Edition. Fcap. 8vo, 4s. 6d.

STOCKEN.—Elements of Dental Materia Medica and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S., Lecturer on Dental Materia Medica and Therapeutics and Dental Surgeon to the National Dental Hospital. Second Edition. Fcap. 8vo, 6s. 6d.

DENTISTRY—*continued.*

TAFT.—**A Practical Treatise on Operative Dentistry.** By JONATHAN TAFT, D.D.S., Professor of Operative Surgery in the Ohio College of Dental Surgery. Third Edition. With 134 Engravings. 8vo, 18s.

TOMES (C. S.).—**Manual of Dental Anatomy, Human and Comparative.** By CHARLES S. TOMES, M.A., M.R.C.S., Lecturer on Anatomy and Physiology at the Dental Hospital of London. With 179 Engravings. Crown 8vo, 10s. 6d.

TOMES (J. and C. S.).—**A Manual of Dental Surgery.** By JOHN TOMES, M.R.C.S., F.R.S., Consulting Surgeon-Dentist to Middlesex Hospital; and CHARLES S. TOMES, M.A., M.R.C.S., Lecturer on Anatomy and Physiology at the Dental Hospital of London. Second Edition. With 262 Engravings. Fcap. 8vo, 14s.

EAR, DISEASES OF.

BURNETT.—**The Ear: its Anatomy, Physiology, and Diseases.** A Practical Treatise for the Use of Medical Students and Practitioners. By CHARLES H. BURNETT, M.D., Aural Surgeon to the Presbyterian Hospital, Philadelphia. With 87 Engravings. 8vo, 18s.

DALBY.—**On Diseases and Injuries of the Ear.** By WILLIAM B. DALBY, F.R.C.S., Aural Surgeon to, and Lecturer on Aural Surgery at, St. George's Hospital. With Engravings. Fcap. 8vo, 6s. 6d.

JONES.—**A Practical Treatise on Aural Surgery.** By H. MACNAUGHTON JONES, M.D., Professor of the Queen's University in Ireland, Surgeon to the Cork Ophthalmic and Aural Hospital. With 46 Engravings. Crown 8vo, 5s.

By the same Author.

Atlas of the Diseases of the Membrana Tympani. In Coloured Plates, containing 59 Figures. With Explanatory Text. Crown 4to, 21s.

NEW BURLINGTON STREET.

FORENSIC MEDICINE.

OGSTON.—Lectures on Medical Jurisprudence.

By FRANCIS OGSTON, M.D., Professor of Medical Jurisprudence and Medical Logic in the University of Aberdeen. Edited by FRANCIS OGSTON, Jun., M.D., Assistant to the Professor of Medical Jurisprudence and Lecturer on Practical Toxicology in the University of Aberdeen. With 12 Plates. 8vo, 18s.

TAYLOR.—The Principles and Practice of

Medical Jurisprudence. By ALFRED S. TAYLOR, M.D., F.R.S., Professor of Medical Jurisprudence to Guy's Hospital. Second Edition. With 189 Engravings. 2 Vols. 8vo, 31s. 6d.

By the same Author.

A Manual of Medical Jurisprudence.

Tenth Edition. With 55 Engravings. Crown 8vo, 14s.

ALSO,

On Poisons, in relation to Medical Juris-

prudence and Medicine. Third Edition. With 104 Engravings. Crown 8vo, 16s.

WOODMAN AND TIDY.—A Handy-Book of

Forensic Medicine and Toxicology. By W. BATHURST WOODMAN, M.D., F.R.C.P.; and C. MEYMOTT TIDY, M.B., Professor of Chemistry and of Medical Jurisprudence, &c., at the London Hospital. With 8 Lithographic Plates and 116 Wood Engravings. 8vo, 31s. 6d.

HYGIENE.

WILSON.—A Handbook of Hygiene and Sani-

tary Science. By GEORGE WILSON, M.A., M.D., Medical Officer of Health for Mid Warwickshire. Fourth Edition. With Engravings. Crown 8vo, 10s. 6d.

NEW BURLINGTON STREET.

HYGIENE—*continued.*

PARKES.—A Manual of Practical Hygiene.

By EDMUND A. PARKES, M.D., F.R.S. Fifth Edition by F. DE CHAUMONT, M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. With 9 Plates and 112 Engravings. 8vo, 18s.

By the same Author.

Public Health: being a Concise Sketch of the Sanitary Considerations connected with the Land, with Cities, Villages, Houses, and Individuals. Revised by WILLIAM AITKEN, M.D., F.R.S., Professor of Pathology in the Army Medical School. Crown 8vo, 2s. 6d.

MATERIA MEDICA AND THERAPEUTICS.

BINZ AND SPARKS.—The Elements of Thera-

peutics: a Clinical Guide to the Action of Medicines. By C. BINZ, M.D., Professor of Pharmacology in the University of Bonn. Translated from the Fifth German Edition, and Edited with Additions, in conformity with the British and American Pharmacopœias, by EDWARD I. SPARKS, M.A., M.B. Oxon., F.R.C.P. Lond. Crown 8vo, 8s. 6d.

ROYLE AND HARLEY.—A Manual of Materia

Medica and Therapeutics. By J. FORBES ROYLE, M.D., F.R.S., formerly Professor of Materia Medica in King's College; and JOHN HARLEY, M.D., F.R.C.P., Physician to, and Joint Lecturer on Clinical Medicine at, St. Thomas's Hospital. Sixth Edition. With 139 Engravings. Crown 8vo, 15s.

THOROWGOOD.—The Student's Guide to

Materia Medica. By JOHN C. THOROWGOOD, M.D., F.R.C.P., Lecturer on Materia Medica at the Middlesex Hospital. With Engravings. Fcap. 8vo, 6s. 6d.

WARING.—A Manual of Practical Therapeu-

tics. By EDWARD J. WARING, M.D., F.R.C.P., Retired Surgeon H.M. Indian Army. Third Edition. Fcap. 8vo, 12s. 6d.

NEW BURLINGTON STREET.

MEDICINE.

BARCLAY.—A Manual of Medical Diagnosis.

By A. WHYTE BARCLAY, M.D., F.R.C.P., Physician to, and Lecturer on Medicine at, St. George's Hospital. Third Edition. Fcap. 8vo, 10s. 6d.

BARLOW.—A Manual of the Practice of

Medicine. By HILARO BARLOW, M.D., Formerly Senior Physician to Guy's Hospital. Second Edition. Fcap. 8vo, 7s. 6d.

CHARTERIS.—The Student's Guide to the

Practice of Medicine. By MATTHEW CHARTERIS, M.D., Professor of Practice of Medicine, Anderson's College; Physician and Lecturer on Clinical Medicine, Royal Infirmary, Glasgow. With Engravings on Copper and Wood. Second Edition. Fcap. 8vo, 6s. 6d.

FENWICK.—The Student's Guide to Medical

Diagnosis. By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Fourth Edition. With 106 Engravings. Fcap. 8vo, 6s. 6d.

By the same Author.

The Student's Outlines of Medical Treatment. Fcap. 8vo, 7s.

FLINT.—Clinical Medicine : a Systematic Treat-

tise on the Diagnosis and Treatment of Disease. By AUSTIN FLINT, M.D., Professor of the Principles and Practice of Medicine, &c., in Bellevue Hospital Medical College. 8vo, 20s.

By the same Author.

A Manual of Percussion and Auscultation ;
of the Physical Diagnosis of Diseases of the Lungs and Heart, and
of Thoracic Aneurism. Post 8vo, 6s. 6d.

HALL.—Synopsis of the Diseases of the Larynx,

Lungs, and Heart : comprising Dr. Edwards' Tables on the Examination of the Chest. With Alterations and Additions. By F. DE HAVILLAND HALL, M.D., Assistant-Physician to the Westminster Hospital. Royal 8vo, 2s. 6d.

MIDWIFERY.

BARNES.—*Lectures on Obstetric Operations*, including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P. Obstetric Physician to, and Lecturer on Diseases of Women, &c., at St. George's Hospital. Third Edition. With 124 Engravings. 8vo, 18s.

CLAY.—*The Complete Handbook of Obstetric Surgery*; or, Short Rules of Practice in every Emergency, from the Simplest to the most formidable Operations connected with the Science of Obstetrics. By CHARLES CLAY, M.D., late Senior Surgeon to, and Lecturer on Midwifery at, St. Mary's Hospital, Manchester. Third Edition. With 91 Engravings. Fcap. 8vo, 6s. 6d.

RAMSBOTHAM.—*The Principles and Practice of Obstetric Medicine and Surgery*. By FRANCIS H. RAMSBOTHAM, M.D., formerly Obstetric Physician to the London Hospital. Fifth Edition. Illustrated with 120 Plates, forming one thick handsome volume. 8vo, 22s.

ROBERTS.—*The Student's Guide to the Practice of Midwifery*. By D. LLOYD ROBERTS, M.D., F.R.C.P., Physician to St. Mary's Hospital, Manchester. Second Edition. With 96 Engravings. Fcap. 8vo, 7s.

SCHROEDER.—*A Manual of Midwifery*; including the Pathology of Pregnancy and the Puerperal State. By KARL SCHROEDER, M.D., Professor of Midwifery in the University of Erlangen. Translated by CHARLES H. CARTER, M.D. With Engravings. 8vo, 12s. 6d.

SWAYNE.—*Obstetric Aphorisms for the Use of Students commencing Midwifery Practice*. By JOSEPH G. SWAYNE, M.D., Lecturer on Midwifery at the Bristol School of Medicine. Sixth Edition. With Engravings. Fcap. 8vo, 3s. 6d.

NEW BURLINGTON STREET.

MICROSCOPY.

CARPENTER.—*The Microscope and its Revelations.* By WILLIAM B. CARPENTER, C.B., M.D., F.R.S., late Registrar to the University of London. Fifth Edition. With more than 500 Engravings. Crown 8vo, 15s.

MARSH.—*Section-Cutting: a Practical Guide* to the Preparation and Mounting of Sections for the Microscope, special prominence being given to the subject of Animal Sections. By Dr. SYLVESTER MARSH. With Engravings. Fcap. 8vo, 2s. 6d.

MARTIN.—*A Manual of Microscopic Mounting.* By JOHN H. MARTIN, Member of the Society of Public Analysts, &c. Second Edition. With several Plates and 144 Engravings. 8vo, 7s. 6d.

WYTHE.—*The Microscopist: a Manual of* Microscopy and Compendium of the Microscopic Sciences, Micro-Mineralogy, Micro-Chemistry, Biology, Histology, and Pathological Histology. By J. H. WYTHE, A.M., M.D., Professor of Microscopy and Biology in the San Francisco Medical College. Third Edition. With 205 Illustrations. Royal 8vo, 18s.

OPHTHALMOLOGY.

HIGGENS.—*Hints on Ophthalmic Out-Patient Practice.* By CHARLES HIGGENS, F.R.C.S., Ophthalmic Assistant-Surgeon to, and Lecturer on Ophthalmology at, Guy's Hospital. Second Edition. Fcap. 8vo, 3s.

JONES.—*A Manual of the Principles and Practice of Ophthalmic Medicine and Surgery.* By T. WHARTON JONES, F.R.C.S., F.R.S., Ophthalmic Surgeon and Professor of Ophthalmology to University College Hospital. Third Edition. With 9 Coloured Plates and 173 Engravings. Fcap. 8vo, 12s. 6d.

MACNAMARA.—*A Manual of the Diseases of the Eye.* By CHARLES MACNAMARA, F.R.C.S., Surgeon to Westminster Hospital. Third Edition. With 7 Coloured Plates and 52 Engravings. Fcap. 8vo, 12s. 6d.

OPHTHALMOLOGY—*continued.*

NETTLESHIP.—The Student's Guide to Diseases of the Eye. By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmic Surgery at, St. Thomas's Hospital. With 48 Engravings. Fcap. 8vo, 7s. 6d.

WELLS.—A Treatise on the Diseases of the Eye. By J. SOELBERG WELLS, F.R.C.S., late Ophthalmic Surgeon to King's College Hospital, and Professor of Ophthalmology at King's College. With Coloured Plates and Engravings. Third Edition. 8vo, 25s.

PATHOLOGY.

JONES AND SIEVEKING.—A Manual of Pathological Anatomy. By C. HANDFIELD JONES, M.B., F.R.S., Physician to St. Mary's Hospital, and EDWARD H. SIEVEKING, M.D., F.R.C.P., Physician to St. Mary's Hospital. Second Edition. Edited by J. F. PAYNE, M.B., Assistant-Physician and Lecturer on General Pathology at St. Thomas's Hospital. With 195 Engravings. Crown 8vo, 16s.

VIRCHOW.—Post-Mortem Examinations: a Description and Explanation of the Method of Performing them, with especial reference to Medico-Legal Practice. By Professor RUDOLPH VIRCHOW, Berlin Charité Hospital. Second Edition, with 4 Plates. Fcap. 8vo, 3s. 6d.

WILKS AND MOXON.—Lectures on Pathological Anatomy. By SAMUEL WILKS, M.D., F.R.S., Physician to, and Lecturer on Medicine at, Guy's Hospital; and WALTER MOXON, M.D., F.R.C.P., Physician to, and Lecturer on Clinical Medicine at, Guy's Hospital. Second Edition. With 5 Steel Plates. 8vo, 18s.

PSYCHOLOGY.

BUCKNILL AND TUKE.—A Manual of Psychological Medicine: containing the Lunacy Laws, Nosology, Ætiology, Statistics, Description, Diagnosis, Pathology, and Treatment of Insanity, with an Appendix of Cases. By JOHN C. BUCKNILL, M.D., F.R.S., and D. HACK TUKE, M.D., F.R.C.P. Fourth Edition, with 12 Plates (30 Figures). 8vo, 25s.

NEW BURLINGTON STREET.

PHYSIOLOGY.

CARPENTER.—**Principles of Human Physiology.** With 3 Steel Plates and 371 Engravings. By WILLIAM B. CARPENTER, C.B., M.D., F.R.S., late Registrar to the University of London. Eighth Edition. Edited by Mr. Henry Power. 8vo, 31s. 6d.

By the same Author.

A Manual of Physiology. With upwards of 250 Illustrations. Fifth Edition. Edited by P. H. PYE-SMITH, M.D., F.R.C.P. Crown 8vo. [*In the press.*]

DALTON.—**A Treatise on Human Physiology :** designed for the use of Students and Practitioners of Medicine. By JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Sixth Edition. With 316 Engravings. Royal 8vo, 20s.

FREY.—**The Histology and Histo-Chemistry of Man.** A Treatise on the Elements of Composition and Structure of the Human Body. By HEINRICH FREY, Professor of Medicine in Zurich. Translated from the Fourth German Edition, by ARTHUR E. BARKER, Assistant-Surgeon to the University College Hospital. With 608 Engravings. 8vo, 21s.

FULTON.—**A Text-Book of Physiology, including Histology.** By J. FULTON, M.D., Professor of Physiology and Sanitary Science in Trinity Medical College, Toronto; Surgeon to the Toronto General Hospital. Second Edition, with 151 Engravings. 8vo, 15s.

RUTHERFORD.—**Outlines of Practical Histology.** By WILLIAM RUTHERFORD, M.D., F.R.S., Professor of the Institutes of Medicine in the University of Edinburgh; Examiner in Physiology in the University of London. Second Edition. With 63 Engravings. Crown 8vo (with additional leaves for Notes), 6s.

SANDERSON.—**Handbook for the Physiological Laboratory :** containing an Exposition of the fundamental facts of the Science, with explicit Directions for their demonstration. By J. BURDON SANDERSON, M.D., F.R.S., Professor and Superintendent of the Brown Institution; E. KLEIN, M.D., F.R.S., Assistant-Professor in the Brown Institution; MICHAEL FOSTER, M.D., F.R.S., Prælector of Physiology at Trinity College, Cambridge; and T. LAUDER BRUNTON, M.D., F.R.S., Lecturer on Materia Medica at St. Bartholomew's Hospital Medical College. 2 Vols., with 123 Plates. 8vo, 24s.

SURGERY.

BRYANT.—A Manual for the Practice of Surgery. By THOMAS BRYANT, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Guy's Hospital. Third Edition. With 672 Engravings (nearly all original, many being coloured). 2 vols. Crown 8vo, 28s.

BELLAMY.—The Student's Guide to Surgical Anatomy; a Text-Book for the Pass Examination. By EDWARD BELLAMY, F.R.C.S., Surgeon to, Lecturer on Anatomy and formerly Teacher of Operative Surgery in, Charing Cross Hospital. With 76 Engravings. Fcap. 8vo, 7s.

CLARK AND WAGSTAFFE.—Outlines of Surgery and Surgical Pathology. By F. LE GROS CLARK, F.R.C.S., F.R.S., Consulting Surgeon to St. Thomas's and the Great Northern Hospitals. Second Edition. Revised and expanded by the Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant-Surgeon to St. Thomas's Hospital. 8vo, 10s. 6d.

DRUITT.—The Surgeon's Vade-Mecum; a Manual of Modern Surgery. By ROBERT DRUITT, F.R.C.S. Eleventh Edition. With 369 Engravings. Fcap. 8vo, 14s.

FERGUSSON.—A System of Practical Surgery. By Sir WILLIAM FERGUSSON, Bart., F.R.C.S., F.R.S., late Surgeon and Professor of Clinical Surgery to King's College Hospital. With 463 Engravings. Fifth Edition. 8vo, 21s.

HEATH.—A Manual of Minor Surgery and Bandaging, for the use of House-Surgeons, Dressers, and Junior Practitioners. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Sixth Edition. With 115 Engravings. Fcap. 8vo. [Nearly ready.]

By the same Author.

A Course of Operative Surgery: with
Twenty Plates drawn from Nature by M. LÉVEILLÉ, and Coloured by hand under his direction. Large 8vo, 40s.

ALSO,

The Student's Guide to Surgical Diagnosis. Fcap. 8vo, 6s. 6d.

NEW BURLINGTON STREET.

SURGERY—continued.

MAUNDER.—Operative Surgery. By Charles

F. MAUNDER, F.R.C.S., late Surgeon to, and Lecturer on Surgery at, the London Hospital. Second Edition. With 164 Engravings. Post 8vo, 6s.

PIRRIE.—The Principles and Practice of

Surgery. By WILLIAM PIRRIE, F.R.S.E., Professor of Surgery in the University of Aberdeen. Third Edition. With 490 Engravings. 8vo, 28s.

TERMINOLOGY.

DUNGLISON.—Medical Lexicon: a Dictionary

of Medical Science, containing a concise Explanation of its various Subjects and Terms, with Accentuation, Etymology, Synonymes, &c. By ROBLEY DUNGLISON, M.D. New Edition, thoroughly revised by RICHARD J. DUNGLISON, M.D. Royal 8vo, 28s.

MAYNE.—A Medical Vocabulary: being an

Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their Derivation, Meaning, Application, and Pronunciation. By ROBERT G. MAYNE, M.D., LL.D., and JOHN MAYNE, M.D., L.R.C.S.E. Fourth Edition. Fcap. 8vo, 10s.

WOMEN, DISEASES OF.

BARNES.—A Clinical History of the Medical

and Surgical Diseases of Women. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Second Edition. With 151 Engravings. 8vo, 28s.

DUNCAN.—Clinical Lectures on the Diseases

of Women. By J. MATTHEWS DUNCAN, M.D., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 8s.

EMMET.—The Principles and Practice of

Gynæcology. By THOMAS ADDIS EMMET, M.D., Surgeon to the Woman's Hospital of the State of New York. With 130 Engravings. Royal 8vo, 24s.

NEW BURLINGTON STREET.

WOMEN, DISEASES OF—*continued.*

GALABIN.—The Student's Guide to the Diseases of Women. By ALFRED L. GALABIN, M.D., F.R.C.P., Assistant Obstetric Physician and Joint Lecturer on Obstetric Medicine to Guy's Hospital. With 63 Engravings. Fcap. 8vo, 7s. 6d.

SMITH.—Practical Gynæcology: a Handbook of the Diseases of Women. By HEYWOOD SMITH, M.D., Physician to the Hospital for Women, and to the British Lying-in Hospital. With Engravings. Crown 8vo, 5s. 6d.

WEST AND DUNCAN.—Lectures on the Diseases of Women. By CHARLES WEST, M.D., F.R.C.P. Fourth Edition. Revised and in part re-written by the Author, with numerous additions, by J. MATTHEWS DUNCAN, M.D., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 16s.

ZOOLOGY.

BRADLEY.—Manual of Comparative Anatomy and Physiology. By S. MESSENGER BRADLEY, F.R.C.S., Lecturer on Practical Surgery in Owen's College, Manchester. Third Edition. With 61 Engravings. Post 8vo, 6s. 6d.

CHAUVEAU AND FLEMING.—The Comparative Anatomy of the Domesticated Animals. By A. CHAUVEAU, Professor at the Lyons Veterinary School; and GEORGE FLEMING, Veterinary Surgeon, Royal Engineers. With 450 Engravings. 8vo, 31s. 6d.


HUXLEY.—Manual of the Anatomy of Invertebrated Animals. By THOMAS H. HUXLEY, LL.D., F.R.S. With 156 Engravings. Fcap. 8vo, 16s.

By the same Author.

Manual of the Anatomy of Vertebrated Animals. With 110 Engravings. Post 8vo, 12s.

WILSON.—The Student's Guide to Zoology: a Manual of the Principles of Zoological Science. By ANDREW WILSON, Lecturer on Natural History, Edinburgh. With Engravings. Fcap. 8vo, 6s. 6d.

NEW BURLINGTON STREET.





Riley Dunn & Wilson Ltd
EXPERT CONSERVATORS & BOOKBINDERS

